



IMPORTANT and PRIVATE INFORMATION ABOUT YOUR HEALTH PLEASE READ THIS VERY CAREFULLY

DIRECTIONS FOR TAKING CEFIXIME

Your sex partner has recently been treated for gonorrhea (sometimes called “the clap”). Gonorrhea is a curable sexually transmitted disease (STD) that you can get from having sex with a person who already has it. Some people with gonorrhea do not know they have it because they have no symptoms and feel fine.

-Men may notice a yellow or greenish discharge (drip) from their penis, pain or discomfort when urinating (peeing), or pain and swelling in their testicles.

-Women may notice a change in their usual vaginal discharge, pain during sex, bleeding between periods or after sex, lower belly or pelvic cramps or pain, or pain when urinating (peeing). Women can become infertile (unable to have children) if they don't get treated.

-Men and women can also get infected with gonorrhea in the throat from oral sex or in the rectum from anal sex.

Whether or not you are having any of these symptoms at this time; you could have gonorrhea. It is important that you take this medicine to prevent you from getting sick. Also, you can re-infect your partner if you have unprotected sex before you are cured of the infection.

After you take the medicine it is best for you to see your own doctor or medical care provider for an STD exam, counseling, and, possibly, to get tested for other sexually transmitted diseases. If you do not have a medical provider, or if you are uninsured, you can receive free care at the Department of Health (call the closest office to find out when you can be seen).

Before you take the medicine, please read the following:

Cefixime is very safe cephalosporin antibiotic. However, **DO NOT TAKE** if any of the following are true:

- You are female and having lower belly pain, pain during sex, vomiting, or fever.
- You are pregnant.
- You are male and having pain or swelling in the testicles (balls) or fever.

- You ever had a bad reaction, hives (a kind of rash), or allergy to penicillin or to a related antibiotic (such as ampicillin, amoxicillin, or another cephalosporin antibiotic).
- You have a serious long-term illness like kidney, heart or liver disease, or a seizure disorder.
- You are currently taking another prescription medication.

If any of these circumstances exist, you should talk to your healthcare provider as soon as possible.

Some people may get diarrhea, nausea, abdominal cramps, indigestion, rash, headache, vomiting, itching, or a vaginal yeast infection after taking cefixime. These won't last long, if they do, contact your healthcare provider.

If you experience any other side effects or an allergic reaction (rash, itching, swelling, dizziness, or trouble breathing) - you should call your healthcare provider immediately.

Take the tablet of cefixime with a glass of water or with food. It is important that you take this medicine as soon as possible to get cured.

DO NOT share or give this medicine to anyone else.

Do not have sex for the next 7 days. **It takes 7 days to be sure that the medicine has cured gonorrhea.** If you have unprotected sex during the 7 days after taking the medicine, you could still pass the infection to your sex partners. While condoms are effective, the safest way to make sure you don't pass the infection on to anyone is to not have sex for 7 days.

If you have any questions about cefixime or gonorrhea, please call the public health location closest to you:

- | | |
|--------------------------------------|-------------------------------|
| • Albuquerque: Stanford STD Clinic | Phone 505: 841-4180 |
| • Roswell: Public Health DPS | Phone 575: 347-2409 Ext. 6216 |
| • Las Cruces: Dona Ana Public Health | Phone 575: 528-5000 |
| • Farmington: San Juan Public Health | Phone 505: 327-4461 |
| • Santa Fe: Santa Fe Public Health | Phone 505: 476-2600 |

All calls are confidential. **No one will ask for your name.**