Monkeypox Screening & Testing Checklist for Healthcare Providers

PATIENT PRECAUTIONS

☐ Ha	ave suspect pat	ient wear a m	ask and cov	er any exposed	d skin lesion:	s prior to arr	ival
------	-----------------	---------------	-------------	----------------	----------------	----------------	------

- Do not place suspect patient in general waiting area, bring patient directly back to exam room
- Place suspect patient directly into a closed, single-person room. Special air handling is not required.

PATIENT SCREENING

Consider testing if the person has a rash/lesions and in the last 21 days has had:

- ☐ Direct, skin-to-skin contact with a person who has confirmed/suspected monkeypox
- Close, in-person contact with other men who have sex with men
- Traveled outside the US or in the US where there has been monkeypox noted
- Lymphadenopathy, fever, and fatigue
- Well-circumscribed, deep seated lesions that often have a central umbilication (can be in mucosal areas such as mouth and anogenital areas)

HEALTHCARE PERSONNEL PRECAUTIONS

- Prior to seeing a suspect patient, healthcare personnel should don:
 - Disposable gown, gloves, eye protection, N95 or higher-level respirator mask
 - Pregnant or immune compromised staff should avoid interaction with suspect patients
- ☐ See Sequence for Donning and Doffing Personal Protective Equipment.



Providers should report any suspect cases to the NM Department of Health by calling the 24/7/365 EPIDEMIOLOGY HOTLINE at:

(505) 827-0006

TESTING DIRECTIONS FOR SLD

The goal is to sample 2 to 3 lesions from 2 to 3 different locations on the body and/or lesions with differing
appearances.

- ☐ Vigorously swab or scrub each lesion **with two** separate sterile, dry, polyester swabs (one for preliminary and one for confirmatory testing). **Do not use cotton swabs and do not swab more than one lesion with each swab.** You do not need to unroof the lesion.
- Each swab should each be placed in a separate dry container (such as sterile urine cup or tube).
- Do not add or store in viral or universal transport media (for SLD).
- Each lesion that is sampled requires a separate test requisition form.
- If you swab 3 lesions, you will have 6 swabs in 6 separate dry containers and 3 requisitions.
- Immediately refrigerate samples (2-8°C), and contact NMDOH at (505) 827-0006 for shipping instructions.

PATIENT PRECAUTIONS

- Advise patients waiting for results to isolate at home, avoid contact with others, and monitor symptoms. People should wear a mask and cover lesions if they must be around others.
- If positive for monkeypox, refer patients to CDC's Isolation and Prevention Practices for People with Monkeypox | Monkeypox | Poxvirus | CDC

