

Reported Cases of Gonorrhea and Chlamydia, New Mexico, 1970-2007

Year	Gonorrhea					Chlamydia*				
	New Mexico		U.S.			New Mexico		U.S.		
	Cases	Rate** per 100,000	Cases	Rate** per 100,000	NM Ranking	Cases	Rate** per 100,000	Cases	Rate** per 100,000	NM Ranking
1970	2882	283.7	600,072	297.2						
1971	2969	284.1	670,268	327.2						
1972	4377	410.6	767,215	369.7						
1973	4805	437.2	842,621	402.0						
1974	5433	484.2	906,121	428.2						
1975	7023	612.3	999,937	467.7						
1976	7137	611.0	1,001,994	464.1						
1977	6941	499.2	1,002,219	459.5						
1978	2600	461.7	1,013,436	459.7						
1979	4918	405.4	1,004,058	450.3						
1980	4676	362.5	1,004,029	445.1						
1981	4636	356.6	990,864	435.2						
1982	4402	338.6	960,633	417.9						
1983	3662	259.9	900,435	387.6						
1984	3372	245.2	878,556	372.5	33rd					
1985	3122	217.9	911,419	383.0	36th					
1986	2601	175.9	892,229	371.5	36th					
1987	2167	139.0	787,532	325.0	35th					
1988	1546	103.0	738,160	301.9	36th					
1989	1323	87.8	733,294	297.1	39th	2855	189.4	200,904	102.5	10th
1990	1235	81.2	690,042	276.4	41st	3597	236.4	323,663	160.2	9th
1991	1120	72.0	621,918	245.8	36th	4676	300.7	381,228	180.0	3rd
1992	921	57.8	502,858	196.0	39th	4482	280.9	409,694	182.4	2nd
1993	1014	62.0	444,649	171.7	36th	4754	290.1	405,332	178.1	1st
1994	1130	67.2	419,602	163.9	33rd	5037	299.4	451,785	192.5	2nd
1995	1054	61.3	392,651	147.5	34th	4285	249.1	478,577	187.8	3rd
1996	890	50.8	328,169	121.8	35th	4007	228.7	492,631	190.6	8th
1997	857	48.3	327,665	120.2	35th	4021	226.6	537,904	205.5	13th
1998	957	53.4	356,492	129.2	36th	3793	211.5	614,250	231.8	25th
1999	952	52.7	360,813	129.3	35th	4657	257.6	662,647	247.2	17th
2000	1136	62.5	363,136	128.7	35th	5052	277.7	709,452	251.4	13th
2001	1074	58.0	361,705	126.8	34th	6433	347.2	783,242	274.5	5th
2002	1472	78.5	351,852	122.0	32nd	7158	381.5	834,555	289.4	7th
2003	1143	60.2	335,104	115.2	37th	7294	383.9	877,478	301.7	7th
2004	1328	68.8	330,132	112.4	32nd	9182	475.8	929,462	316.5	4th
2005	1579	80.2	339,953	114.6	27th	8635	438.5	976,445	329.4	4th
2006	1758	87.3	358,366	120.9	25th	10054	499.2	1,030,911	347.8	5th
2007***	1797	87.3	350,475	117.1	23rd	9462	459.5	1,090,019	364.1	5th

Data as of 7/15/2008

* Chlamydia was not a reportable disease in New Mexico until mid 1988. Data are presented beginning in 1989.

** Rates are estimates based on data from the U.S. Census Bureau and the University of New Mexico, Bureau of Business and Economic Research (www.unm.edu/~bber)

*** For 2007, all U.S. numbers and rates as well as the New Mexico ranking are preliminary

During 2007 and 2008, the New Mexico STD Program conducted extensive quality assurance review of STD case investigations from 1999 through 2007. These QA activities have resulted in identifying hundreds of STD cases (mostly chlamydia and most from 2004-2006) that had not been entered into the STD database, hundreds of miscodes for Race/Ethnicity (mostly cases among Native American incorrectly coded as Other) and numerous incorrect ages and county of residence. These errors have been fixed. (See note below about errors that could not be corrected) As result, the numbers of cases on these tables will differ from what had been posted on this website prior to September 2008.

NOTE: Information for Cases from Santa Fe County for 2003 through 2005 are inaccurate. Data were not entered and have been lost. Thus, the numbers of cases for Santa Fe County in these tables are only a small proportion of the true number of cases. Please use these data with caution. Information for Native American Cases from McKinley County is missing from Gallup Indian Medical Center for 1999 through 2002. Thus, the numbers of cases for in McKinley County for these years is inaccurate. Please use these data with caution.