EPIDEMIOLOGY PROGRAM	Newly Diagnosed in 2013*				Persons Living with HIV or AIDS in New Mexico							
	Newly Diagnosed in 2013*			HIV			AIDS			Total		
	Ν	%	$Rate^{\dagger}$	N	%	$Rate^{\dagger}$	Ν	%	$Rate^{\dagger}$	N	%	$Rate^{\dagger}$
Total	145	100	8.4	1265	100	73.6	1861	100	108.2	3126	100	181.
at birth												
Male	126	87%	14.9	1,077	85%	127.7	1,663	89%	197.2	2,740	88%	325
Female	19	13%	2.2	188	15%	21.5	198	11%	22.6	386	12%	44.
e/Ethnicity												
American Indian/Alaska Native	20	14%	14.0	106	8%	74.3	144	8%	100.9	250	8%	175
Asian/Pacific Islander	1	1%	0	8	1%	29.9	11	1%	41.2	19	1%	71.
Black	7	5%	20.2	100	8%	288.2	98	5%	282.5	198	6%	570
Hispanic	80	55%	10.6	595	47%	78.9	810	44%	107.4	1405	45%	186
White	36	25%	4.7	443	35%	58.2	766	41%	100.7	1209	39%	158
Multi-race	1	1%	CNC	13	1%	CNC	32	2%	CNC	45	1%	CN
2 [‡]												
13-24	32	22%	9.1	71	6%	20.2	17	1%	4.8	88	3%	25.
25-34	44	30%	16.3	279	22%	103.2	136	7%	50.3	415	13%	153
35-44	32	22%	12.8	311	25%	124.3	350	19%	139.9	661	21%	264
45-54	23	16%	8.0	358	28%	124.2	736	40%	255.3	1094	35%	379
55+	14	10%	2.5	246	19%	44.0	622	33%	111.1	868	28%	156
gion [§]												
Northwest	17	12%	9.4	115	9%	63.7	155	8%	85.8	270	9%	149
Northeast	29	20%	11.6	232	18%	92.5	378	20%	150.8	610	20%	243
Metro	68	47%	9.1	603	48%	80.7	959	52%	128.4	1,562	50%	209
Southeast	7	5%	3.0	63	5%	26.9	110	6%	47.0	173	6%	73.
Southwest	24	17%	7.8	252	20%	82.1	259	14%	84.4	511	16%	166
nsmission Category			-	-					-	-		1
Men who have sex with men (MSM)	91	63%	CNC	744	59%	CNC	1172	63%	CNC	1916	61%	CN
Intravenous Drug Use (IDU)	9	6%	CNC	93	7%	CNC	165	9%	CNC	258	8%	CN
MSM/IDU	14	10%	CNC	91	7%	CNC	191	10%	CNC	282	9%	CN
High-Risk-Hetrosexual	14	10%	CNC	154	12%	CNC	180	10%	CNC	334	11%	CN
Other	0	0%	CNC	2	0%	CNC	10	1%	CNC	12	0%	CN
Perinatal Exposure	0	0%	CNC	4	0%	CNC	8	0%	CNC	12	0%	CN
No Risk Identified	17	12%	CNC	177	14%	CNC	135	7%	CNC	312	10%	CN

*Rates per 100,000 based on Bureau of Business and Economic Research Geospatial and Population Studies Program population estimates for 2013.

‡For new diagnoses, age at time of diagnosis. For persons living with HIV or AIDS, current age.

§For new diagnoses, residence at time of diagnosis. For persons living with HIV or AIDS, current residence.

CNC means Can Not Calculate - population data is

unavailable

Red - Indicates greater than 20% change from previous year for values > 20.

N < 20: Rate calculations for small numbers may be statistically unreliable. Fluctuations in the associated rates Region: New Mexico Department of Health constructed new health regions in 2012. Regional comparison to previous years is not recommended. A map depicting New Mexico's Health Regions is available for download from the New Mexico Department of Health at http://bis.health.state.nm.us/image/HealthRegions2012.png.