Includes Confirmed and Probable Cases Only

Table 1: New Mexico Pertussis Cases for 2012

Case S		
Confirmed	Probable	Total Cases
717	141	858

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 2: Pertussis Cases by County

County Name	Case Number	Percent of Total Cases	Incidence per 100,000
Bernalillo County	513	59.8	77.1
Chaves County	8	0.9	12.2
Cibola County	6	0.7	22.0
Colfax County	11	1.3	80.1
Curry County	4	0.5	8.3
Dona Ana County	12	1.4	5.7
Eddy County	1	0.1	1.9
Grant County	10	1.2	33.9
Lea County	10	1.2	15.4
Los Alamos County	4	0.5	22.3
Luna County	1	0.1	4.0
McKinley County	26	3.0	36.4
Mora County	1	0.1	20.5
Quay County	1	0.1	11.1
Rio Arriba County	7	0.8	17.4
San Juan County	18	2.1	13.8
San Miguel County	37	4.3	126.0
Sandoval County	15	1.7	11.3
Santa Fe County	64	7.5	44.3
Sierra County	6	0.7	50.2

Includes Confirmed and Probable Cases Only

Table 2: Pertussis Cases by County

County Name	Case Number	Percent of Total Cases	Incidence per 100,000
Socorro County	6	0.7	33.5
Taos County	6	0.7	18.2
Torrance County	1	0.1	6.1
Unknown	1	0.1	
Valencia County	89	10.4	115.9

Includes Confirmed and Probable Cases Only

Table 3: Pertussis Cases by Age Group

Age Group	Case Number	Percent
< 1 Year	72	8.4
1 to 5 Years	103	12.0
6 to 10 Years	179	20.9
11 to 18 Years	209	24.4
19 to 65 Years	260	30.3
65 Years or Older	34	4.0
Unknown	1	0.1

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 4: Pertussis Cases by Race/Ethnicity

Race/Ethnicity	Case Number	Percent
American Indian or Alaska Native	60	7.0
Asian or Pacific Islander	6	0.7
Black or African American	30	3.5
Hispanic	505	58.9
White	227	26.5
Multi-race	9	1.0
Unknown/Missing	21	2.4

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS).

Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 5: Cases of Pertussis by Race/Ethnicity

	Ethnicity			
Race	Hispanic or Latino	Not Hispanic or Latino	Unknown/ Missing	Total
American Indian or Alaska Native	10	48	2	60
	1.17	5.59	0.23	6.99
Asian or Pacific Islander	1	5	0	6
	0.12	0.58	0.00	0.70
Black or African American	6	23	1	30
	0.70	2.68	0.12	3.50
White	348	223	4	575
	40.56	25.99	0.47	67.02
Multi-race	5	4	0	9
	0.58	0.47	0.00	1.05
Unknown/Missing	160	7	11	178
	18.65	0.82	1.28	20.75
Total	530	310	18	858
	61.77	36.13	2.10	100.00

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 6: Pertussis Cases by Sex

Gender	Case Number	Percent
Female	487	56.8
Male	370	43.1
Unknown	1	0.1

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 7: Hospitalizations of Pertussis Cases

Yes		No		
Count	Percent Count		Percent	
40	4.7	807	95.3	

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only


	Hospitalized Cases					
Age Group	Yes No Unknown Tota					
< 1 Year	24 33.33	46 63.89	2 2.78	72		
1 to 5 Years	2 1.94	99 96.12	2 1.94	103		
6 to 10 Years	0 0.00	175 97.77	4 2.23	179		
11 to 18 Years	2 0.96	205 98.09	2 0.96	209		
19 to 65 Years	7 2.69	252 96.92	1 0.38	260		
65 Years or Older	5 14.71	29 85.29	0 0.00	34		
Unknown	0 0.00	1 100.00	0 0.00	1		
Total	40	807	11	858		

Table 8: Cases of Pertussis by Hospitalization and Age Group

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Case Counts of Pertussis by Week of Cough Onset for 2012



2012 --- Average, last 3 years