Table 1: New Mexico Pertussis Cases for 2012

| Case Status | | |
|-------------|----------|-------------|
| Confirmed | Probable | Total Cases |
| 12 | 6 | 18 |

Table 2: Pertussis Cases by County

| County Name | Case Number | Percent of Total Cases | Incidence per 100,000 |
|-------------------|----------------|------------------------------|-----------------------------|
| Bernalillo County | 9 | 50.0 | 1.4 |
| Colfax County | 3 | 16.7 | 21.8 |
| San Juan County | 1 | 5.6 | 0.8 |
| Santa Fe County | 3 | 16.7 | 2.1 |
| Unknown | 1 | 5.6 | |
| Valencia County | 1 | 5.6 | 1.3 |

Table 3: Pertussis Cases by Age Group

| Age Group | Case Number | Percent |
|----------------|----------------|---------|
| < 1 Year | 1 | 5.6 |
| 1 to 5 Years | 1 | 5.6 |
| 6 to 10 Years | 8 | 44.4 |
| 11 to 18 Years | 2 | 11.1 |
| 19 to 65 Years | 6 | 33.3 |

Table 4: Pertussis Cases by Race/Ethnicity

| Race/Ethnicity | Case Number | Percent |
|-----------------|----------------|---------|
| Hispanic | 12 | 66.7 |
| White | 5 | 27.8 |
| Unknown/Missing | 1 | 5.6 |

Table 5: Cases of Pertussis by Race/Ethnicity

| | Ethnicity | | | |
|----------------------------------|--------------------------|---------------------------------|---------------------|--------|
| Race | Hispanic or Latino | Not Hispanic or Latino | Unknown/ Missing | Total |
| American Indian or Alaska Native | 10 | 48 | 2 | 60 |
| | 1.14 | 5.47 | 0.23 | 6.84 |
| Asian or Pacific Islander | 1 | 5 | 0 | 6 |
| | 0.11 | 0.57 | 0.00 | 0.68 |
| Black or African American | 6 | 23 | 1 | 30 |
| | 0.68 | 2.62 | 0.11 | 3.42 |
| White | 357 | 229 | 4 | 590 |
| | 40.71 | 26.11 | 0.46 | 67.27 |
| Multi-race | 5 | 4 | 0 | 9 |
| | 0.57 | 0.46 | 0.00 | 1.03 |
| Unknown/Missing | 163 | 7 | 12 | 182 |
| | 18.59 | 0.80 | 1.37 | 20.75 |
| Total | 542 | 316 | 19 | 877 |
| | 61.80 | 36.03 | 2.17 | 100.00 |

Table 6: Pertussis Cases by Sex

| Gender | Case Number | Percent |
|--------|----------------|---------|
| Female | 10 | 55.6 |
| Male | 8 | 44.4 |

Table 7: Hospitalizations of Pertussis Cases

| Yes | | No | |
|-------|---------|-------|---------|
| Count | Percent | Count | Percent |
| 1 | 5.6 | 17 | 94.4 |


Table 8: Cases of Pertussis by Hospitalization and Age Group

| | Hospitalized Cases | | | | |
|----------------|--------------------|-------------|----|--|--|
| Age Group | Yes No Tota | | | | |
| < 1 Year | 1 100.00 | 0.00 | 1 | | |
| 1 to 5 Years | 0.00 | 1 100.00 | 1 | | |
| 6 to 10 Years | 0.00 | 8 100.00 | 8 | | |
| 11 to 18 Years | 0.00 | 2 100.00 | 2 | | |
| 19 to 65 Years | 0.00 | 6 100.00 | 6 | | |
| Total | 1 | 17 | 18 | | |

Pertussis Report for Dates 01/01/2013 to 01/27/2013

Includes Confirmed and Probable Cases Only

Case Counts of Pertussis by Week of Cough Onset for 2013



Week of Cough Onset

2013 --- Average, last 3 years