Includes Confirmed and Probable Cases Only

Table 1: New Mexico Pertussis Cases for 2013

Case S		
Confirmed	Probable	Total Cases
288	76	364

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 2: Pertussis Cases by County

County Name	Case Number	Percent of Total Cases	Incidence per 100,000
Bernalillo County	128	35.2	19.2
Chaves County	2	0.5	3.0
Colfax County	7	1.9	51.0
Curry County	2	0.5	4.1
Dona Ana County	5	1.4	2.4
Eddy County	5	1.4	9.3
Grant County	16	4.4	54.3
Guadalupe County	2	0.5	
Lea County	15	4.1	23.1
Los Alamos County	20	5.5	111.5
McKinley County	8	2.2	11.2
Quay County	2	0.5	22.2
Rio Arriba County	8	2.2	19.9
Roosevelt County	1	0.3	5.0
San Juan County	11	3.0	8.4
San Miguel County	1	0.3	3.4
Sandoval County	12	3.3	9.1
Santa Fe County	71	19.5	49.1
Sierra County	2	0.5	16.7
Socorro County	3	0.8	16.8

Includes Confirmed and Probable Cases Only

Table 2: Pertussis Cases by County

County Name	Case Number	Percent of Total Cases	Incidence per 100,000
Taos County	20	5.5	60.6
Torrance County	8	2.2	48.9
Union County	1	0.3	
Unknown	2	0.5	
Valencia County	12	3.3	15.6

Includes Confirmed and Probable Cases Only

Table 3: Pertussis Cases by Age Group

Age Group	Case Number	Percent
<1 Year	28	7.7
1 to 5 Years	47	12.9
6 to 10 Years	85	23.4
11 to 18 Years	90	24.7
19 to 65 Years	98	26.9
65 Years or Older	16	4.4

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 4: Pertussis Cases by Race/Ethnicity

Race/Ethnicity	Case Number	Percent
American Indian or Alaska Native	22	6.0
Black or African American	10	2.7
Hispanic	188	51.6
White	133	36.5
Multi-race	4	1.1
Unknown/Missing	7	1.9

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification

of historical cases.

Includes Confirmed and Probable Cases Only

Table 5: Cases of Pertussis by Race/Ethnicity

	Ethnicity				
Race	Hispanic or Latino	Not Hispanic or Latino	Unknown/ Missing	Total	
American Indian or Alaska Native	3	19	0	22	
	0.82	5.22	0.00	6.04	
Black or African American	4	6	0	10	
	1.10	1.65	0.00	2.75	
White	150	132	1	283	
	41.21	36.26	0.27	77.75	
Multi-race	3	1	0	4	
	0.82	0.27	0.00	1.10	
Unknown/Missing	39	4	2	45	
	10.71	1.10	0.55	12.36	
Total	199	162	3	364	
	54.67	44.51	0.82	100.00	

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 6: Pertussis Cases by Sex

Gender	Case Number	Percent
Female	197	54.1
Male	166	45.6
Unknown	1	0.3

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

Table 7: Hospitalizations of Pertussis Cases

Yes		No		
Count	t Percent Cou		Percent	
16	4.4	346	95.6	


*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

	Hospitalized Cases				
Age Group	Yes No Unknown To				
< 1 Year	11 39.29	17 60.71	0 0.00	28	
1 to 5 Years	0 0.00	47 100.00	0 0.00	47	
6 to 10 Years	1 1.18	84 98.82	0 0.00	85	
11 to 18 Years	1 1.11	87 96.67	2 2.22	90	
19 to 65 Years	1 1.02	97 98.98	0 0.00	98	
65 Years or Older	2 12.50	14 87.50	0 0.00	16	
Total	16	346	2	364	

Table 8: Cases of Pertussis by Hospitalization and Age Group

*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.



Case Counts of Suspected Pertussis by Week of Cough Onset for 2013