Includes Confirmed and Probable Cases Only

### Table 1: New Mexico Pertussis Cases for 2013

Case S		
Confirmed	Probable	Total Cases
322	88	410

\*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

# Table 2: Pertussis Cases by County

County Name	Case Number	Percent of Total Cases	Incidence per 100,000
Bernalillo County	145	35.4	21.8
Chaves County	2	0.5	3.0
Colfax County	7	1.7	51.0
Curry County	4	1.0	8.3
Dona Ana County	6	1.5	2.9
Eddy County	5	1.2	9.3
Grant County	19	4.6	64.5
Guadalupe County	2	0.5	
Lea County	17	4.1	26.2
Los Alamos County	20	4.9	111.5
McKinley County	19	4.6	26.6
Quay County	2	0.5	22.2
Rio Arriba County	8	2.0	19.9
Roosevelt County	1	0.2	5.0
San Juan County	12	2.9	9.2
San Miguel County	1	0.2	3.4
Sandoval County	16	3.9	12.1
Santa Fe County	74	18.0	51.2
Sierra County	2	0.5	16.7
Socorro County	4	1.0	22.3

Includes Confirmed and Probable Cases Only

# Table 2: Pertussis Cases by County

County Name	Case Number	Percent of Total Cases	Incidence per 100,000
Taos County	21	5.1	63.6
Torrance County	10	2.4	61.1
Union County	1	0.2	•
Valencia County	12	2.9	15.6

**Includes Confirmed and Probable Cases Only** 

#### Table 3: Pertussis Cases by Age Group

Age Group	Case Number	Percent
<1 Year	29	7.1
1 to 5 Years	51	12.4
6 to 10 Years	89	21.7
11 to 18 Years	106	25.9
19 to 65 Years	116	28.3
65 Years or Older	19	4.6

\*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

**Includes Confirmed and Probable Cases Only** 

#### Table 4: Pertussis Cases by Race/Ethnicity

Race/Ethnicity	Case Number	Percent
American Indian or Alaska Native	32	7.8
Black or African American	11	2.7
Hispanic	205	50.0
White	148	36.1
Multi-race	5	1.2
Unknown/Missing	9	2.2

\*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS).

Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

**Includes Confirmed and Probable Cases Only** 

#### Table 5: Cases of Pertussis by Race/Ethnicity

	Ethnicity				
Race	Hispanic or Latino	Not Hispanic or Latino	Unknown/ Missing	Total	
American Indian or Alaska Native	3	29	0	32	
	0.73	7.07	0.00	7.80	
Black or African American	5	6	0	11	
	1.22	1.46	0.00	2.68	
White	165	146	2	313	
	40.24	35.61	0.49	76.34	
Multi-race	3	2	0	5	
	0.73	0.49	0.00	1.22	
Unknown/Missing	41	4	4	49	
	10.00	0.98	0.98	11.95	
Total	217	187	6	410	
	52.93	45.61	1.46	100.00	

\*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

**Includes Confirmed and Probable Cases Only** 

### **Table 6: Pertussis Cases by Sex**

Gender	Case Number	Percent
Female	219	53.4
Male	191	46.6

\*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

Includes Confirmed and Probable Cases Only

### **Table 7: Hospitalizations of Pertussis Cases**

Yes		No		
Count	Percent	Count	Percent	
17	4.2	389	95.8	


\*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.

**Includes Confirmed and Probable Cases Only** 

	Hospitalized Cases					
Age Group	Yes No Unknown Total					
< 1 Year	10 34.48	19 65.52	0 0.00	29		
1 to 5 Years	0 0.00	51 100.00	0 0.00	51		
6 to 10 Years	1 1.12	87 97.75	1 1.12	89		
11 to 18 Years	1 0.94	102 96.23	3 2.83	106		
19 to 65 Years	2 1.72	114 98.28	0 0.00	116		
65 Years or Older	3 15.79	16 84.21	0 0.00	19		
Total	17	389	4	410		

#### Table 8: Cases of Pertussis by Hospitalization and Age Group

\*This report is prepared by the Infectious Disease Epidemiology Bureau (IDEB) using the New Mexico Electronic Disease Surveillance System (NM-EDSS). Data are provisional and may change significantly upon the completion of ongoing investigations. Data for prior years are also subject to minor revisions due to corrections or identification of historical cases.



Case Counts of Suspected Pertussis by Week of Cough Onset for 2013