


NEW MEXICO INFLUENZA SURVEILLANCE UPDATE 2015 - 2016 Influenza Season

Epidemiology and Response Division, New Mexico Department of Health (NMDOH)

Influenza Activity in New Mexico for Week Ending January 9, 2016 (MMWR Week 1)

Influenza in the state was at low activity by several indicators¹:


1. Outpatient visits for influenza-like illness (ILI): Twenty-four sentinel outpatient provider sites reported a total of 11,106 patient visits, of which 182 (1.6%) were identified as visits for ILI²; which is below the national baseline of 2.1%. Nationally, the percentage of ILI visits was 2.0%; which is also below baseline (see graph below).



Weekly ILI and lab data may change as additional reports are compiled.

2. ILI Activity by State Health Regions: by NMDOH regions within the state (refer to page 4 table footnote defining the regions by county). Weekly ILI ranged from 0.7% (NE Region) to 2.2% (SE and SW Regions). See graph below for the current season's weekly ILI by region.


² Influenza-like Activity (ILI) is defined as Fever (≥ 100°F [37.8° C], oral or equivalent) AND cough and/or sore throat in the absence of a KNOWN cause other than influenza.



Sentinel Outpatient Reporting Sites by Region:

NW Region San Juan Regional Medical Center, Farmington and Acoma-Canoncito-Laguna Indian Hea			
NE Region	Hospital, Acoma Raton Family Practice Associates, Raton; Taos-Picuris Indian Health Clinic, Taos; Pecos Valley Medical Center, Pecos; La Familia Medical Clinic, Santa Fe; Los Alamos Family Care Clinic, Los		
Metro Region	Alamos; Jicarilla Apache Indian Health Clinic, Dulce; Children's Clinic PA, Los Alamos Lovelace Downtown Medical Center, Albuquerque; University of New Mexico Student Health Clinic, Albuquerque; Presbyterian Medical Group-Atrisco, Northside, Pediatric Urgent Care; Albuquerque Health Partners Urgent Care, Rio Rancho		
SE Region	Carlsbad Medical Center, Carlsbad; Lea Regional Medical Center, Hobbs; Eastern New Mexico Medical Center, Roswell; Roosevelt General Hospital Clinic, Portales		
SW Region	Gila Regional Medical Center, Silver City; Hidalgo Medical Services Clinic, Lordsburg; Ben Archer Health Center, Deming, Columbus, and Dona Ana; La Clinica de Familia, Sunland Park; Mescalero Apache Indian Health Hospital, Mescalero; New Mexico Mining and Technology Institute, Socorro		

- 3. Laboratory tests for influenza: Since the start of the influenza season, the New Mexico Scientific Laboratory Division reported three respiratory specimens tested positive for Influenza A. Two were subtype 2009 H1N1 and one was subtype H3N2.
- **4. Influenza and pneumonia-related deaths:** There have been three influenza-related deaths in adults and 50 pneumonia-related deaths among New Mexico residents reported since the start of the 2015-2016 influenza season.
- 5. Geographic Spread of Influenza in the State: NMDOH reported the state influenza activity level as 'Local" to the Centers for Disease Control and Prevention (CDC). See the table on page 4 for the activity level definitions. The map below displays New Mexico activity in comparison to influenza activity in bordering states, regions, and nationally for the week ending January 9, 2015.



^{*} This map indicates geographic spread & does not measure the severity of influenza activity

This information is collected by the Infectious Disease Epidemiology Bureau, Epidemiology Response Division, New Mexico Department of Health.

For questions, please call 505-827-0006.

For more information on influenza go to the NMDOH web page: http://www.health.state.nm.us/flu/ or the CDC web page: http://www.cdc.gov/flu/about/disease/

TABLE OF INFLUENZA ACTIVITY LEVELS

Activity Level	ILI activity*/Outbreaks		Laboratory data	
No activity	Low	And	No lab confirmed cases [†]	
	Not increased	And	Isolated lab-confirmed cases	
Sporadic	OR			
	Not increased	And	Lab confirmed outbreak in one institution [‡]	
	Increased ILI in 1 region**;		Recent (within the past 3 weeks) lab evidence	
	ILI activity in other regions is	And	of influenza in region with increased ILI	
	not increased		OR	
Local				
Locai	2 or more institutional		Recent (within the past 3 weeks) lab evidence	
	outbreaks (ILI or lab		of influenza in region with the outbreaks; virus	
	confirmed) in 1 region; ILI	And	activity is no greater than sporadic in other	
	activity in other regions is		regions	
	not increased			
	Increased ILI in less than	And	Recent (within the past 3 weeks) lab confirmed	
	half of the regions (2 or less)		influenza in the affected regions	
Regional	OR			
i togiona.	Institutional outbreaks (ILI or		Recent (within the past 3 weeks) lab confirmed	
	lab confirmed) in less than	And	influenza in the affected regions	
	half of the regions (2 or less)			
	Increased ILI and/or		Recent (within the past 3 weeks) lab confirmed	
Widosproad	institutional outbreaks (ILI or	And	influenza in the state.	
Widespread	lab confirmed) in at least 3			
	of the 5 regions			

Influenza-like illness: Fever (\geq 100°F [37.8°C], oral or equivalent) <u>and</u> cough <u>and/or</u> sore throat (in the absence of a known cause other than influenza).

[†]Lab confirmed case = case confirmed by rapid diagnostic test, antigen detection, culture, or PCR. Care should be given when relying on results of point of care rapid diagnostic test kits during times when influenza is not circulating widely. The sensitivity and specificity of these tests vary and the predicative value positive may be low outside the time of peak influenza activity. Therefore, at the start of the season, NMDOH uses lab confirmation by PCR or culture to declare the presence of lab-confirmed influenza in the state. Positivity by other lab methods is recognized and tracked as another influenza activity indicator as the season progresses.

[‡] Institution includes nursing home, hospital, prison, school, etc.

^{**}Region: population under surveillance in a defined geographical subdivision of a state. A region could be comprised of 1 or more counties and would be based on each state's specific circumstances. In New Mexico, the NMDOH regions are: Northwest Region: San Juan, McKinley, and Cibola counties; Northeast Region: Los Alamos, Rio Arriba, Taos, Colfax, Union, Mora, Harding, Santa Fe, San Miguel and Guadalupe counties; Metro Region: Bernalillo, Sandoval, Torrance and Valencia counties; Southeast Region: Quay, Curry, DeBaca, Lincoln, Roosevelt, Chaves, Eddy and Lea counties, and Southwest Region: Socorro, Catron, Grant, Sierra, Otero, Dona Ana, Luna and Hidalgo counties.