

**NEW MEXICO SELECTED HEALTH STATISTICS
ANNUAL REPORT 2012**

**THE STATE CENTER FOR HEALTH STATISTICS
Bureau of Vital Records and Health Statistics**



STATE OF NEW MEXICO



Susana Martinez, Governor

Department of Health

Retta Ward, M.P.H., Cabinet Secretary

Brad McGrath Lynn Gallagher
Chief Deputy Secretary Deputy Secretary

Epidemiology and Response Division

Michael G. Landen, M.D., M.P.H.
Director and State Epidemiologist

Bureau of Vital Records and Health Statistics

Mark Kassouf
Bureau Chief and Director, State Center for Health Statistics

Renee Valencia, M.A.

Acting State Registrar and Registration Manager

Issuance Manager

The 2012 Annual Report was prepared by the Statistics and Epidemiology Unit of the Bureau of Vital Records and Health Statistics: Irma Montoya Project Coordinator, Camille Clifford, Tony Ortiz, Rosemary Perez-Mendoza, Anne Worthington and assisted by Nancy Blake.

Much of the data contained in the tables and figures of this report were generated using dataset queries from New Mexico's Indicator-Based Information System (NM-IBIS). <http://ibis.health.state.nm.us/>

This publication is available at <http://www.vitalrecordsnm.org/StatReports.shtml>.

Photos were provided by Jim Orr (jamesorr.smugmug.com), Randal K. Roberts, the New Mexico Department of Tourism, Sally King, Bandelier National Monument and Jon Branch.

The mission of the New Mexico Department of Health is to promote health and sound health policy, prevent diseases and disability, improve health service systems and assure that essential public health functions and safety net services are available to New Mexicans.

Executive Summary

Population Highlights

New Mexico's estimated 2012 population reached 2,091,432, reflecting an increase of 1.3% since 2010. The state's population is projected to reach 2,613,332 by 2030.

For race/ethnicity in 2012, 41.4% of New Mexicans were White; 46.4% were Hispanic; 8.8% were American Indian or Alaska Native; 2.0% were Black or African American; and 1.5% were Asian or Pacific Islander.

Natality Highlights

In 2012 there were 26,992 births to New Mexico resident mothers, translating to a birth rate of 12.9 births per 1,000 population, a record low for New Mexico. The state birth rate has been consistently higher than the national rate.

New Mexico teen birth rates continue to be higher than the national teen birth rates. The 2012 teen birth rate was 22.7 per 1,000 females ages 15-17 years and 77.3 per 1,000 females ages 18-19 years. The United States rates for 2011 were 15.4 for 15-17 year olds and 54.1 for 18-19 years.

More than half (51.9%) of 2012 New Mexico births were to single mothers, compared to 40.7% nationally.

The proportion of infants weighing less than 2,500 grams was unchanged from 2011 at 77.3%.

The lowest level of prenatal care for New Mexican mothers occurred in the less than 15 years of age group followed by the 15-17 year old age group and then the 18-19 year old age group.

Mortality Highlights

In 2012, there were 16,640 deaths among New Mexico residents, translating to an age-adjusted death rate of 749.4 deaths per 100,000, population adjusted to U.S. standard population.

The leading cause of death among New Mexico residents was Malignant Neoplasm (Cancer) with 3,422 deaths, exceeding Heart Disease with 3,306 deaths. Accidents (Unintentional Injuries) accounted for 1,331 deaths.

Infant mortality, which is deaths of children under 1 year of age per 1,000 live births, was higher in New Mexico in 2012 (6.9) than in 2011 (5.2). New Mexico's rate was higher than the 2010 U.S. rate of 6.2.

Table of Contents

Population Section	11
Population Estimates and Projections	12
Population Distribution	13
Age	13
Race and Ethnicity	14
Natality Section	25
Birth Numbers and Rates	26
Race and Ethnicity	27
Age of Mother	27
Teen Mothers	28
Births to Single Mothers	28
Birth Order	30
Multiple Births	31
Birthweight	32
Gestational Age	34
Prenatal Care	36
Kessner Index	36
Facility and Attendant	37
Method of Delivery	38
Mortality Section	71
All Causes of Death	72
Leading Causes of Death	75
Selected Causes	79
Injury Deaths and Alcohol- and Drug-Induced Deaths	79
Maternal Mortality	82
Infant Mortality	83
Disposition of Remains	86
Fetal Mortality Section	125
Abortion Section	131
Technical Appendix	135

Population Section

Figure P-1	Population Counts and Projections, New Mexico	12
Figure P-2	Population Pyramids, New Mexico and United States	13
Figure P-3	Population Distribution by Race/Ethnicity, New Mexico	14
Figure P-4	Population Distribution by Race/Ethnicity, Age, and Sex, New Mexico	15
Table P-1	Population Percent by Age Group, New Mexico and United States	16
Table P-2	Population Number by Age and County, New Mexico – Both Sexes	17
Table P-3	Population Number by Age and County, New Mexico – Males	18
Table P-4	Population Number by Age and County, New Mexico – Females	19
Table P-5	Population Number by Race/Ethnicity, Sex, and Age, New Mexico and United States	20
Table P-6	Population Number by Race/Ethnicity, County, and Health Region, New Mexico	21
Table P-7	Births and Deaths by City, New Mexico	22
Table P-8	Summary and Trends in New Mexico Health Statistics	23

Natality Section

Figure N-1	Birth Rates, New Mexico and United States	26
Figure N-2	Birth Rates by Race/Ethnicity, New Mexico	27
Figure N-3	Birth Rates by Mother's Age, New Mexico	27
Figure N-4	Teen (15-17 Years and 18-19 Years) Birth Rates, New Mexico and United States	28
Figure N-5	Percent of Births to Single Mothers, New Mexico and United States	29
Figure N-6	Percent of Births to Single Mothers by Race/Ethnicity, New Mexico	29
Figure N-7	Percent of Births to Single Mothers by Age, New Mexico	30
Figure N-8	Percent of Birth Order, New Mexico	31
Figure N-9	Number of Births in Single and Multiple Deliveries by Year, New Mexico	31
Figure N-10	Percent of Births with Low Birthweight, New Mexico and United States	32
Figure N-11	Percent of Births with Low Birthweight by Mother's Race/Ethnicity, New Mexico	33
Figure N-12	Percent of Births with Low Birthweight by Mother's Age, New Mexico and United States	34
Figure N-13	Percent of Preterm Births by Race/Ethnicity, New Mexico	35
Figure N-14	Percent Gestational Age by Birthweight, New Mexico	35
Figure N-15	Percent of Births by Level of Prenatal Care (Modified Kessner Index), New Mexico	36

Figure N-16	Percent of Births with Low or No Prenatal Care by Mother’s Age, New Mexico	37
Figure N-17	Percent of Births Occurring in New Mexico by Attendant Type, New Mexico	38
Figure N-18	Distribution of Delivery Method by Year, New Mexico	39
Table N-1	Natality Characteristics by Resident County and Health Region, New Mexico Residents and United States	40
Table N-2	Number of Births by Year, Mother's Age, and Race/Ethnicity, New Mexico and United States	41
Table N-3	Birth Rates by Mother's Age and Race/Ethnicity, New Mexico and United States	42
Table N-4(a)	Number of Births by Mother's Race/Ethnicity, County, and Health Region, New Mexico and United States	43
Table N-4(b)	Percent of Births by Mother’s Race/Ethnicity, County , and Health Region, New Mexico and United States	44
Table N-5(a)	Number of Births by Year, County, and Health Region, New Mexico and United States	45
Table N-5(b)	Birth Rates by Year, County, and Health Region, New Mexico and United States	46
Table N-6(a)	Number of Births by Mother’s Age, County, and Health Region, New Mexico	47
Table N-6(b)	Birth Rates by Mother’s Age, County, and Health Region, New Mexico	48
Table N-7	Number and Percent of Births to Single Mother’s by Mother’s Age and Race/Ethnicity, New Mexico	49
Table N-8	Number and Percent of Births to Single Mothers by County and Health Region, New Mexico	50
Table N-9	Number and Percent of Births to Single Mothers by Mother’s Age, New Mexico	51
Table N-10	Percent Distribution of Birth Order by Mother’s Race/Ethnicity, New Mexico	52
Table N-11(a)	Number of Births by Mother’s Age, Race/Ethnicity, and Infant’s Birthweight, New Mexico	53
Table N-11(b)	Percent of Births by Mother’s Age, Race/Ethnicity, and Infant’s Birthweight, New Mexico	54
Table N-12	Number of Single and Multiple Births by Mother’s Age and Race/Ethnicity, New Mexico	55
Table N-13(a)	Number of Births by Birthweight (Grams), County, and Health Region, New Mexico and United States	56
Table N-13(b)	Percent of Births by Birthweight (Grams), County, and Health Region, New Mexico and United States	57
Table N-14(a)	Number of Births by Level of Prenatal Care, County, and Health Region, New Mexico	58
Table N-14(b)	Percent of Births by Level of Prenatal Care, County, and Health Region, New Mexico	59
Table N-15(a)	Number of Births with Low or No Prenatal Care by Mother’s Race/Ethnicity, County, and Health Region, New Mexico	60
Table N-15(b)	Percent of Births with Low or No Prenatal Care by Mother’s Race/Ethnicity, County, and Health Region, New Mexico	61
Table N-16(a)	Number of Births with Low or No Prenatal Care by Mother's Age, County, and Health Region, New Mexico	62

Table N-16(b)	Percent of Births with Low or No Prenatal Care by Mother's Age, County, and Health Region, New Mexico	63
Table N-17(a)	Number of Births by Trimester Prenatal Care Began, County, and Health Region, New Mexico	64
Table N-17(b)	Percent of Births by Trimester Prenatal Care Began, County, and Health Region, New Mexico	65
Table N-18	Number and Percent of Resident Births that Occurred and Did Not Occur in New Mexico by County of Residence	66
Table N-19	Number and Percent of Births by Facility Type and Attendant Type, New Mexico	67
Table N-20(a)	Number of Births by Birth Method, Mother's Age, and Race/Ethnicity, New Mexico	68
Table N-20(b)	Percent of Births by Birth Method, Mother's Age, and Race/Ethnicity, New Mexico	69
Table N-21	Number of Births by Birth Method, County, and Health Region, New Mexico	70

Mortality Section

Figure M-1	Percent of Deaths by Age Group and Sex, New Mexico	72
Figure M-2	Death Rates for All Causes by Sex and Race/Ethnicity, New Mexico	73
Figure M-3	Death Rates for All Causes by Race/Ethnicity, New Mexico	74
Figure M-4	Death Rates, New Mexico and United States	75
Figure M-5	Leading Causes of Death, New Mexico and United States	76
Figure M-6	Death Rates by Cause, New Mexico	77
Figure M-7(a)	Malignant Neoplasm Rates by Race/Ethnicity, New Mexico	78
Figure M-7(b)	Heart Disease Rates by Race/Ethnicity, New Mexico	78
Figure M-7(c)	Unintentional Injury (Accidents) Rates by Race/Ethnicity, New Mexico	79
Figure M-8	Death Rates for Unintentional Injuries by Type, New Mexico	80
Figure M-9	Death Rates for Intentional Injuries by Type, New Mexico	81
Figure M-10	Percent of Firearm Injury Deaths by Manner, New Mexico	82
Figure M-11	Infant Mortality Rates, New Mexico and United States	83
Figure M-12	Infant Mortality Rates by Race/Ethnicity, New Mexico	84
Figure M-13(a)	Neonatal Mortality Rates, New Mexico and United States	85
Figure M-13(b)	Postneonatal Mortality Rates, New Mexico and United States	85
Figure M-14	Percent Distribution by Method of Disposition, New Mexico Occurrence	86

Table M-1(a)	Number of Deaths by Age, Race/Ethnicity, and Sex, New Mexico and United States	87
Table M-1(b)	Death Rates by Age, Race/Ethnicity, and Sex, New Mexico and United States	88
Table M-2(a)	Number of Deaths by Age, Sex, County, and Health Region, New Mexico and United States – Both Sexes	89
Table M-2(b)	Number of Deaths by Age, Sex, County, and Health Region, New Mexico and United States – Females	90
Table M-2(c)	Number of Deaths by Age, Sex, County, and Health Region, New Mexico and United States – Males	91
Table M-3(a)	Death Rates by Age, Sex, County, and Health Region, New Mexico and United States – Both Sexes	92
Table M-3(b)	Death Rates by Age, Sex, County and Health Region, New Mexico and United States – Females	93
Table M-3(c)	Death Rates by Age, Sex, County and Health Region, New Mexico and United States – Males	94
Table M-4(a)	Number of Deaths Ranked by Leading Causes of Death by County and Health Region, New Mexico and United States	95
Table M-4(b)	Death Rates for Ranked Leading Causes of Death by County and Health Region, New Mexico and United States	96
Table M-5	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, New Mexico – All Ages New Mexico and United States	97
Table M-6	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, New Mexico – Ages 1 to 4 Years	98
Table M-7	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, New Mexico – Ages 5 to 14 Years	99
Table M-8	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, New Mexico – Ages 15 to 24 Years	100
Table M-9	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, New Mexico – Ages 25 to 44 Years	101
Table M-10	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, New Mexico – Ages 45 to 64 Years	102
Table M-11	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, New Mexico – Ages 65 to 84 Years	103
Table M-12	Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death, New Mexico – Ages 85 Plus Years	104
Table M-13(a)	Number of Deaths by Race/Ethnicity and Sex, New Mexico and United States	105
Table M-13(b)	Death Rates by Race/Ethnicity and Sex, New Mexico and United States	106
Table M-14(a)	Number of Deaths for Leading Causes of Death, New Mexico	107
Table M-14(b)	Death Rates for Leading Causes of Death, New Mexico	108
Table M-15(a)	Number of Deaths for 39 Selected Causes by Age, New Mexico	109
Table M-15(b)	Death Rates for 39 Selected Causes by Age, New Mexico	110
Table M-16(a)	Number of Deaths for 39 Selected Causes by Sex, New Mexico	111
Table M-16(b)	Death Rates for 39 Selected Causes by Sex, New Mexico	112

Table M-17(a)	Number of Deaths for 39 Selected Causes by Health Region, New Mexico	113
Table M-17(b)	Death Rates for 39 Selected Causes by Health Region, New Mexico	114
Table M-18(a)	Number of Death for 39 Selected Causes by Race/Ethnicity, New Mexico	115
Table M-18(b)	Death Rates for 39 Selected Causes by Race/Ethnicity, New Mexico	116
Table M-19	Number of Injury Deaths by Sex, County, and Health Region, New Mexico and United States	117
Table M-20	Death Rates for Injury by Sex, County, and Health Region, New Mexico and United States	118
Table M-21	Number of Deaths due to Firearms, Alcohol, Drugs, and Injury at Work by Sex, County, and Health Region, New Mexico and United States	119
Table M-22	Death Rates due to Firearms, Alcohol, Drugs, and Injury at Work by Sex, County, and Health Region, New Mexico and United States	120
Table M-23	Maternal Mortality Deaths and Death Rates, New Mexico	121
Table M-24(a)	Number of Infant Deaths by Sex and Race/Ethnicity, New Mexico	122
Table M-24(b)	Infant Death Rates by Sex and Race/Ethnicity, New Mexico	123
Table M-25	Number of Infant Deaths by Cause, New Mexico and United States	124
Table M-26	Method of Final Disposition for Deaths Occurring in New Mexico, New Mexico	124

Fetal Mortality Section

Figure F-1	Fetal Mortality Rates, New Mexico and United States	126
Figure F-2	Fetal Mortality Rates by Mother's Age, New Mexico and United States	127
Table F-1	Number of Fetal Deaths and Fetal Death Rates by Mother's Race/Ethnicity, New Mexico and United States	128
Table F-2	Number of Fetal Deaths and Fetal Death Rates by Cause, New Mexico	129

Abortion Section

Figure A-1	Percent Distribution of Induced Abortions by Weeks Gestation, New Mexico Occurrence	132
Figure A-2	Ratio of Induced Abortions, New Mexico Occurrence	133
Table A-1	Number and Percent of Induced Abortions by Woman's Age and Race/Ethnicity, New Mexico and United States	134

Intentionally Left Blank

Population Section




POPULATION SECTION

POPULATION ESTIMATES AND PROJECTIONS

The total United States population estimate for 2012 was 313,914,040. This total represents a 1.7% increase in the nation's population since 2010 (U.S. Census Bureau, released June 2013). New Mexico's 2012 estimated population was 2,091,432, which is a 1.3% increase from 2010. This is also an increase of 14.4% since 2000 (Table P-8).

New Mexico's population is projected to reach 2,351,724 in 2020 and 2,613,332 in 2030. These are projected increases from 2012 of 12.4% and 25.0%, respectively (Figure P-1).

**Figure P-1, Population Counts and Projections,
New Mexico, 1850 to 2030**




POPULATION DISTRIBUTION

Age

In 2012, for the United States, the largest portion of the population was between the ages of 45 and 54 years. The largest portion for New Mexico was 65 years and older, and age group 15 to 24 years; each with 14% (Table P-1).

In 1900, both the United States and New Mexico were characterized by low life expectancy and high fertility rates resulting in a high proportion of young people and a low proportion of elderly. After World War II, the percent of the population in the younger age groups declined to produce a more uniform pyramid. At the same time, the national figure becomes more rectangular in shape, indicating a more even distribution of ages in the population. New Mexico's older population was slower to increase than that of the United States. By 2010, the State population distribution more closely reflected that of the Nation (Figure P-2).


**Figure P-2, Population Pyramids
New Mexico and United States, 1900, 1950, and 2010**



Race and Ethnicity

The New Mexico Department of Health combines race and ethnicity for reporting purposes. The Department’s race and ethnicity guidelines, which were revised for 2012, are described in the Technical Appendix.

Figure P-3, Population Distribution by Race/Ethnicity, New Mexico, 2012




According to 2012 state population estimates, 46.4% of New Mexicans were Hispanic, and 41.4 % were White (Figure P-3). For 2012, the Hispanic category now includes Black, American Indian, or Asian or Pacific Islander populations who reported Hispanic ethnicity. The American Indian or Alaska Native population comprised 8.8% of New Mexico’s population; the Black or African American population made up 2.0%; and the Asian or Pacific Islander population constituted another 1.5%.

The American Indian or Alaska Native and the Hispanic populations had higher percents of individuals under age 25 years compared to the other racial/ethnic groups. The White population had the highest percent of persons 65 years of age and older (Figure P-4).

Population Section

**Figure P-4, Population Distribution by Race/Ethnicity, Age, and Sex
New Mexico, 2012**



Population Section

**Table P-1 Population Percent by Age Group
New Mexico and United States, 1990, 2000, and 2012**

Age Group	1990		2000		2012	
	NM	US	NM	US	NM	US
< 5	8.5	7.4	7.2	6.8	7.0	6.4
5 to 14	16.6	14.2	15.8	14.6	13.7	13.1
15 to 24	14.6	14.8	14.6	13.9	14.0	14.0
25 to 34	16.9	17.4	12.9	14.2	12.9	13.5
35 to 44	15.0	15.1	15.4	16.0	12.0	12.9
45 to 54	9.7	10.1	13.4	13.4	13.8	14.1
55 to 64	8.0	8.5	8.7	8.6	12.7	12.3
65 +	10.7	12.6	11.8	12.4	14.0	13.7

See Technical Appendix for information on population sources.

Population Section

Table P-2 Population Number by Age and County

New Mexico, 2012

Both Sexes

	Less													
	All ages	than 1	1-4	5-14	15-17	18-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
New Mexico	2,091,432	29,104	116,447	286,237	90,215	60,137	142,305	270,260	250,021	288,110	265,593	168,771	91,352	32,881
County														
Bernalillo	675,550	9,174	36,697	88,644	27,787	18,525	50,476	97,611	85,020	93,359	82,338	47,819	26,988	11,112
Catron	3,745	29	121	304	96	63	92	268	241	546	878	722	301	85
Chaves	66,120	1,057	4,227	10,053	3,333	2,222	4,250	8,042	7,433	8,463	7,562	4,944	3,214	1,320
Cibola	27,353	375	1,499	3,665	1,152	768	1,795	3,694	3,529	3,939	3,316	2,115	1,163	344
Colfax	13,518	140	561	1,524	494	330	605	1,359	1,392	2,059	2,244	1,612	876	323
Curry	49,596	874	3,497	7,571	2,138	1,426	4,028	7,449	5,928	6,125	4,840	3,123	1,908	689
De Baca	1,991	21	89	254	64	42	55	200	176	295	330	251	152	63
Dona Ana	214,878	3,078	12,311	30,378	10,855	7,237	18,370	27,159	23,603	26,176	23,543	18,831	10,354	2,984
Eddy	54,815	776	3,102	7,864	2,376	1,584	3,208	6,937	6,471	7,739	6,940	4,162	2,594	1,062
Grant	29,389	349	1,397	3,493	1,092	728	1,495	2,840	2,778	3,823	4,823	3,685	2,141	745
Guadalupe	4,664	51	211	564	172	113	269	660	551	743	587	416	253	73
Harding	695	5	22	44	17	11	16	49	55	106	159	96	81	35
Hidalgo	4,807	64	255	650	224	149	273	463	550	692	658	456	278	95
Lea	66,322	1,211	4,843	10,487	3,005	2,003	4,641	9,672	8,128	8,468	6,679	3,960	2,435	790
Lincoln	20,578	193	772	2,155	670	447	818	1,775	2,017	3,182	3,845	2,952	1,386	368
Los Alamos	18,219	193	772	2,540	683	455	500	1,745	2,332	3,331	2,862	1,514	885	407
Luna	25,138	361	1,445	3,581	1,143	762	1,433	2,639	2,643	2,983	3,152	2,745	1,743	508
McKinley	73,426	1,252	5,010	12,171	4,144	2,763	5,546	9,163	8,663	9,743	7,587	4,318	2,313	752
Mora	4,822	47	195	582	176	116	210	439	548	717	845	550	312	86
Otero	64,768	957	3,827	8,582	2,606	1,737	4,886	8,500	7,205	8,587	7,798	5,822	3,319	943
Quay	8,852	101	403	1,054	342	228	356	860	957	1,275	1,405	1,085	596	190
Rio Arriba	40,257	536	2,146	5,415	1,671	1,114	2,420	4,624	4,886	5,944	5,546	3,527	1,827	602
Roosevelt	20,577	326	1,302	2,869	1,170	780	2,304	2,775	2,198	2,299	2,023	1,363	814	354
Sandoval	135,950	1,844	7,375	20,213	5,664	3,776	7,120	16,146	17,399	19,929	18,410	11,096	5,069	1,909
San Juan	128,805	2,149	8,595	20,475	6,078	4,052	8,646	17,706	15,104	17,368	14,452	8,127	4,395	1,657
San Miguel	29,160	315	1,261	3,406	1,443	962	1,976	3,157	3,165	4,441	4,342	2,890	1,368	433
Santa Fe	146,559	1,642	6,568	16,753	5,005	3,337	7,498	16,719	18,189	22,036	24,731	14,920	6,773	2,389
Sierra	11,908	111	444	994	313	209	469	796	994	1,600	2,230	2,075	1,252	420
Socorro	17,764	236	943	2,302	868	579	1,444	2,094	1,915	2,460	2,333	1,570	765	254
Taos	33,122	359	1,435	3,747	1,113	742	1,547	3,348	3,906	5,025	5,853	3,716	1,720	613
Torrance	16,187	180	718	2,183	681	454	827	1,661	1,902	2,521	2,597	1,570	675	217
Union	4,535	48	199	479	160	105	239	667	581	694	549	434	266	113
Valencia	77,361	1,052	4,207	11,241	3,479	2,319	4,490	9,043	9,561	11,441	10,137	6,307	3,135	949

See Technical Appendix for information on population sources.

Population Section

Table P-3 Population Number by Age and County
New Mexico, 2012
Males

	All Ages	Less												
		than 1	1-4	5-14	15-17	18-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
New Mexico	1,032,728	14,874	59,509	145,481	46,198	30,794	72,816	137,797	124,710	140,928	127,739	79,450	40,689	11,743
County														
Bernalillo	330,545	4,714	18,855	44,939	14,078	9,386	25,320	49,218	42,427	45,420	39,084	21,932	11,377	3,795
Catron	1,961	15	64	166	49	32	45	145	134	261	452	401	163	34
Chaves	32,691	544	2,176	5,124	1,788	1,192	2,148	4,040	3,603	4,230	3,612	2,356	1,387	492
Cibola	13,897	187	749	1,891	576	384	872	2,007	1,965	2,012	1,647	954	526	130
Colfax	6,894	73	293	777	273	182	316	708	747	1,038	1,136	822	413	115
Curry	24,939	442	1,769	3,877	1,118	745	2,190	3,915	2,957	3,048	2,385	1,457	808	229
De Baca	975	10	40	129	32	21	19	96	86	150	174	130	68	21
Dona Ana	104,705	1,561	6,244	15,331	5,480	3,653	9,288	13,726	11,194	12,365	11,198	8,656	4,849	1,159
Eddy	27,430	408	1,633	3,946	1,239	826	1,653	3,607	3,292	3,862	3,483	1,983	1,152	347
Grant	14,433	184	735	1,839	575	383	746	1,391	1,331	1,825	2,288	1,840	1,024	273
Guadalupe	2,643	27	111	297	99	65	148	436	344	416	339	197	130	35
Harding	368	3	11	21	8	5	8	30	29	61	82	47	47	17
Hidalgo	2,412	33	132	337	115	77	131	240	273	341	330	233	139	33
Lea	33,905	612	2,446	5,373	1,521	1,014	2,397	5,246	4,229	4,443	3,378	1,868	1,077	301
Lincoln	10,113	102	408	1,103	359	239	406	910	974	1,505	1,802	1,437	701	166
Los Alamos	9,173	97	389	1,301	360	240	256	914	1,150	1,654	1,478	757	416	161
Luna	12,530	184	738	1,844	564	376	734	1,367	1,309	1,426	1,536	1,373	855	225
McKinley	35,530	642	2,570	6,115	2,075	1,384	2,837	4,534	4,233	4,574	3,486	1,878	933	269
Mora	2,470	24	99	314	93	61	116	216	268	373	422	291	148	46
Otero	32,820	493	1,970	4,367	1,394	929	2,732	4,477	3,692	4,216	3,845	2,823	1,539	342
Quay	4,367	54	216	549	183	122	185	412	475	617	670	538	278	68
Rio Arriba	19,886	262	1,046	2,713	874	582	1,218	2,355	2,457	2,984	2,694	1,701	802	198
Roosevelt	10,374	174	697	1,463	586	391	1,196	1,445	1,142	1,133	991	662	367	129
Sandoval	66,370	952	3,807	10,325	2,883	1,922	3,602	7,908	8,502	9,686	8,653	5,272	2,183	677
San Juan	63,972	1,081	4,323	10,391	3,124	2,083	4,382	9,045	7,537	8,656	7,091	3,775	1,929	553
San Miguel	14,559	155	619	1,745	744	496	1,040	1,637	1,555	2,246	2,127	1,440	610	145
Santa Fe	71,400	822	3,286	8,415	2,533	1,688	3,974	8,600	9,159	10,573	11,572	6,934	3,045	798
Sierra	5,997	57	226	523	158	106	240	402	489	800	1,095	1,064	644	194
Socorro	9,131	118	470	1,175	473	315	833	1,117	953	1,197	1,215	798	377	90
Taos	16,305	182	729	1,974	585	390	819	1,658	1,984	2,412	2,822	1,783	751	215
Torrance	8,350	86	343	1,116	367	245	477	892	1,037	1,237	1,324	815	332	81
Union	2,636	26	109	261	88	58	157	479	384	421	291	209	111	42
Valencia	38,947	552	2,210	5,743	1,805	1,203	2,331	4,625	4,801	5,744	5,036	3,025	1,507	365

See Technical Appendix for information on population sources.

Population Section

Table P-4 Population Number by Age and County
New Mexico, 2012
Females

	Less													
	All Ages	than 1	1-4	5-14	15-17	18-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
New Mexico	1,058,704	14,230	56,937	140,756	44,017	29,343	69,489	132,463	125,311	147,182	137,854	89,321	50,663	21,138
County														
Bernalillo	345,005	4,460	17,842	43,705	13,709	9,139	25,155	48,393	42,594	47,938	43,254	25,888	15,611	7,316
Catron	1,784	14	57	137	47	31	47	124	107	285	426	321	138	50
Chaves	33,429	513	2,050	4,929	1,546	1,030	2,102	4,002	3,830	4,234	3,950	2,589	1,827	828
Cibola	13,456	188	750	1,774	576	384	923	1,687	1,564	1,927	1,669	1,161	637	214
Colfax	6,625	67	268	748	222	148	288	650	644	1,022	1,108	790	463	208
Curry	24,657	432	1,728	3,694	1,020	680	1,839	3,534	2,971	3,077	2,455	1,666	1,101	459
De Baca	1,016	12	49	125	32	21	36	104	90	144	156	122	83	42
Dona Ana	110,173	1,517	6,067	15,047	5,375	3,583	9,082	13,433	12,410	13,811	12,345	10,174	5,504	1,825
Eddy	27,384	367	1,470	3,917	1,137	758	1,555	3,329	3,179	3,878	3,457	2,179	1,442	715
Grant	14,957	166	662	1,655	517	345	749	1,448	1,448	1,997	2,535	1,845	1,117	473
Guadalupe	2,022	24	101	268	73	48	121	224	206	327	249	219	123	38
Harding	327	3	11	23	8	5	8	19	26	45	77	49	34	18
Hidalgo	2,394	31	124	314	109	73	143	222	278	351	328	222	139	62
Lea	32,417	599	2,397	5,114	1,483	989	2,244	4,427	3,899	4,025	3,301	2,092	1,359	489
Lincoln	10,466	91	363	1,052	311	208	412	865	1,042	1,677	2,042	1,515	685	202
Los Alamos	9,047	96	383	1,239	324	216	245	831	1,182	1,676	1,384	756	469	245
Luna	12,608	177	708	1,737	579	386	699	1,272	1,334	1,557	1,616	1,372	888	283
McKinley	37,896	610	2,440	6,056	2,069	1,379	2,710	4,629	4,430	5,169	4,101	2,440	1,380	483
Mora	2,352	23	96	268	83	55	94	223	279	344	423	260	164	40
Otero	31,948	464	1,857	4,215	1,212	808	2,154	4,023	3,513	4,371	3,952	2,999	1,780	601
Quay	4,485	47	188	505	159	106	171	448	482	658	735	547	318	122
Rio Arriba	20,371	275	1,099	2,702	797	531	1,202	2,269	2,429	2,960	2,852	1,826	1,024	405
Roosevelt	10,203	151	605	1,406	584	390	1,107	1,330	1,056	1,166	1,032	701	447	225
Sandoval	69,580	892	3,568	9,888	2,781	1,854	3,519	8,238	8,897	10,243	9,757	5,824	2,886	1,232
San Juan	64,833	1,068	4,272	10,084	2,954	1,969	4,264	8,661	7,566	8,712	7,361	4,353	2,465	1,104
San Miguel	14,601	161	643	1,661	699	466	936	1,520	1,610	2,195	2,215	1,450	758	288
Santa Fe	75,159	820	3,282	8,338	2,473	1,648	3,524	8,119	9,030	11,463	13,159	7,985	3,728	1,591
Sierra	5,911	54	217	472	154	103	229	394	506	800	1,135	1,011	608	226
Socorro	8,634	118	473	1,127	395	263	611	978	963	1,263	1,118	772	388	164
Taos	16,817	176	706	1,773	528	352	728	1,690	1,921	2,614	3,031	1,932	968	398
Torrance	7,837	94	375	1,067	314	209	351	769	864	1,284	1,273	756	343	137
Union	1,899	22	90	217	72	48	82	188	198	272	259	225	155	71
Valencia	38,414	499	1,997	5,498	1,674	1,116	2,159	4,419	4,760	5,696	5,100	3,283	1,628	584

See Technical Appendix for information on population sources.

Population Section

Table P-5 Population Number by Race/Ethnicity, Sex, and Age
New Mexico, 2012 and United States, 2012

	Less than													
	All Ages	1	1-4	5-14	15-17	18-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
New Mexico														
All Races														
Male	1,032,728	14,874	59,509	145,481	46,198	30,794	72,816	137,797	124,710	140,928	127,739	79,450	40,689	11,743
Female	1,058,704	14,230	56,937	140,756	44,017	29,343	69,489	132,463	125,311	147,182	137,854	89,321	50,663	21,138
Both Sexes	2,091,432	29,104	116,447	286,237	90,215	60,137	142,305	270,260	250,021	288,110	265,593	168,771	91,352	32,881
American Indian or Alaska Native														
Male	88,308	1,572	6,288	15,795	5,541	3,694	7,202	12,566	10,412	11,667	7,358	3,948	1,767	496
Female	94,862	1,557	6,226	15,684	5,418	3,612	7,292	12,967	11,257	12,892	8,979	5,373	2,666	937
Both Sexes	183,169	3,128	12,514	31,480	10,959	7,306	14,495	25,534	21,669	24,559	16,337	9,321	4,434	1,433
Asian or Pacific Islander														
Male	14,190	182	726	1,871	698	465	1,296	2,656	1,979	1,836	1,395	741	275	71
Female	16,911	168	674	1,898	670	447	1,317	3,064	2,552	2,315	2,119	1,029	540	118
Both Sexes	31,101	350	1,400	3,769	1,368	912	2,612	5,720	4,531	4,151	3,514	1,770	815	189
Black or African American														
Male	23,667	345	1,379	3,496	1,298	866	2,046	3,609	3,072	3,132	2,505	1,265	516	137
Female	19,159	308	1,232	3,176	929	619	1,415	2,451	2,232	2,746	1,964	1,115	719	251
Both Sexes	42,825	653	2,612	6,673	2,228	1,485	3,461	6,060	5,304	5,878	4,469	2,380	1,235	389
Hispanic														
Male	480,143	8,845	35,387	81,651	23,328	15,549	36,619	68,039	56,720	69,974	43,009	25,065	12,682	3,275
Female	489,274	8,525	34,111	79,489	22,852	15,233	35,675	66,713	56,740	72,290	46,072	29,266	16,548	5,760
Both Sexes	969,417	17,370	69,498	161,140	46,179	30,783	72,294	134,752	113,460	142,265	89,081	54,331	29,230	9,036
White														
Male	426,421	3,931	15,729	42,668	15,332	10,220	25,653	50,927	52,528	54,318	73,471	48,431	25,449	7,763
Female	438,498	3,672	14,694	40,509	14,148	9,431	23,789	47,267	52,529	56,939	78,720	52,538	30,190	14,071
Both Sexes	864,919	7,603	30,423	83,176	29,481	19,652	49,442	98,194	105,057	111,257	152,192	100,969	55,638	21,835
United States														
All Races														
Male	154,492,067	2,016,738	8,199,397	21,026,407	6,447,066	4,515,795	11,549,456	21,338,792	20,173,607	21,806,870	18,602,894	11,202,862	5,648,150	1,964,033
Female	159,421,973	1,926,339	7,856,870	20,118,347	6,136,924	4,260,917	11,033,747	20,970,529	20,342,813	22,461,868	19,983,308	12,782,530	7,624,484	3,923,297
Both Sexes	313,914,040	3,943,077	16,056,267	41,144,754	12,583,990	8,776,712	22,583,203	42,309,321	40,516,420	44,268,738	38,586,202	23,985,392	13,272,634	5,887,330

See Technical Appendix for information on race/ethnicity and population sources.

Population Section

**Table P-6 Population Number by Race/Ethnicity, County, and Health Region
New Mexico, 2012**

	All Races	American Indian or Alaska Native	Asian or Pacific Islander	Black or African American	Hispanic	White
New Mexico	2,091,432	183,169	31,101	42,825	969,417	864,919
County						
Bernalillo	675,550	28,243	17,065	19,491	324,302	286,449
Catron	3,745	98	8	35	716	2,888
Chaves	66,120	568	451	1,301	34,447	29,352
Cibola	27,353	10,882	162	276	10,002	6,031
Colfax	13,518	157	57	61	6,415	6,828
Curry	49,596	384	703	3,150	19,622	25,737
De Baca	1,991	12	2	18	778	1,181
Dona Ana	214,878	1,773	2,478	3,240	139,652	67,736
Eddy	54,815	518	367	730	24,269	28,930
Grant	29,389	247	144	254	14,221	14,524
Guadalupe	4,664	55	62	72	3,729	747
Harding	695	3	0	3	299	390
Hidalgo	4,807	15	27	34	2,721	2,010
Lea	66,322	564	348	2,643	33,950	28,817
Lincoln	20,578	452	73	126	6,203	13,724
Los Alamos	18,219	116	1,075	156	2,755	14,117
Luna	25,138	157	125	237	15,423	9,195
McKinley	73,426	54,352	624	569	9,806	8,075
Mora	4,822	18	12	23	3,904	865
Otero	64,768	3,988	987	2,406	22,358	35,030
Quay	8,852	79	85	109	3,772	4,807
Rio Arriba	40,257	5,768	148	150	28,742	5,449
Roosevelt	20,577	215	185	359	8,210	11,609
Sandoval	135,950	17,046	2,237	3,043	47,674	65,950
San Juan	128,805	46,965	609	977	24,645	55,609
San Miguel	29,160	261	204	392	22,420	5,883
Santa Fe	146,559	3,609	1,837	1,209	74,425	65,478
Sierra	11,908	156	56	83	3,372	8,240
Socorro	17,764	1,933	225	194	8,637	6,776
Taos	33,122	1,885	216	186	18,611	12,224
Torrance	16,187	325	83	213	6,361	9,204
Union	4,535	51	24	86	1,808	2,567
Valencia	77,361	2,273	422	999	45,168	28,499
Health Region						
Northwest	229,584	112,200	1,395	1,822	44,453	69,714
Northeast	295,552	11,924	3,635	2,339	163,108	114,547
Metro	905,047	47,887	19,807	23,746	423,505	390,102
Southeast	288,851	2,792	2,214	8,436	131,251	144,158
Southwest	372,398	8,367	4,050	6,483	207,100	146,399

See Technical Appendix for information on race/ethnicity, health regions, and population sources.

Population Section

**Table P-7 Births and Deaths by City
Population, 2010 and 2012; Births and Deaths, 2012**

City	Population			Births	Deaths	City	Population			Births	Deaths
	2010 Population Estimate	2012 Population Estimate	Percent Change 2011- 2012				2010 Population Estimate	2012 Population Estimate	Percent Change 2011- 2012		
Alamogordo	30,403	31,500	3.6	511	361	Lake Arthur	436	437	0.2	12	5
Albuquerque	545,852	555,417	1.8	8,150	4,887	Las Cruces	97,618	101,047	3.5	1,840	1,094
Artesia	11,301	11,365	0.6	255	146	Las Vegas	13,753	13,529	-1.6	265	204
Aztec	6,763	6,683	-1.2	211	160	Logan	1,042	1,008	-3.3	12	14
Bayard	2,328	2,316	-0.5	33	19	Lordsburg	2,797	2,740	-2.0	44	31
Belen	7,269	7,255	-0.2	256	226	Los Lunas	14,835	15,168	2.2	535	260
Bernalillo	8,320	8,413	1.1	121	81	Los Ranchos De Albuquerque	6,024	6,087	1.0	1	14
Bloomfield	8,112	7,968	-1.8	223	93	Loving	1,413	1,411	-0.1	25	19
Bosque Farms	3,904	3,889	-0.4	42	41	Lovington	11,009	11,275	2.4	238	116
Capitan	1,489	1,470	-1.3	18	21	Magdalena	938	926	-1.3	19	18
Carlsbad	26,138	26,697	2.1	506	355	Maxwell	254	244	-3.9	3	1
Carrizozo	996	984	-1.2	11	14	Melrose	651	662	1.7	11	7
Causey	104	106	1.9	0	0	Mesilla	2,196	1,913	-12.9	6	14
Chama	1,022	1,025	0.3	17	18	Milan	3,245	3,251	0.2	35	15
Cimarron	1,021	978	-4.2	5	9	Moriarty	1,910	1,868	-2.2	77	73
Clayton	2,980	2,910	-2.3	34	35	Mosquero	93	95	2.2	2	2
Cloudcroft	674	697	3.4	6	8	Mountainair	928	908	-2.2	19	27
Clovis	37,775	39,197	3.8	876	364	Pecos	1,392	1,367	-1.8	39	24
Columbus	1,664	1,662	-0.1	40	18	Portales	12,280	12,723	3.6	264	149
Corona	172	170	-1.2	3	3	Questa	1,770	1,760	-0.6	33	16
Corrales	8,329	8,453	1.5	32	60	Raton	6,885	6,607	-4.0	77	100
Cuba	731	740	1.2	59	13	Red River	477	475	-0.4	5	2
Deming	14,855	14,793	-0.4	383	255	Reserve	289	284	-1.7	7	11
Des Moines	143	137	-4.2	6	0	Rio Rancho	87,521	90,818	3.8	1,004	588
Dexter	1,266	1,273	0.6	68	36	Roswell	48,366	48,477	0.2	846	559
Dora	133	135	1.5	1	1	Roy	234	238	1.7	3	3
Eagle Nest	290	278	-4.1	3	2	Ruidoso	8,029	8,005	-0.3	80	85
Edgewood	3,735	3,779	1.2	105	96	Ruidoso Downs	2,815	2,739	-2.7	50	23
Elida	197	201	2.0	0	10	San Jon	216	210	-2.8	3	3
Encino	82	80	-2.4	0	2	Santa Clara	1,686	1,684	-0.1	25	29
Espanola	10,224	10,240	0.2	220	142	Santa Fe	67,947	69,204	1.8	1,238	924
Estancia	1,655	1,632	-1.4	29	15	Santa Rosa	2,848	2,802	-1.6	35	23
Eunice	2,922	2,993	2.4	50	29	San Ysidro	193	195	1.0	3	2
Farmington	45,877	45,854	-0.1	870	361	Silver City	10,315	10,273	-0.4	199	201
Floyd	133	135	1.5	0	4	Socorro city	9,051	8,906	-1.6	123	90
Folsom	56	57	1.8	2	1	Springer	1,047	1,006	-3.9	10	30
Fort Sumner	1,031	983	-4.7	11	35	Sunland Park	14,106	14,776	4.7	218	61
Gallup	21,678	22,088	1.9	376	160	Taos	5,716	5,676	-0.7	127	105
Grady	107	109	1.9	3	4	Tatum	798	817	2.4	13	12
Grants	9,182	9,224	0.5	196	106	Texico	1,130	1,148	1.6	31	7
Grenville	38	37	-2.6	2	2	Tijeras	541	547	1.1	61	64
Hagerman	1,257	1,259	0.2	20	11	Truth or Consequences	6,475	6,411	-1.0	59	146
Hatch	1,648	1,639	-0.5	62	19	Tucumcari	5,363	5,204	-3.0	102	92
Hobbs	34,122	35,007	2.6	819	353	Tularosa	2,842	2,943	3.6	44	63
Hope	105	105	0.0	4	6	Vaughn	446	437	-2.0	4	2
House	68	66	-2.9	2	1	Virden	152	149	-2.0	3	5
Hurley	1,297	1,292	-0.4	27	23	Wagon Mound	314	303	-3.5	2	9
Jal	2,047	2,096	2.4	31	21	Willard	253	247	-2.4	2	0
Jemez Springs	250	253	1.2	5	8	Williamsburg	449	447	-0.4	3	16

See Technical Appendix for information on population sources.

Population Section

Table P-8 Summary of Health Statistics Trends

New Mexico, 1960-2012

Year	Population	Births	Deaths	Maternal Mortality		Infant Mortality				Fetal Mortality	
				Number	Rate	All Infants Deaths	Mortality Rate	Neonatal Rate	Post-neonatal Rate	Number	Rate
2012	2,091,432	26,992	16,640	1	3.7	186	6.9	4.7	2.2	69	2.5
2011	2,075,056	27,251	16,245	10	36.7	143	5.2	3.2	2.0	68	2.5
2010	2,065,171	27,795	15,866	4	14.4	155	5.6	3.4	2.2	75	2.7
2009	2,036,112	28,873	15,392	4	13.9	145	5.0	3.0	2.0	90	3.1
2008	2,013,046	30,156	15,400	6	19.9	154	5.1	3.0	2.1	89	2.9
2007	1,989,979	30,605	15,400	4	13.1	188	6.1	3.8	2.4	82	2.7
2006	1,966,876	29,918	15,231	1	3.3	170	5.7	3.6	2.1	69	2.3
2005	1,943,810	28,822	14,866	2	6.9	175	6.1	3.6	2.5	84	2.9
2004	1,920,743	28,355	14,197	4	14.1	178	6.3	3.2	2.2	78	2.7
2003	1,897,640	27,799	14,493	4	14.4	150	5.4	3.2	2.2	89	3.2
2002	1,874,575	27,708	14,114	6	21.7	168	6.1	4.2	1.9	89	3.2
2001	1,851,512	27,101	14,016	8	29.5	174	6.4	4.0	2.4	64	2.4
2000	1,828,560	27,206	13,384	8	29.4	180	6.6	3.7	2.9	104	3.8
1999	1,808,082	27,133	13,433	5	18.4	185	6.8	3.9	2.9	90	3.3
1998	1,793,484	27,294	12,858	3	11.0	194	7.1	4.3	2.8	84	3.1
1997	1,774,839	26,844	12,613	1	3.7	165	6.1	3.3	2.9	103	3.8
1996	1,752,326	27,216	12,456	3	11.0	169	6.2	3.9	2.4	84	3.1
1995	1,720,394	26,914	12,500	3	11.1	162	6.0	4.0	2.0	106	3.9
1994	1,682,398	27,585	12,106	5	18.1	228	8.3	4.9	3.4	108	3.9
1993	1,636,453	27,831	11,689	2	7.2	233	8.4	4.6	3.8	108	3.9
1992	1,595,442	27,910	11,130	2	7.2	209	7.5	3.9	3.6	104	3.7
1991	1,555,305	27,783	11,225	3	10.8	224	8.1	4.9	3.1	100	3.6
1990	1,521,574	27,318	10,549	9	32.9	243	8.9	5.2	3.7	123	4.5
1989	1,503,901	27,265	10,473	5	18.3	232	8.5	5.4	3.1	111	4.1
1988	1,490,336	26,935	10,381	2	7.4	268	9.9	6.1	3.9	147	5.4
1987	1,478,519	27,246	10,324	1	3.7	218	8.0	5.0	3.0	121	4.4
1986	1,462,728	27,281	10,007	3	11.0	254	9.3	5.4	4.0	111	4.1
1985	1,438,360	27,449	9,637	3	10.9	291	10.6	6.2	4.4	157	5.7
1984	1,416,719	27,350	9,504	5	18.3	263	9.6	6.0	3.6	186	6.8
1983	1,394,362	27,508	9,138	3	10.9	274	10.0	5.5	4.5	192	6.9
1982	1,363,822	27,630	9,186	4	14.5	316	11.4	6.7	4.7	202	7.3
1981	1,332,747	26,565	8,668	2	7.5	256	9.6	5.7	3.9	180	6.7
1980	1,303,303	26,589	9,032	3	11.3	293	11.0	7.0	4.0	204	7.6
1979	1,283,000	24,821	8,617	10	40.3	352	14.2	8.8	5.4	206	8.2
1978	1,254,000	23,907	8,331	2	8.4	330	13.8	8.7	5.1	199	8.3
1977	1,227,000	23,100	8,223	2	8.7	333	14.4	9.8	4.6	201	8.6
1976	1,196,000	22,180	8,204	3	13.5	347	15.6	10.0	5.6	196	8.8
1975	1,164,000	21,078	8,003	2	9.5	357	16.9	11.0	6.0	169	8.0
1974	1,130,000	21,339	8,029	10	46.9	391	18.3	12.2	6.1	184	8.5
1973	1,104,000	20,852	8,139	2	9.6	421	20.2	12.9	7.3	211	10.0
1972	1,078,000	20,813	7,877	5	24.0	403	19.4	14.4	5.0	248	11.8
1971	1,053,000	22,205	7,638	5	22.5	460	20.7	15.0	5.8	254	11.3
1970	1,017,055	22,004	7,411	6	27.3	463	21.0	15.0	6.0	245	11.0
1969	1,011,000	21,543	7,180	5	23.2	508	23.6	16.1	7.5	266	12.2
1968	994,000	20,346	7,128	7	34.4	487	23.9	16.8	7.2	264	12.8
1967	1,000,000	21,243	6,897	4	18.8	527	24.8	16.1	8.7	249	11.6
1966	1,007,000	22,363	6,971	9	40.2	601	26.9	16.7	10.2	278	12.3
1965	1,012,000	24,352	6,801	10	41.1	657	27.0	17.8	9.2	326	13.2
1964	1,006,000	26,862	6,902	14	52.1	781	29.1	19.2	9.9	349	12.8
1963	989,000	27,820	6,837	10	35.9	848	30.5	20.0	10.5	371	13.2
1962	979,000	29,226	6,507	16	54.7	877	30.0	19.5	10.5	409	13.8
1961	965,000	30,009	6,344	19	63.3	880	29.3	20.0	9.3	370	12.2
1960	951,023	30,747	6,503	13	42.3	1,022	33.2	20.2	13.0	414	13.3

Maternal mortality rates are per 100,000 live births; infant mortality rates are per 1,000 live births; and fetal mortality rates are per 1,000 live births + fetal deaths. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution. See Technical Appendix for information on rates and population sources.

Intentionally Left Blank

Natality Section




NATALITY SECTION

BIRTH NUMBERS AND RATES

There were 26,992 births to New Mexico resident mothers in 2012, translating to a birth rate of 12.9 births per 1,000 population (Table N-1). New Mexico’s birth rate has generally declined since 1994 when the birth rate was 16.7. In 2011, the latest year for which United States data are available, the national birth rate (12.7) was slightly lower than the New Mexico 2012 birth rate. These are the lowest birth rates on record for both the United States and New Mexico.

The fertility rate is calculated as the number of births per 1,000 females ages 15-44 years. New Mexico’s fertility rate decreased 7.7% between 2011 (68.5) and 2012 (63.2). New Mexico’s birth rate for mothers in the 20-24 year-old age category has decreased from 137.4 in 2008 to 113.2 in 2012. However, the 2012 New Mexico rate was 24.6 % higher than the 2011 United States rate (85.3) for the same age group (Table N-3).

**Figure N-1, Birth Rates
New Mexico, 1994-2012, and United States, 1994-2011**



Population note: Birth rates for years 2001-2010 differ from those shown in reports published in previous years. This reflects adjustments to 2001-2010 population estimates following the 2010 Census. See the *Technical Appendix* for more information.


RACE AND ETHNICITY

NOTE: Birth rates by race and ethnicity differ from those shown in reports published in previous years. This reflects a revision to the Department of Health’s standard definition for presentation of race/ethnicity as a single construct. (See technical appendix for more information.)

Natality Section


In 2012, Hispanic mothers had the highest percent of births (54.0%) (Table N-4(b)). Since 2000, American Indians or Alaska Natives had the highest birth rates among the state's racial/ethnic groups (Figure N-2).

**Figure N-2, Birth Rates by Race/Ethnicity
New Mexico, 1994-2012**



AGE OF MOTHER

**Figure N-3, Birth Rates by Mother's Age
New Mexico, 1990, 2000, and 2012**




Natality Section

Between 1990 and 2012 birth rates decreased for women ages 15-29 years and increased for ages 30-44 years. The most significant birth rate decreases were among 15-19 year olds (42.7%). The highest rates of birth rate decreases were among women ages 40-44 years (42.6%) (Figure N-3).

Teen Mothers

**Figure N-4, Teen (15-17 Years and 18-19 Years) Birth Rates
New Mexico, 1995-2012, and United States, 1995-2011**




Birth rates for New Mexico teens ages 15-17 years decreased 55.6% and rates for teens ages 18-19 years decreased 32.1% between 1995 and 2012. However, New Mexico's teen birth rates continue to be higher than national rates (Figure N-4).

BIRTHS TO SINGLE MOTHERS


From 1985 to 2012 in New Mexico, the percent of births to single mothers increased from 26.4% to 51.9%. Nationally, 40.7% of births were to single women in 2011, compared to 51.9% in New Mexico in 2012 (Figure N-5). The percent of births to single women in 2012 was highest for American Indian or Alaska Native women (78.8). Hispanic women (58.2) and Black or African American women (53.4) were the next highest. The increase in the percent of single mothers was highest among White women (70.1%) and second highest among Asian or Pacific Islander White women (45.7%) between 1990 and 2012. Black or African American women were the only group to have had a decrease in the proportion of births to single mothers (4.1%) (Figure N-6).

Natality Section

**Figure N-5, Percent of Births to Single Mothers
New Mexico, 1985-2012, and United States, 1985-2011**




**Figure N-6, Percent of Births to Single Mothers by Race/Ethnicity
New Mexico, 1990, 2000 and 2012**



AI=American Indian; AN=Alaska Native; PI=Pacific Islander; and AA=African American.
See Technical Appendix for information on race/ethnicity.

Natality Section

**Figure N-7, Percent of Births to Single Mothers by Age
New Mexico, 1990, 2000 and 2012**




BIRTH ORDER

Birth order refers to the order in which the child is born to the mother: first child, second child, etc. A population that has a high proportion of second or higher births may have greater fertility rates than a population where most newborns are first born.

In 2012, 37.1% of all births in New Mexico were first born (Figure N-8). This is up from 25.3% in 1960. In contrast, 6.1% of the births in 2012 were fifth born or later, compared to 21.8% in 1960.

Natality Section


**Figure N-8, Percent of Birth Order
New Mexico, 1960-2012**



MULTIPLE BIRTHS

The multiple birth rate, defined as the number of twins, triplets, or higher multiple births per 1,000 total live births, was 25.7 in 2012 in New Mexico (Figure N-9). The 2012 multiple birth rate in New Mexico was 25.7% lower than the 2011 United States multiple birth rate (34.6).¹

**Figure N-9, Number of Births in Single and Multiple Deliveries by Year
New Mexico, 1990-2012**



¹ Martin JA, Hamilton BE, et al. Births: Final data for 2010. National Vital Statistics Report, Vol. 61, No. 1. Hyattsville, MD: National Center for Health Statistics. 2012.

Natality Section


The number of singleton births decreased in New Mexico by 1.8% between 1990 and 2012; whereas, the number of multiple births increased 29.9%. The percent of multiple births to total births increased from 2.0% in 1990 to 2.6% in 2012 (Figure N-9).

BIRTHWEIGHT

Low birthweight	Less than 2,500 grams
Normal birthweight	2,500 – 3,999 grams
High birthweight	Equal to or greater than 4,000 grams

Birthweight is an important predictor of infant survival. Low birthweight is a factor in approximately two-thirds of infant deaths nationally.² Low birthweight newborns are also more likely to have serious health problems and increased risks of long-term disabilities.


**Figure N-10, Percent of Births with Low Birthweight
New Mexico, 1999-2012, and United States, 1999-2011**



² March of Dimes. 2006. "Professional and Researchers: Low Birth Weight," <http://www.marchofdimes.com>.

Natality Section


**Figure N-11, Percent of Births with Low Birthweight by Mother's Race/Ethnicity
New Mexico, 2008-2012**



From 1999 to 2006, the percent of low birthweight infants generally increased in New Mexico and in the United States. The percent of low birthweight infants in the United States reached a high of 8.3% in 2006. The steady increase in low birthweight births in New Mexico went from 7.7% in 1999 to 8.8% in 2012. (Figure N-10).

In 2012, Black or African American women had the highest percentages of low birthweight infants (13.9%), followed by Asians or Pacific Islanders (13.3%) (Figure N-11).

**Figure N-12, Percent of Births with Low Birthweight by Mother’s Age
New Mexico, 2010-2012, and United States, 2011**



Demographic factors associated with increased risk of low birthweight include mother’s age (19 years and younger or 35 years and older), marital status of the mother (single), and gestational age.³ For New Mexico and the nation, mothers 40 years of age and older showed the highest percentages of low birthweight births (Figure N-12).

GESTATIONAL AGE

Ninety percent of term births (born at 37-41 weeks gestation) were born with normal birthweight, while 64.4% of early preterm births were very low birthweight in 2012 (Figure N-14). Preterm birth is a leading cause of infant death and is associated with congenital neurological defects.

Preterm births were highest in Black or African American (15.5%) and American Indian (14.2%) populations in 2012 (Figure N-13).

The calculation for gestational age in New Mexico uses the United States National Center for Health Statistics (NCHS) methodology and measures the number of weeks from the date of the last normal menses to delivery. Clinical estimates of gestation are used instead of calculated gestation for unknown or “out of usual range” dates of last normal menses. (See *Technical Appendix* for additional information.)

³ Martin JA, Hamilton BE, Sutton PD. et al. Births: Final data for 2005. National Vital Statistics Report, Vol. 56, No. 6. Hyattsville, MD: National Center for Health Statistics. 2007.

Natality Section

Figure N-13
Percent of Preterm Births by Race/Ethnicity
New Mexico, 2009-2012





Figure N-14
Percent Gestational Age by Birthweight
New Mexico, 2012




PRENATAL CARE

There are two primary ways that New Mexico reports on prenatal care: the trimester prenatal care began and the Kessner Index. These measures rely heavily on when prenatal care was initiated.

Kessner Index

The traditional measure of prenatal care used in New Mexico has been a modified Kessner index. Levels of prenatal care are defined by using a combination of factors: the month prenatal care began and the number of prenatal visits made. A low level of care is defined as care that begins in the third trimester with less than five or no prenatal care visits. A high level of care is defined as care that began during the first trimester of pregnancy with nine or more prenatal care visits occurring during that period. Moderate care is defined as care that began during the first trimester with 5-8 prenatal visits, or care beginning in the fourth to sixth month of pregnancy with 5 or more visits.


**Figure N-15, Percent of Births by Level of Prenatal Care (Modified Kessner Index)
New Mexico, 2009-2012**



Nativity Section

The Revised Birth Certificate: Changes to the New Mexico Birth File, in the New Mexico Selected Health Statistics, Annual Report, 2008 at http://www.vitalrecordsnm.org/reports/2008_AR.pdf

**Figure N-16, Percent of Births with Low or No Prenatal Care by Mother's Age
New Mexico, 2012**




New Mexico mothers less than 18 years of age received the highest percentage of no or low levels of prenatal care (Figure N-16).

FACILITY AND ATTENDANT

Of all the births to New Mexico resident mothers in 2012, 98.1% occurred in hospitals. Medical doctors attended 66.3% of the births, and certified nurse midwives (CNMs) attended 27.0% of the births. The percent of births delivered by medical doctors decreased from 88.0% in 1990 to 66.3% in 2012. With this decrease, the number of births delivered by other types of attendants increased during this time period. Specifically, the percent of deliveries by CNMs increased from 10.6% in 1990 to 27.0% in 2012 (Table N-19)

Natality Section

**Figure N-17, Percent of Births Occurring in New Mexico by Attendant Type
New Mexico, 2003-2012**




METHOD OF DELIVERY

The percent of vaginal births decreased 6.6% since 1998, while the percent of cesarean births increased 43.1% (Figure N-18).

Generally, the younger the mother is, the higher the percent of vaginal births. For 15-17 year old mothers, 86.1% had vaginal deliveries (excluding vaginal after previous cesarean) while only 13.0% had cesarean sections; 77.9% of 20-24 year old mothers delivered vaginally (excluding vaginal after previous cesarean) and 15.3% delivered cesarean. For 35-39 year old mothers, 63.6% had vaginal deliveries (excluding vaginal after previous cesarean) while 23.6% had cesareans (Table N-20(b)).

Natality Section

**Figure N-18, Distribution of Delivery Method by Year
New Mexico, 1998-2012**



Natality Section

**Table N-1 Natality Characteristics by County and Health Region
New Mexico, 2012, and United States, 2011**

	Number of Live Births	Crude Rate	Percent	Fertility Rate	Sex Ratio		Male to Female
					Male	Female	
United States	3,953,590	12.7	100.0	63.2	2,024,052	1,929,538	1.05
New Mexico	26,992	12.9	100.0	67.4	13,815	13,177	1.05
County							
Bernalillo	8,379	12.4	31.0	60.3	4,297	4,082	1.05
Catron	22	5.9	0.1	61.8	11	11	1.00
Chaves	948	14.3	3.5	75.8	479	469	1.02
Cibola	429	15.7	1.6	83.6	219	210	1.04
Colfax	115	8.5	0.4	58.9	60	55	1.09
Curry	924	18.6	3.4	92.0	488	436	1.12
De Baca	12	6	0.0	42.4	6	6	1.00
Dona Ana	3,071	14.3	11.4	70.0	1,563	1,508	1.04
Eddy	796	14.5	2.9	79.9	401	395	1.02
Grant	322	11.0	1.2	71.4	158	164	0.96
Guadalupe	47	10.1	0.2	69.8	21	26	0.81
Harding	5	7.2	0.0	75.0	2	3	0.67
Hidalgo	59	12.3	0.2	71.6	26	33	0.79
Lea	1,161	17.5	4.3	89.0	609	552	1.10
Lincoln	196	9.5	0.7	69.1	100	96	1.04
Los Alamos	178	9.8	0.7	63.6	102	76	1.34
Luna	419	16.7	1.6	98.1	205	214	0.96
McKinley	1,239	16.9	4.6	81.4	628	611	1.03
Mora	41	8.5	0.2	55.8	27	14	1.93
Otero	888	13.7	3.3	75.8	435	453	0.96
Quay	120	13.6	0.4	87.8	45	75	0.60
Rio Arriba	610	15.2	2.3	84.4	292	318	0.92
Roosevelt	271	13.2	1.0	60.7	139	132	1.05
Sandoval	1,459	10.7	5.4	57.7	773	686	1.13
San Juan	1,862	14.5	6.9	73.3	941	921	1.02
San Miguel	352	12.1	1.3	67.3	178	174	1.02
Santa Fe	1,358	9.3	5.0	54.8	695	663	1.05
Sierra	91	7.6	0.3	65.6	50	41	1.22
Socorro	211	11.9	0.8	65.8	117	94	1.24
Taos	319	9.6	1.2	61.1	157	162	0.97
Torrance	160	9.9	0.6	63.8	94	66	1.42
Union	53	11.7	0.2	90.1	32	21	1.52
Valencia	873	11.3	3.2	61.8	464	409	1.13
Unknown County	2	--	0.0	--	1	1	1.00
Health Region							
Northwest	3,530	15.4	13.1	77.1	1,788	1,742	1.03
Northeast	3,078	10.4	11.4	62.4	1,566	1,512	1.04
Metro	10,871	12	40.3	60.1	5,628	5,243	1.07
Southeast	4,428	15.3	16.4	81.2	2,267	2,161	1.05
Southwest	5,083	13.6	18.8	72.5	2,565	2,518	1.02
Unknown Region	2	--	0.0	--	1	1	1.00

See Technical Appendix for information on rates, race/ethnicity, and health regions.

Natality Section

**Table N-2 Number of Births by Year, Mother's Age, and Race/Ethnicity
New Mexico, 2008-2012, and United States, 2008-2011**

Year	Mother's age group											Unknown/ Not Reported
	All Ages	10 to 14	15 to 17	18 to 19	20 to 24	25 to 29	30 to 34	35 to 39	40 to 44	45 to 49	50+	
United States - All Races												
2011	3,953,590	3,974	95,538	234,234	925,200	1,127,583	986,682	463,849	108,920	7,025	585	0
2010	3,999,386	4,497	109,173	258,505	951,688	1,133,713	962,170	464,870	107,045	7,154	571	0
2009	4,130,665	5,029	124,247	285,555	1,005,982	1,166,787	955,246	474,103	105,827	7,320	569	0
2008	4,247,694	5,764	135,664	299,094	1,052,184	1,195,774	956,716	488,875	105,973	7,109	541	0
New Mexico - All Races												
2012	26,992	50	998	2,267	7,867	7,503	5,346	2,367	547	29	5	13
2011	27,251	44	1,107	2,340	7,877	7,731	5,209	2,318	583	27	6	9
2010	27,795	46	1,287	2,578	8,185	7,570	5,088	2,425	570	39	6	1
2009	28,873	52	1,479	2,930	8,586	7,994	4,913	2,318	552	45	1	3
2008	30,156	63	1,597	2,940	9,272	8,096	5,151	2,388	593	48	4	4
American Indian or Alaska Native												
2012	3,277	12	127	347	1,063	875	536	251	62	4	0	0
2011	3,388	2	151	333	1,140	876	544	262	75	4	0	1
2010	3,415	9	163	360	1,086	858	555	319	62	3	0	0
2009	3,698	6	201	425	1,188	941	567	284	77	8	0	1
2008	3,741	7	206	433	1,236	940	557	279	73	10	0	0
Asian or Pacific Islander												
2012	556	0	1	12	63	146	192	124	16	1	0	1
2011	534	1	4	10	72	158	184	84	20	1	0	0
2010	546	0	4	11	61	170	168	113	19	0	0	0
2009	486	1	6	9	69	148	150	75	28	0	0	0
2008	534	0	9	10	70	149	177	93	25	1	0	0
Black or African American												
2012	459	1	9	40	148	119	98	34	9	1	0	0
2011	465	2	13	35	163	118	82	45	4	2	1	0
2010	488	0	21	46	176	112	92	31	8	2	0	0
2009	504	3	22	55	145	139	84	45	11	0	0	0
2008	489	1	16	58	182	127	68	29	6	2	0	0
Hispanic												
2012	14,580	35	717	1,430	4,629	3,937	2,473	1,084	261	7	2	5
2011	14,798	35	774	1,502	4,589	4,037	2,480	1,098	268	12	0	3
2010	15,173	32	927	1,673	4,827	3,985	2,360	1,092	264	13	0	0
2009	15,848	41	1,060	1,868	5,041	4,196	2,335	1,085	207	15	0	0
2008	16,620	50	1,148	1,904	5,453	4,235	2,503	1,063	240	18	4	2
White												
2012	7,901	2	135	422	1,906	2,365	2,005	852	192	16	2	4
2011	7,895	4	160	448	1,873	2,502	1,873	810	209	8	5	3
2010	8,025	5	169	473	1,986	2,407	1,889	856	212	21	6	1
2009	8,186	1	187	557	2,097	2,535	1,742	820	224	22	1	0
2008	8,630	4	209	528	2,287	2,607	1,823	913	241	17	0	1

See Technical Appendix for information on race/ethnicity.

Note: Other and unknown races or ages, if any, are included in the "All Race" or "All Ages" categories

For the 10 to 14 year old age group, U.S. data are for mothers under 15 years of age.

Natality Section

**Table N-3 Birth Rates by Mother's Age and Race/Ethnicity
New Mexico, 2008-2012, and United States, 2008-2011**

Year	Mother's age group									
	10 to 14	15 to 17	18 to 19	20 to 24	25 to 29	30 to 34	35 to 39	40 to 44	45 to 49	50+
United States - All Races										
2011	0.4	15.4	54.1	85.3	107.2	96.5	47.2	10.3	0.7	*
2010	0.4	17.3	58.2	90.0	108.3	96.5	45.9	10.2	0.7	*
2009	0.5	19.6	64.0	96.2	111.5	97.5	46.1	10.0	0.7	*
2008	0.6	21.1	68.2	101.8	115.0	99.4	46.8	9.9	0.7	*
New Mexico - All Races										
2012	0.7	22.7	77.3	113.2	109.1	83.9	37.8	8.7	0.4	0.0
2011	0.6	25.2	80.0	113.8	112.9	82.1	37.2	9.3	0.4	0.0
2010	0.7	29.3	88.1	117.5	110.5	80.7	39.3	9.1	0.5	0.0
2009	0.7	33.8	100.5	125.4	118.9	78.4	36.9	8.6	0.6	0.0
2008	0.9	36.6	101.2	137.4	122.2	82.6	37.5	9.2	0.7	0.0
American Indian or Alaska Native										
2012	1.4	23.4	96.1	145.8	129.3	86.5	43.4	11.3	0.6	0.0
2011	0.2	27.9	92.3	156.6	129.7	87.9	45.3	13.7	0.6	0.0
2010	1.1	30.2	99.9	148.8	127.6	90.8	55.9	11.3	0.4	0.0
2009	0.7	37.5	118.9	164.1	141.1	93.0	48.7	13.8	1.2	0.0
2008	0.8	38.8	122.2	171.7	141.8	91.6	47.0	13.0	1.5	0.0
Asian or Pacific Islander										
2012	0.0	1.5	26.9	47.8	93.2	128.2	88.5	13.9	0.9	0.0
2011	1.2	6.0	22.5	54.9	101.3	123.4	60.2	17.5	0.9	0.0
2010	0.0	6.0	24.8	46.5	109.2	113.4	81.9	16.5	0.0	0.0
2009	1.2	9.4	21.0	55.0	98.6	104.6	55.1	24.6	0.0	0.0
2008	0.0	14.5	24.2	58.3	102.5	127.2	69.4	22.2	0.9	0.0
Black or African American										
2012	0.6	9.7	64.6	104.6	89.6	87.3	30.0	8.2	0.7	0.0
2011	1.2	14.1	56.8	115.8	89.3	73.5	41.3	3.6	1.3	0.2
2010	0.0	22.8	74.9	124.8	84.9	83.2	28.8	7.3	1.3	0.0
2009	1.9	24.0	90.0	103.6	108.2	76.5	42.6	10.0	0.0	0.0
2008	0.6	17.5	95.2	130.5	101.1	62.2	27.0	5.4	1.4	0.0
Hispanic										
2012	0.9	31.4	93.9	129.8	112.6	77.9	36.8	9.6	0.2	0.0
2011	0.9	34.0	99.0	129.1	115.9	78.4	37.5	9.8	0.3	0.0
2010	0.8	40.6	110.1	134.8	114.2	75.0	37.6	9.7	0.3	0.0
2009	1.1	46.9	124.0	143.8	123.3	75.3	37.0	7.5	0.4	0.0
2008	1.3	51.2	127.4	158.5	127.2	81.9	36.1	8.7	0.4	0.0
White										
2012	0.1	9.5	44.7	80.1	98.0	86.6	34.4	6.9	0.8	0.0
2011	0.2	11.4	47.7	79.1	104.2	81.3	32.8	7.6	0.4	0.0
2010	0.2	12.0	50.4	83.4	100.2	82.5	35.0	7.6	1.1	0.0
2009	0.0	13.2	58.8	89.2	106.6	75.6	32.5	7.8	1.0	0.0
2008	0.2	14.6	55.3	98.2	110.3	78.5	35.2	8.2	0.7	0.0

See Technical Appendix for information on rates and race/ethnicity. * indicates U.S. data not available.

Rates based on few er than 20 events may be statistically unreliable and should be interpreted w ith caution.

Natality Section

**Table N-4(a) Number of Births by Mother's Race/Ethnicity, County, and Health Region
New Mexico, 2012, and United States, 2011**

	All Races	American Indian or Alaska Native	Asian or Pacific Islander	Black or African American	Hispanic	White	Unknown or Not Stated
United States	3,953,590	46,419	253,915	632,901	873,789	2,146,566	0
New Mexico	26,992	3,277	556	459	14,580	7,901	219
County							
Bernalillo	8,379	488	292	187	4,772	2,520	120
Catron	22	1	0	1	4	16	0
Chaves	948	6	10	7	632	287	6
Cibola	429	210	6	3	151	58	1
Colfax	115	1	1	0	79	33	1
Curry	924	9	23	64	387	440	1
De Baca	12	0	0	0	7	5	0
Dona Ana	3,071	23	45	45	2,158	788	12
Eddy	796	11	13	9	450	311	2
Grant	322	3	0	4	206	106	3
Guadalupe	47	1	0	0	43	3	0
Harding	5	0	0	0	3	2	0
Hidalgo	59	2	3	0	38	16	0
Lea	1,161	7	6	17	781	344	6
Lincoln	196	13	2	1	93	87	0
Los Alamos	178	3	27	2	36	110	0
Luna	419	4	2	0	362	51	0
McKinley	1,239	1,012	6	6	142	73	0
Mora	41	0	0	0	33	8	0
Otero	888	86	19	47	308	418	10
Quay	120	2	1	2	66	49	0
Rio Arriba	610	104	1	2	465	34	4
Roosevelt	271	2	2	3	110	152	2
Sandoval	1,459	300	36	26	596	487	14
San Juan	1,862	834	19	7	377	614	11
San Miguel	352	7	4	4	300	37	0
Santa Fe	1,358	38	24	7	934	349	6
Sierra	91	0	0	1	39	51	0
Socorro	211	53	2	2	111	39	4
Taos	319	18	5	0	219	75	2
Torrance	160	4	2	3	80	70	1
Union	53	0	0	0	16	37	0
Valencia	873	34	5	9	582	231	12
Unknown County	2	1	0	0	0	0	1
Health Region							
Northwest	3,530	2,056	31	16	670	745	12
Northeast	3,078	172	62	15	2,128	688	13
Metro	10,871	826	335	225	6,030	3,308	147
Southeast	4,428	50	57	103	2,526	1,675	17
Southwest	5,083	172	71	100	3,226	1,485	29
Unknown Region	2	1	0	0	0	0	1

See Technical Appendix for information on race/ethnicity and health regions.

Natality Section

**Table N-4(b) Percent of Births by Mother's Race/Ethnicity, County, and Health Region
New Mexico, 2012, and United States, 2011**

	American Indian or Alaska Native	Asian or Pacific Islander	Black or African American	Hispanic	White	Unknown or Not Stated
United States	1.2	6.4	16.0	22.1	54.3	0.0
New Mexico	12.1	2.1	1.7	54.0	29.3	0.8
County						
Bernalillo	5.8	3.5	2.2	57.0	30.1	1.4
Catron	4.5	0.0	4.5	18.2	72.7	0.0
Chaves	0.6	1.1	0.7	66.7	30.3	0.6
Cibola	49.0	1.4	0.7	35.2	13.5	0.2
Colfax	0.9	0.9	0.0	68.7	28.7	0.9
Curry	1.0	2.5	6.9	41.9	47.6	0.1
De Baca	0.0	0.0	0.0	58.3	41.7	0.0
Dona Ana	0.7	1.5	1.5	70.3	25.7	0.4
Eddy	1.4	1.6	1.1	56.5	39.1	0.3
Grant	0.9	0.0	1.2	64.0	32.9	0.9
Guadalupe	2.1	0.0	0.0	91.5	6.4	0.0
Harding	0.0	0.0	0.0	60.0	40.0	0.0
Hidalgo	3.4	5.1	0.0	64.4	27.1	0.0
Lea	0.6	0.5	1.5	67.3	29.6	0.5
Lincoln	6.6	1.0	0.5	47.4	44.4	0.0
Los Alamos	1.7	15.2	1.1	20.2	61.8	0.0
Luna	1.0	0.5	0.0	86.4	12.2	0.0
McKinley	81.7	0.5	0.5	11.5	5.9	0.0
Mora	0.0	0.0	0.0	80.5	19.5	0.0
Otero	9.7	2.1	5.3	34.7	47.1	1.1
Quay	1.7	0.8	1.7	55.0	40.8	0.0
Rio Arriba	17.0	0.2	0.3	76.2	5.6	0.7
Roosevelt	0.7	0.7	1.1	40.6	56.1	0.7
Sandoval	20.6	2.5	1.8	40.8	33.4	1.0
San Juan	44.8	1.0	0.4	20.2	33.0	0.6
San Miguel	2.0	1.1	1.1	85.2	10.5	0.0
Santa Fe	2.8	1.8	0.5	68.8	25.7	0.4
Sierra	0.0	0.0	1.1	42.9	56.0	0.0
Socorro	25.1	0.9	0.9	52.6	18.5	1.9
Taos	5.6	1.6	0.0	68.7	23.5	0.6
Torrance	2.5	1.3	1.9	50.0	43.8	0.6
Union	0.0	0.0	0.0	30.2	69.8	0.0
Valencia	3.9	0.6	1.0	66.7	26.5	1.4
Health Region						
Northwest	58.2	0.9	0.5	19.0	21.1	0.3
Northeast	5.6	2.0	0.5	69.1	22.4	0.4
Metro	7.6	3.1	2.1	55.5	30.4	1.4
Southeast	1.1	1.3	2.3	57.0	37.8	0.4
Southwest	3.4	1.4	2.0	63.5	29.2	0.6

See Technical Appendix for information on race/ethnicity and health regions.

The symbol -- indicates the percent cannot be calculated because no events occurred in the denominator value of this category.

Nativity Section

**Table N-5(a) Number of Births by Year, County, and Health Region
New Mexico, 2005-2012, and United States, 2005-2011***

	2005	2006	2007	2008	2009	2010	2011	2012
United States	4,138,349	4,265,555	4,316,233	4,247,694	4,130,665	3,999,386	3,953,590	*
New Mexico	28,822	29,918	30,605	30,156	28,873	27,795	27,251	26,992
County								
Bernalillo	9,297	9,633	9,809	9,416	9,122	8,589	8,385	8,379
Catron	21	27	19	15	24	19	22	22
Chaves	989	1,015	1,094	1,074	1,042	986	929	948
Cibola	433	439	458	428	420	418	395	429
Colfax	142	151	166	154	146	138	143	115
Curry	927	897	846	852	845	935	940	924
De Baca	17	18	18	29	22	25	24	12
Dona Ana	3,331	3,385	3,460	3,310	3,323	3,323	3,240	3,071
Eddy	659	763	789	822	766	729	768	796
Grant	358	405	363	406	367	355	292	322
Guadalupe	45	48	40	52	36	46	38	47
Harding	3	6	6	8	4	3	7	5
Hidalgo	63	57	66	83	60	39	76	59
Lea	1,046	1,126	1,226	1,230	1,202	1,044	1,063	1,161
Lincoln	214	213	235	234	233	190	198	196
Los Alamos	195	178	188	178	168	181	153	178
Luna	412	404	454	429	412	373	357	419
McKinley	1,332	1,415	1,404	1,362	1,343	1,248	1,305	1,239
Mora	45	49	42	52	36	44	44	41
Otero	846	857	866	889	869	874	934	888
Quay	116	121	98	115	100	109	111	120
Rio Arriba	708	744	549	705	630	622	576	610
Roosevelt	320	333	328	302	294	303	282	271
Sandoval	1,340	1,446	1,609	1,628	1,585	1,579	1,554	1,459
San Juan	2,069	2,188	2,242	2,320	2,163	1,928	1,910	1,862
San Miguel	365	382	358	389	314	342	317	352
Santa Fe	1,662	1,688	1,852	1,686	1,452	1,511	1,394	1,358
Sierra	117	97	110	115	107	97	87	91
Socorro	266	275	242	259	246	256	238	211
Taos	363	370	375	389	329	327	328	319
Torrance	167	178	200	186	165	163	173	160
Union	49	39	41	47	48	50	40	53
Valencia	905	971	1,047	990	999	947	928	873
Unknown, Not Reported	0	0	5	2	1	2	0	2
Health Regions								
Northwest	3,834	4,042	4,104	4,110	3,926	3,594	3,610	3,530
Northeast	3,577	3,655	3,617	3,660	3,163	3,264	3,040	3,078
Metro	11,709	12,228	12,665	12,220	11,871	11,278	11,040	10,871
Southeast	4,288	4,486	4,634	4,658	4,504	4,321	4,315	4,428
Southwest	5,414	5,507	5,580	5,506	5,408	5,336	5,246	5,083
Unknown, Not Reported	0	0	5	2	1	2	0	2

Unknown county and unknown region are included in New Mexico total.

See Technical Appendix for information on health regions. * indicates 2012 U.S. data not available at publication time.

Natality Section

**Table N-5(b) Birth Rates by Year, County, and Health Region
New Mexico, 2005-2012, and United States, 2005-2011***

	2005	2006	2007	2008	2009	2010	2011	2012
United States	14.0	14.3	14.3	14.0	13.5	13.0	12.7	*
New Mexico	14.8	15.2	15.4	15.0	14.2	13.5	13.1	12.9
County								
Bernalillo	15.2	15.5	15.5	14.7	14.0	12.9	12.5	12.4
Catron	5.7	7.3	5.1	4.0	6.5	5.1	5.9	5.9
Chaves	15.5	15.8	16.9	16.5	16.0	15.0	14.1	14.3
Cibola	16.3	16.5	17.1	15.9	15.5	15.3	14.4	15.7
Colfax	10.1	10.8	11.9	11.1	10.6	10.0	10.4	8.5
Curry	19.8	19.0	17.8	17.8	17.6	19.3	19.1	18.6
De Baca	8.0	8.5	8.6	14.0	10.8	12.4	12.1	6.0
Dona Ana	17.3	17.3	17.4	16.3	16.1	15.8	15.2	14.3
Eddy	12.4	14.4	14.8	15.4	14.3	13.5	14.1	14.5
Grant	11.8	13.4	12.1	13.6	12.4	12.0	9.9	11.0
Guadalupe	9.6	10.2	8.5	11.1	7.7	9.8	8.1	10.1
Harding	4.0	8.1	8.2	11.2	5.7	4.3	10.1	7.2
Hidalgo	11.6	10.7	12.7	16.3	12.0	8.0	15.7	12.3
Lea	17.3	18.4	19.7	19.5	18.8	16.1	16.2	17.5
Lincoln	10.7	10.6	11.6	11.5	11.4	9.3	9.6	9.5
Los Alamos	10.7	9.8	10.4	9.9	9.3	10.1	8.4	9.8
Luna	16.4	16.1	18.1	17.1	16.4	14.9	14.2	16.7
McKinley	18.2	19.4	19.4	18.9	18.7	17.5	17.7	16.9
Mora	8.9	9.8	8.4	10.5	7.3	9.0	9.1	8.5
Otero	13.4	13.5	13.7	14.0	13.6	13.7	14.5	13.7
Quay	12.1	12.7	10.4	12.4	10.9	12.1	12.3	13.6
Rio Arriba	17.3	18.3	13.5	17.4	15.6	15.5	14.3	15.2
Roosevelt	16.8	17.4	17.0	15.5	14.9	15.3	13.7	13.2
Sandoval	12.0	12.5	13.5	13.2	12.4	11.9	11.5	10.7
San Juan	17.0	17.7	17.9	18.3	16.8	14.8	14.9	14.5
San Miguel	12.2	12.8	12.1	13.2	10.6	11.6	10.8	12.1
Santa Fe	12.2	12.2	13.3	11.9	10.2	10.4	9.6	9.3
Sierra	9.2	7.8	8.9	9.4	8.8	8.1	7.3	7.6
Socorro	14.8	15.3	13.5	14.4	13.8	14.3	13.3	11.9
Taos	11.5	11.6	11.7	12.0	10.1	9.9	9.9	9.6
Torrance	10.0	10.7	12.1	11.3	10.0	10.0	10.6	9.9
Union	11.2	8.8	9.2	10.5	10.6	11.0	8.8	11.7
Valencia	12.6	13.4	14.2	13.3	13.2	12.3	12.0	11.3
Health Region								
Northwest	17.3	18.1	18.3	18.2	17.3	15.7	15.7	15.4
Northeast	12.5	12.7	12.5	12.6	10.8	11.1	10.3	10.4
Metro	14.4	14.8	15.1	14.3	13.6	12.7	12.2	12.0
Southeast	15.6	16.2	16.6	16.6	16.0	15.2	15.0	15.3
Southwest	15.4	15.6	15.6	15.3	14.9	14.6	14.1	13.6

See Technical Appendix for information on health regions. * indicates 2012 U.S. data not available at publication time.

Rates per 1,000 population for specified county or region.

Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Natality Section

**Table N-6(a) Number of Births by Mother's Age, County, and Health Region
New Mexico, 2012**

	All Ages	10-14	15 - 17	18 - 19	20 - 24	25 - 29	30 - 34	35 - 39	40 - 44	45 - 49	Unknown or 50+ Not Reported	
New Mexico	26,992	50	998	2,267	7,867	7,503	5,346	2,367	547	29	5	13
County												
Bernalillo	8,379	10	243	536	2,053	2,404	1,993	926	197	12	1	4
Catron	22	0	0	2	5	4	8	1	1	0	1	0
Chaves	948	1	39	109	342	240	139	68	10	0	0	0
Cibola	429	0	13	41	151	117	68	33	6	0	0	0
Colfax	115	0	4	12	38	34	24	3	0	0	0	0
Curry	924	0	45	83	303	269	160	56	7	0	0	1
De Baca	12	0	0	1	5	4	0	1	1	0	0	0
Dona Ana	3,071	3	129	312	992	802	567	204	58	3	1	0
Eddy	796	2	40	96	280	207	121	39	9	2	0	0
Grant	322	1	16	29	106	91	55	20	4	0	0	0
Guadalupe	47	0	5	4	19	8	8	3	0	0	0	0
Harding	5	0	0	0	1	1	2	0	0	0	0	1
Hidalgo	59	0	4	3	21	16	7	7	1	0	0	0
Lea	1,161	4	61	137	370	346	161	65	14	0	0	3
Lincoln	196	1	8	23	58	52	26	20	7	1	0	0
Los Alamos	178	0	2	2	26	44	70	25	8	0	0	1
Luna	419	2	41	51	145	116	35	21	7	0	0	1
McKinley	1,239	4	45	155	372	300	221	114	27	1	0	0
Mora	41	0	1	2	14	15	6	2	1	0	0	0
Otero	888	4	22	72	269	281	169	56	14	0	1	0
Quay	120	0	4	11	50	32	14	7	2	0	0	0
Rio Arriba	610	0	26	51	214	163	95	51	10	0	0	0
Roosevelt	271	1	11	21	92	92	36	14	4	0	0	0
Sandoval	1,459	0	39	105	375	439	319	141	37	3	0	1
San Juan	1,862	9	57	158	617	506	332	153	28	1	1	0
San Miguel	352	4	16	33	117	104	56	19	3	0	0	0
Santa Fe	1,358	1	54	85	348	336	296	176	57	5	0	0
Sierra	91	0	2	11	25	26	17	9	1	0	0	0
Socorro	211	2	10	25	66	54	34	16	4	0	0	0
Taos	319	0	9	16	85	78	82	39	10	0	0	0
Torrance	160	0	10	16	54	43	22	13	1	1	0	0
Union	53	0	0	2	17	20	9	4	1	0	0	0
Valencia	873	1	42	63	236	259	194	61	17	0	0	0
Unknown County	2	0	0	0	1	0	0	0	0	0	0	1
Health Region												
Northwest	3,530	13	115	354	1,140	923	621	300	61	2	1	0
Northeast	3,078	5	117	207	879	803	648	322	90	5	0	2
Metro	10,871	11	334	720	2,718	3,145	2,528	1,141	252	16	1	5
Southeast	4,428	9	208	481	1,500	1,242	657	270	54	3	0	4
Southwest	5,083	12	224	505	1,629	1,390	892	334	90	3	3	1
Unknown County	2	0	0	0	1	0	0	0	0	0	0	1

See Technical Appendix for information on health regions.

Natality Section

**Table N-6(b) Birth Rates by Mother's Age, County, and Health Region
New Mexico, 2012 (REVISED January 26, 2016)**

	All Ages	10-14	15 - 17	18 - 19	20 - 24	25 - 29	30 - 34	35 - 39	40 - 44	45 - 49	50+
New Mexico	12.9	0.7	22.7	77.3	113.2	109.1	83.9	37.8	8.7	0.4	0.0
County											
Bernalillo	12.4	0.5	17.7	58.6	81.6	93.5	87.9	43.4	9.3	0.5	0.0
Catron	5.9	0.0	0.0	64.2	106.7	77.8	110.6	19.0	18.4	0.0	0.9
Chaves	14.3	0.4	25.2	105.8	162.7	114.1	73.2	34.9	5.3	0.0	0.0
Cibola	15.7	0.0	22.6	106.7	163.6	138.9	80.5	42.1	7.7	0.0	0.0
Colfax	8.5	0.0	18.0	81.1	131.9	100.0	77.4	9.5	0.0	0.0	0.0
Curry	18.6	0.0	44.1	122.0	164.8	141.7	97.8	35.7	5.0	0.0	0.0
De Baca	6.0	0.0	0.0	47.6	138.9	72.4	0.0	29.1	18.0	0.0	0.0
Dona Ana	14.3	0.4	24.0	87.1	109.2	111.2	91.2	32.8	9.4	0.4	0.0
Eddy	14.5	1.0	35.2	126.6	180.0	124.0	72.9	24.5	5.7	1.1	0.0
Grant	11.0	1.2	31.0	84.2	141.4	126.4	75.5	27.6	5.5	0.0	0.0
Guadalupe	10.1	0.0	68.2	82.7	156.7	69.8	73.3	31.6	0.0	0.0	0.0
Harding	7.2	0.0	0.0	0.0	127.1	93.6	242.0	0.0	0.0	0.0	0.0
Hidalgo	12.3	0.0	36.8	41.4	147.0	148.9	60.9	55.7	6.6	0.0	0.0
Lea	17.5	1.6	41.1	138.5	164.9	152.9	74.4	33.1	7.2	0.0	0.0
Lincoln	9.5	1.9	25.7	110.8	140.9	112.4	64.6	38.4	13.4	1.2	0.0
Los Alamos	9.8	0.0	6.2	9.3	106.3	130.3	141.8	43.4	13.2	0.0	0.0
Luna	16.7	2.2	70.8	132.2	207.4	178.3	56.4	32.6	10.1	0.0	0.0
McKinley	16.9	1.3	21.8	112.4	137.3	127.5	97.1	51.6	12.2	0.4	0.0
Mora	8.5	0.0	12.0	36.4	148.3	118.1	62.6	15.7	6.6	0.0	0.0
Otero	13.7	1.9	18.2	89.1	124.9	131.5	89.6	32.3	7.9	0.0	0.1
Quay	13.6	0.0	25.2	103.8	292.7	143.6	62.2	30.3	8.0	0.0	0.0
Rio Arriba	15.2	0.0	32.6	96.0	178.1	145.7	82.6	41.4	8.4	0.0	0.0
Roosevelt	13.2	1.4	18.8	53.9	83.1	129.7	58.0	25.3	8.0	0.0	0.0
Sandoval	10.7	0.0	14.0	56.6	106.6	109.2	75.6	31.3	8.4	0.6	0.0
San Juan	14.5	1.8	19.3	80.2	144.7	110.5	81.4	39.4	7.6	0.2	0.0
San Miguel	12.1	4.6	22.9	70.8	125.0	129.3	78.2	22.9	3.8	0.0	0.0
Santa Fe	9.3	0.2	21.8	51.6	98.7	86.6	69.8	40.7	12.1	0.9	0.0
Sierra	7.6	0.0	13.0	106.8	108.9	121.8	94.2	40.9	3.5	0.0	0.0
Socorro	11.9	3.6	25.3	94.9	108.1	109.8	70.0	33.8	8.2	0.0	0.0
Taos	9.6	0.0	17.1	45.5	116.8	94.8	94.5	41.8	10.1	0.0	0.0
Torrance	9.9	0.0	31.8	76.4	154.0	111.4	57.4	30.4	2.3	1.6	0.0
Union	11.7	0.0	0.0	41.8	207.6	197.3	103.4	43.1	9.6	0.0	0.0
Valencia	11.3	0.3	25.1	56.5	109.3	116.3	88.5	26.5	6.9	0.0	0.0
Health Region											
Northwest	15.4	1.4	20.5	94.8	144.4	118.7	86.2	43.7	9.1	0.3	0.0
Northeast	10.4	0.6	22.2	58.8	121.6	104.9	80.2	37.7	10.0	0.5	0.0
Metro	12.0	0.4	18.1	58.4	87.2	97.2	85.8	40.0	8.8	0.5	0.0
Southeast	15.3	0.9	33.2	115.0	158.5	132.4	75.9	32.1	6.6	0.3	0.0
Southwest	13.6	1.0	26.7	90.3	118.8	120.0	86.5	32.8	8.7	0.2	0.0

The birth rates for "All Ages" are births per 1,000 population and are taken from Table N-5(b).

Birth Rates by mother's age groups are per 1,000 women in the specified age category.

Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Please be aware that this table has been updated and the corrected page inserted on January 26, 2016.

Natality Section

**Table N-7 Number and Percent of Births to Single Mothers by Mother's Age and Race/Ethnicity
New Mexico, 2012**

Number	All Ages	Percent										Unknown or Not Reported
		10-14	15 - 17	18 - 19	20 - 24	25 - 29	30-34	35-39	40-44	45 - 49	50+ Reported	
All Races	14,016	48	934	1,919	5,181	3,251	1,735	755	180	12	0	1
American Indian or Alaska Native	2,583	12	125	333	923	640	352	158	37	3	0	0
Asian or Pacific Islander	94	0	1	7	22	29	20	12	2	1	0	0
Black or African American	245	1	9	33	99	57	35	8	2	1	0	0
Hispanic	8,479	33	664	1,207	3,160	1,941	992	388	92	2	0	0
White	2,511	2	126	329	941	561	321	181	44	5	0	1
Unknown or Not Reported	104	0	9	10	36	23	15	8	3	0	0	0
Percent												
All Races	100.0	0.3	6.7	13.7	37.0	23.2	12.4	5.4	1.3	0.1	0.0	0.0
American Indian or Alaska Native	100.0	0.5	4.8	12.9	35.7	24.8	13.6	6.1	1.4	0.1	0.0	0.0
Asian or Pacific Islander	100.0	0.0	1.1	7.4	23.4	30.9	21.3	12.8	2.1	1.1	0.0	0.0
Black or African American	100.0	0.4	3.7	13.5	40.4	23.3	14.3	3.3	0.8	0.4	0.0	0.0
Hispanic	100.0	0.4	7.8	14.2	37.3	22.9	11.7	4.6	1.1	0.0	0.0	0.0
White	100.0	0.1	5.0	13.1	37.5	22.3	12.8	7.2	1.8	0.2	0.0	0.0
Unknown or Not Reported	100.0	0.0	8.7	9.6	34.6	22.1	14.4	7.7	2.9	0.0	0.0	0.0

See Technical Appendix for information on race/ethnicity.

Natality Section

**Table N-8 Number and Percent of Births to Single Mothers by County and Health Region
New Mexico, 1990, 2000, and 2012**

	1990		2000		2012	
	Number	Percent	Number	Percent	Number	Percent
New Mexico	9,683	35.4	12,400	45.6	14,016	51.9
County						
Bernalillo	2,601	32.1	3,484	42.0	3,992	47.6
Catron	6	24.0	5	29.4	11	50.0
Chaves	355	37.8	439	45.8	480	50.6
Cibola	265	54.6	277	62.8	302	70.4
Colfax	45	27.4	74	46.0	75	65.2
Curry	237	24.9	283	36.2	382	41.3
De Baca	6	42.9	6	27.3	5	41.7
Dona Ana	906	31.8	1,435	47.4	1,643	53.5
Eddy	279	35.2	306	41.7	423	53.1
Grant	146	36.7	194	46.2	186	57.8
Guadalupe	26	35.1	42	62.7	36	76.6
Harding	1	14.3	1	16.7	2	40.0
Hidalgo	27	31.4	35	45.5	30	50.8
Lea	321	32.8	409	46.9	597	51.4
Lincoln	49	26.5	98	47.1	114	58.2
Los Alamos	15	8.3	25	12.9	29	16.3
Luna	133	42.9	202	53.0	245	58.5
McKinley	1,112	60.2	891	65.7	930	75.1
Mora	22	44.0	30	56.6	21	51.2
Otero	241	24.1	268	31.0	351	39.5
Quay	54	38.0	59	53.6	63	52.5
Rio Arriba	322	44.5	402	57.8	441	72.3
Roosevelt	74	29.4	113	37.3	109	40.2
Sandoval	411	32.6	543	43.5	715	49.0
San Juan	817	41.8	991	50.7	1,032	55.4
San Miguel	182	45.0	221	57.4	231	65.6
Santa Fe	470	32.6	665	40.6	663	48.8
Sierra	27	27.3	45	45.0	46	50.5
Socorro	100	37.7	159	59.1	134	63.5
Taos	139	37.0	191	54.6	201	63.0
Torrance	44	31.4	85	45.2	80	50.0
Union	14	21.2	14	36.8	18	34.0
Valencia	236	31.8	408	41.1	428	49.0
Unknown, Not Reported	0	0.0	0	0.0	1	50.0
Health Region						
Northwest	2,194	51.2	2,159	57.6	2,264	64.1
Northeast	1,236	35.5	1,665	46.4	1,717	55.8
Metro	3,292	32.1	4,520	42.1	5,215	48.0
Southeast	1,375	32.3	1,713	43.0	2,173	49.1
Southwest	1,586	31.5	2,343	45.5	2,646	52.1
Unknown, Not Reported	0	0.0	0	0.0	1	50.0

See Technical Appendix for information on health regions. Unknown county and region are included in New Mexico total.

Nativity Section

**Table N-9 Number and Percent of Births to Single Mothers by Mother's Age
New Mexico, 1990, 2000, and 2012**

Number	1990	2000	2012
All Ages	9,683	12,400	14,016
10 to 14	83	76	48
15 to 17	1,266	1,535	934
18 to 19	1,772	2,263	1,919
20 to 24	3,353	4,511	5,181
25 to 29	1,861	2,217	3,251
30 to 34	877	1,063	1,735
35 to 39	384	555	755
40 to 44	73	155	180
45 to 49	6	8	12
50+	0	0	0
Unknown or Not Reported Age	8	17	1

Percent	1990	2000	2012
All Ages	35.4	45.6	51.9
10 to 14	94.3	97.4	96.0
15 to 17	81.3	90.5	93.6
18 to 19	63.4	76.6	84.6
20 to 24	42.4	55.3	65.9
25 to 29	23.9	32.8	43.3
30 to 34	17.9	23.1	32.5
35 to 39	19.9	23.5	31.9
40 to 44	22.1	28.3	32.9
45 to 49	31.6	34.8	41.4
50+	--	0.0	0.0
Unknown or Not Reported Age	50.0	58.6	7.7

The symbol -- indicates the percent cannot be calculated because no events occurred in the denominator value of this category.

Natality Section

**Table N-10 Percent Distribution of Birth Order by Mother's Race/Ethnicity
New Mexico, 1990, 2000, and 2012**

Birth Order and Race/Ethnicity	1990	2000	2012
All Races			
Total	27,317	27,206	26,992
1	38.0	39.4	37.1
2	31.5	31.7	27.3
3	17.3	17.1	17.5
4	7.8	7.2	8.3
5+	5.4	4.5	6.1
American Indian or Alaska Native			
Total	3,871	3,357	3,277
1	27.3	32.5	35.1
2	24.5	25.9	24.7
3	20.0	17.4	15.9
4	14.3	11.7	10.2
5+	13.9	12.5	10.9
Asian or Pacific Islander			
Total	284	394	556
1	43.0	44.2	43.2
2	36.3	37.8	32.7
3	13.0	12.2	13.3
4	6.7	3.8	4.5
5+	1.1	2.0	2.5
Black or African American			
Total	510	471	459
1	40.4	45.4	38.6
2	28.4	28.2	26.6
3	18.6	16.3	16.3
4	9.2	7.0	7.2
5+	3.3	3.0	7.4
Hispanic			
Total	12,145	13,940	14,580
1	38.8	38.8	35.7
2	31.2	32.2	26.9
3	17.5	18.2	18.8
4	7.5	7.1	9.0
5+	5.0	3.7	5.7
White			
Total	10,501	9,044	7,901
1	40.9	42.4	40.5
2	34.3	33.1	29.2
3	16.1	15.6	16.3
4	5.6	5.9	6.7
5+	3.1	3.0	5.1

See Technical Appendix for information on race/ethnicity and birth order.

Natality Section

**Table N-11(a) Number of Births by Mother's Age, Race/Ethnicity, and Infant's Birthweight
New Mexico, 2012**

Birthweight (grams)	All Ages	10 to 14	15 to 17	18 to 19	20 to 24	25 to 29	30 to 34	35 to 39	40 to 44	45 to 49	50+	Unknown or Not Reported
												Age
All Races												
Less than 1500	344	1	10	33	84	102	70	33	11	0	0	0
1500 to 2499	2,044	6	78	172	585	535	390	205	60	5	2	6
2500 to 3999	23,129	42	877	1,989	6,872	6,425	4,493	1,966	434	22	3	6
4000+	1,363	0	30	66	304	413	365	144	39	2	0	0
Unknown	112	1	3	7	22	28	28	19	3	0	0	1
Total	26,992	50	998	2,267	7,867	7,503	5,346	2,367	547	29	5	13
American Indian or Alaska Native												
Less than 1500	40	1	1	4	7	10	11	4	2	0	0	0
1500 to 2499	227	1	8	20	71	45	47	27	8	0	0	0
2500 to 3999	2,803	10	114	312	936	756	428	196	47	4	0	0
4000+	196	0	4	10	47	61	47	22	5	0	0	0
Unknown	11	0	0	1	2	3	3	2	0	0	0	0
Total	3,277	12	127	347	1,063	875	536	251	62	4	0	0
Asian or Pacific Islander												
Less than 1500	9	0	0	0	0	2	3	4	0	0	0	0
1500 to 2499	65	0	1	1	7	15	31	9	0	1	0	0
2500 to 3999	461	0	0	11	54	120	151	109	15	0	0	1
4000+	17	0	0	0	2	8	6	1	0	0	0	0
Unknown	4	0	0	0	0	1	1	1	1	0	0	0
Total	556	0	1	12	63	146	192	124	16	1	0	1
Black or African American												
Less than 1500	7	0	0	0	2	2	2	0	1	0	0	0
1500 to 2499	57	0	1	4	17	19	13	2	1	0	0	0
2500 to 3999	382	1	8	35	126	95	77	32	7	1	0	0
4000+	13	0	0	1	3	3	6	0	0	0	0	0
Unknown	0	0	0	0	0	0	0	0	0	0	0	0
Total	459	1	9	40	148	119	98	34	9	1	0	0
Hispanic												
Less than 1500	184	0	6	26	51	52	31	14	4	0	0	0
1500 to 2499	1,103	5	56	110	339	292	171	99	26	0	1	4
2500 to 3999	12,582	29	630	1,259	4,081	3,371	2,102	894	208	6	1	1
4000+	642	0	22	32	144	206	151	64	22	1	0	0
Unknown	69	1	3	3	14	16	18	13	1	0	0	0
Total	14,580	35	717	1,430	4,629	3,937	2,473	1,084	261	7	2	5
White												
Less than 1500	94	0	3	2	22	32	23	8	4	0	0	0
1500 to 2499	575	0	12	37	146	163	123	64	24	4	1	1
2500 to 3999	6,723	2	116	358	1,629	2,029	1,700	721	153	11	1	3
4000+	484	0	4	22	104	133	153	57	10	1	0	0
Unknown	25	0	0	3	5	8	6	2	1	0	0	0
Total	7,901	2	135	422	1,906	2,365	2,005	852	192	16	2	4

See Technical Appendix for information on birthweight and race/ethnicity.

Natality Section

**Table N-11(b) Percent of Births by Mother's Age, Race/Ethnicity, and Infant's Birthweight
New Mexico, 2012**

Birthweight (grams)	10 to 14	15 to 17	18 to 19	20 to 24	25 to 29	30 to 34	35 to 39	40 to 44	45 to 49	50+	Unknown or Not Reported Age
All Races											
Less than 1500	0.3	2.9	9.6	24.4	29.7	20.3	9.6	3.2	0.0	0.0	0.0
1500 to 2499	0.3	3.8	8.4	28.6	26.2	19.1	10.0	2.9	0.2	0.1	0.3
2500 to 3999	0.2	3.8	8.6	29.7	27.8	19.4	8.5	1.9	0.1	0.0	0.0
4000+	0.0	2.2	4.8	22.3	30.3	26.8	10.6	2.9	0.1	0.0	0.0
Unknown	0.9	2.7	6.3	19.6	25.0	25.0	17.0	2.7	0.0	0.0	0.9
Total	0.2	3.7	8.4	29.1	27.8	19.8	8.8	2.0	0.1	0.0	0.0
American Indian or Alaska Native											
Less than 1500	2.5	2.5	10.0	17.5	25.0	27.5	10.0	5.0	0.0	0.0	0.0
1500 to 2499	0.4	3.5	8.8	31.3	19.8	20.7	11.9	3.5	0.0	0.0	0.0
2500 to 3999	0.4	4.1	11.1	33.4	27.0	15.3	7.0	1.7	0.1	0.0	0.0
4000+	0.0	2.0	5.1	24.0	31.1	24.0	11.2	2.6	0.0	0.0	0.0
Unknown	0.0	0.0	9.1	18.2	27.3	27.3	18.2	0.0	0.0	0.0	0.0
Total	0.4	3.9	10.6	32.4	26.7	16.4	7.7	1.9	0.1	0.0	0.0
Asian or Pacific Islander											
Less than 1500	0.0	0.0	0.0	0.0	22.2	33.3	44.4	0.0	0.0	0.0	0.0
1500 to 2499	0.0	1.5	1.5	10.8	23.1	47.7	13.8	0.0	1.5	0.0	0.0
2500 to 3999	0.0	0.0	2.4	11.7	26.0	32.8	23.6	3.3	0.0	0.0	0.2
4000+	0.0	0.0	0.0	11.8	47.1	35.3	5.9	0.0	0.0	0.0	0.0
Unknown	0.0	0.0	0.0	0.0	25.0	25.0	25.0	25.0	0.0	0.0	0.0
Total	0.0	0.2	2.2	11.3	26.3	34.5	22.3	2.9	0.2	0.0	0.2
Black or African American											
Less than 1500	0.0	0.0	0.0	28.6	28.6	28.6	0.0	14.3	0.0	0.0	0.0
1500 to 2499	0.0	1.8	7.0	29.8	33.3	22.8	3.5	1.8	0.0	0.0	0.0
2500 to 3999	0.3	2.1	9.2	33.0	24.9	20.2	8.4	1.8	0.3	0.0	0.0
4000+	0.0	0.0	7.7	23.1	23.1	46.2	0.0	0.0	0.0	0.0	0.0
Unknown	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Total	0.2	2.0	8.7	32.2	25.9	21.4	7.4	2.0	0.2	0.0	0.0
Hispanic											
Less than 1500	0.0	3.3	14.1	27.7	28.3	16.8	7.6	2.2	0.0	0.0	0.0
1500 to 2499	0.5	5.1	10.0	30.7	26.5	15.5	9.0	2.4	0.0	0.1	0.4
2500 to 3999	0.2	5.0	10.0	32.4	26.8	16.7	7.1	1.7	0.0	0.0	0.0
4000+	0.0	3.4	5.0	22.4	32.1	23.5	10.0	3.4	0.2	0.0	0.0
Unknown	1.4	4.3	4.3	20.3	23.2	26.1	18.8	1.4	0.0	0.0	0.0
Total	0.2	4.9	9.8	31.7	27.0	17.0	7.4	1.8	0.0	0.0	0.0
White											
Less than 1500	0.0	3.2	2.1	23.4	34.0	24.5	8.5	4.3	0.0	0.0	0.0
1500 to 2499	0.0	2.1	6.4	25.4	28.3	21.4	11.1	4.2	0.7	0.2	0.2
2500 to 3999	0.0	1.7	5.3	24.2	30.2	25.3	10.7	2.3	0.2	0.0	0.0
4000+	0.0	0.8	4.5	21.5	27.5	31.6	11.8	2.1	0.2	0.0	0.0
Unknown	0.0	0.0	12.0	20.0	32.0	24.0	8.0	4.0	0.0	0.0	0.0
Total	0.0	1.7	5.3	24.1	29.9	25.4	10.8	2.4	0.2	0.0	0.1

See Technical Appendix for information on birthweight and race/ethnicity.

Natality Section

**Table N-12 Number of Single and Multiple Births by Mother's Age and Race/Ethnicity
New Mexico, 2010-2012 Total**

	All Ages	10 to 14	15 to 17	18 to 19	20 to 24	25 to 29	30 to 34	35 to 39	40 to 44	45 to 49	50+	Unknown or Not Reported Age
All Races												
Single	79,884	140	3,347	7,104	23,500	22,192	15,092	6,776	1,616	87	11	19
Twins	2,106	0	42	78	423	606	539	322	78	8	6	4
Triplets	48	0	3	3	6	6	12	12	6	0	0	0
Total	82,038	140	3,392	7,185	23,929	22,804	15,643	7,110	1,700	95	17	23
American Indian or Alaska Native												
Single	9,890	23	439	1,028	3,244	2,547	1,592	806	199	11	0	1
Twins	190	0	2	12	45	62	43	26	0	0	0	0
Total	10,080	23	441	1,040	3,289	2,609	1,635	832	199	11	0	1
Asian Pacific Islander												
Single	1,568	1	9	33	194	458	510	309	51	2	0	1
Twins	68	0	0	0	2	16	34	12	4	0	0	0
Total	1,636	1	9	33	196	474	544	321	55	2	0	1
Black or African American												
Single	1,342	3	43	118	463	333	254	103	21	3	1	0
Twins	70	0	0	3	24	16	18	7	0	2	0	0
Total	1,412	3	43	121	487	349	272	110	21	5	1	0
Hispanic												
Single	43,523	102	2,387	4,555	13,799	11,642	7,091	3,145	762	32	2	6
Twins	1,010	0	31	47	243	314	219	126	28	0	0	2
Triplets	18	0	0	3	3	3	3	3	3	0	0	0
Total	44,551	102	2,418	4,605	14,045	11,959	7,313	3,274	793	32	2	8
White												
Single	23,044	11	452	1,327	5,660	7,075	5,537	2,364	564	39	7	8
Twins	747	0	9	16	102	196	221	145	46	6	6	0
Triplets	30	0	3	0	3	3	9	9	3	0	0	0
Total	23,821	11	464	1,343	5,765	7,274	5,767	2,518	613	45	13	8

The number displayed is the total number of births in the specified multiple birth category (not the sets) for years 2010-2012. See Technical Appendix for information on race/ethnicity.

Nativity Section

**Table N-13(a) Number of Births by Birthweight (Grams), County, and Health Region
New Mexico, 2012, and United States, 2011**

	Total Births	Less than 1500	1500 to 2499	2500 to 3999	4000+	Unknown, Not Reported Weight
United States	3,953,590	56,754	262,957	3,322,458	306,872	4,549
New Mexico	26,992	344	2,044	23,129	1,363	112
County						
Bernalillo	8,379	108	715	7,056	431	69
Catron	22	1	2	18	1	0
Chaves	948	14	48	840	46	0
Cibola	429	5	36	378	10	0
Colfax	115	1	11	100	2	1
Curry	924	7	73	773	70	1
De Baca	12	0	1	11	0	0
Dona Ana	3,071	32	190	2,678	170	1
Eddy	796	10	43	685	57	1
Grant	322	9	21	284	7	1
Guadalupe	47	0	4	41	1	1
Harding	5	0	0	5	0	0
Hidalgo	59	2	5	50	1	1
Lea	1,161	11	73	1015	62	0
Lincoln	196	3	22	162	9	0
Los Alamos	178	3	21	141	13	0
Luna	419	5	20	377	16	1
McKinley	1,239	16	85	1,070	62	6
Mora	41	0	1	38	2	0
Otero	888	11	71	759	45	2
Quay	120	4	11	98	7	0
Rio Arriba	610	9	59	523	19	0
Roosevelt	271	4	15	240	12	0
Sandoval	1,459	19	106	1,256	72	6
San Juan	1,862	20	116	1,595	126	5
San Miguel	352	6	33	298	15	0
Santa Fe	1,358	21	117	1,179	35	6
Sierra	91	1	9	80	1	0
Socorro	211	5	23	177	6	0
Taos	319	5	30	274	10	0
Torrance	160	1	12	136	10	1
Union	53	1	3	46	3	0
Valencia	873	10	68	745	42	8
Unknown, Not Reported	2	0	0	1	0	1
Health Region						
Northwest	3,530	41	237	3,043	198	11
Northeast	3,078	46	279	2,645	100	8
Metro	10,871	138	901	9,193	555	84
Southeast	4,428	53	286	3,824	263	2
Southwest	5,083	66	341	4,423	247	6
Unknown, Not Reported	2	0	0	1	0	1

See Technical Appendix for information on infant birthweights.

Natality Section

**Table N-13(b) Percent of Births by Birthweight (Grams), County, and Health Region
New Mexico, 2012, and United States, 2011**

	Less than 1500	1500 to 2499	2500 to 3999	4000+	Unknown, Not Reported Weight
United States	1.4	6.7	84.0	7.8	0.1
New Mexico	1.3	7.6	85.7	5.0	0.4
County					
Bernalillo	1.3	8.5	84.2	5.1	0.8
Catron	4.5	9.1	81.8	4.5	0.0
Chaves	1.5	5.1	88.6	4.9	0.0
Cibola	1.2	8.4	88.1	2.3	0.0
Colfax	0.9	9.6	87.0	1.7	0.9
Curry	0.8	7.9	83.7	7.6	0.1
De Baca	0.0	8.3	91.7	0.0	0.0
Dona Ana	1.0	6.2	87.2	5.5	0.0
Eddy	1.3	5.4	86.1	7.2	0.1
Grant	2.8	6.5	88.2	2.2	0.3
Guadalupe	0.0	8.5	87.2	2.1	2.1
Harding	0.0	0.0	100.0	0.0	0.0
Hidalgo	3.4	8.5	84.7	1.7	1.7
Lea	0.9	6.3	87.4	5.3	0.0
Lincoln	1.5	11.2	82.7	4.6	0.0
Los Alamos	1.7	11.8	79.2	7.3	0.0
Luna	1.2	4.8	90.0	3.8	0.2
McKinley	1.3	6.9	86.4	5.0	0.5
Mora	0.0	2.4	92.7	4.9	0.0
Otero	1.2	8.0	85.5	5.1	0.2
Quay	3.3	9.2	81.7	5.8	0.0
Rio Arriba	1.5	9.7	85.7	3.1	0.0
Roosevelt	1.5	5.5	88.6	4.4	0.0
Sandoval	1.3	7.3	86.1	4.9	0.4
San Juan	1.1	6.2	85.7	6.8	0.3
San Miguel	1.7	9.4	84.7	4.3	0.0
Santa Fe	1.5	8.6	86.8	2.6	0.4
Sierra	1.1	9.9	87.9	1.1	0.0
Socorro	2.4	10.9	83.9	2.8	0.0
Taos	1.6	9.4	85.9	3.1	0.0
Torrance	0.6	7.5	85.0	6.3	0.6
Union	1.9	5.7	86.8	5.7	0.0
Valencia	1.1	7.8	85.3	4.8	0.9
Unknown, Not Reported	0.0	0.0	50.0	0.0	50.0
Health Region					
Northwest	1.2	6.7	86.2	5.6	0.3
Northeast	1.5	9.1	85.9	3.2	0.3
Metro	1.3	8.3	84.6	5.1	0.8
Southeast	1.2	6.5	86.4	5.9	0.0
Southwest	1.3	6.7	87.0	4.9	0.1
Unknown, Not Reported	0.0	0.0	50.0	0.0	50.0

See Technical Appendix for information on infant birthweights.

Natality Section

**Table N-14(a) Number of Births by Level of Prenatal Care, County, and Health Region
New Mexico, 2012**

	No Prenatal					
	Total Births	Care	Low	Moderate	High	Unknown
New Mexico	26,992	341	2,418	7,460	14,331	2,442
County						
Bernalillo	8,379	44	658	2,226	4,243	1,208
Catron	22	0	5	9	6	2
Chaves	948	14	82	300	535	17
Cibola	429	7	88	141	161	32
Colfax	115	4	7	37	64	3
Curry	924	7	91	314	463	49
De Baca	12	0	0	2	10	0
Dona Ana	3,071	107	354	934	1,642	34
Eddy	796	5	59	189	405	138
Grant	322	6	8	86	210	12
Guadalupe	47	0	7	14	22	4
Harding	5	0	0	0	5	0
Hidalgo	59	3	2	24	27	3
Lea	1,161	25	77	333	519	207
Lincoln	196	5	27	58	97	9
Los Alamos	178	1	5	30	135	7
Luna	419	13	38	112	233	23
McKinley	1,239	22	168	418	574	57
Mora	41	0	1	18	21	1
Otero	888	15	55	216	473	129
Quay	120	5	10	42	55	8
Rio Arriba	610	8	52	156	376	18
Roosevelt	271	6	23	87	130	25
Sandoval	1,459	5	100	320	904	130
San Juan	1,862	21	230	588	966	57
San Miguel	352	3	24	77	230	18
Santa Fe	1,358	1	55	231	1,019	52
Sierra	91	2	17	24	44	4
Socorro	211	2	31	86	70	22
Taos	319	2	41	88	180	8
Torrance	160	0	19	42	66	33
Union	53	0	5	16	32	0
Valencia	873	7	79	241	414	132
Unknown, Not Reported	2	1	0	1	0	0
Health Region						
Northwest	3,530	50	486	1,147	1,701	146
Northeast	3,078	19	197	667	2,084	111
Metro	10,871	56	856	2,829	5,627	1,503
Southeast	4,428	67	369	1,325	2,214	453
Southwest	5,083	148	510	1,491	2,705	229
Unknown, Not Reported	2	1	0	1	0	0

See Technical Appendix for information on the Modified Kessner Index.

Note: This table uses the Modified Kessner Index.

Nativity Section

**N-14(b) Percent of Births by Level of Prenatal Care, County, and Health Region
New Mexico, 2012**

	No Prenatal Care	Low	Moderate	High	Unknown
New Mexico	1.3	9.0	27.6	53.1	9.0
County					
Bernalillo	0.5	7.9	26.6	50.6	14.4
Catron	0.0	22.7	40.9	27.3	9.1
Chaves	1.5	8.6	31.6	56.4	1.8
Cibola	1.6	20.5	32.9	37.5	7.5
Colfax	3.5	6.1	32.2	55.7	2.6
Curry	0.8	9.8	34.0	50.1	5.3
De Baca	0.0	0.0	16.7	83.3	0.0
Dona Ana	3.5	11.5	30.4	53.5	1.1
Eddy	0.6	7.4	23.7	50.9	17.3
Grant	1.9	2.5	26.7	65.2	3.7
Guadalupe	0.0	14.9	29.8	46.8	8.5
Harding	0.0	0.0	0.0	100.0	0.0
Hidalgo	5.1	3.4	40.7	45.8	5.1
Lea	2.2	6.6	28.7	44.7	17.8
Lincoln	2.6	13.8	29.6	49.5	4.6
Los Alamos	0.6	2.8	16.9	75.8	3.9
Luna	3.1	9.1	26.7	55.6	5.5
McKinley	1.8	13.6	33.7	46.3	4.6
Mora	0.0	2.4	43.9	51.2	2.4
Otero	1.7	6.2	24.3	53.3	14.5
Quay	4.2	8.3	35.0	45.8	6.7
Rio Arriba	1.3	8.5	25.6	61.6	3.0
Roosevelt	2.2	8.5	32.1	48.0	9.2
Sandoval	0.3	6.9	21.9	62.0	8.9
San Juan	1.1	12.4	31.6	51.9	3.1
San Miguel	0.9	6.8	21.9	65.3	5.1
Santa Fe	0.1	4.1	17.0	75.0	3.8
Sierra	2.2	18.7	26.4	48.4	4.4
Socorro	0.9	14.7	40.8	33.2	10.4
Taos	0.6	12.9	27.6	56.4	2.5
Torrance	0.0	11.9	26.3	41.3	20.6
Union	0.0	9.4	30.2	60.4	0.0
Valencia	0.8	9.0	27.6	47.4	15.1
Unknown, Not Reported	50.0	0.0	50.0	0.0	0.0
Health Region					
Northwest	1.4	13.8	32.5	48.2	4.1
Northeast	0.6	6.4	21.7	67.7	3.6
Metro	0.5	7.9	26.0	51.8	13.8
Southeast	1.5	8.3	29.9	50.0	10.2
Southwest	2.9	10.0	29.3	53.2	4.5
Unknown, Not Reported	50.0	0.0	50.0	0.0	0.0

See Technical Appendix for information on the Modified Kessner Index.

Note: This table uses the Modified Kessner Index.

Nativity Section

Table N-15(a) Number of Births with Low or No Prenatal Care by Mother's Race/Ethnicity, County and Health Region, New Mexico, 2012

	Mother's Race/Ethnicity							
	All Races	American Indian or Alaska Native	Asian or Pacific Islander	Black or African American	Hispanic	White	Unknown or Not Reported Race	
New Mexico	2,759	553	40	54	1,506	579		27
County								
Bernalillo	702	59	23	22	417	167		14
Catron	5	0	0	0	0	5		0
Chaves	96	0	1	0	68	27		0
Cibola	95	59	0	0	27	9		0
Colfax	11	0	0	0	6	5		0
Curry	98	0	2	7	55	34		0
De Baca	0	0	0	0	0	0		0
Dona Ana	461	6	7	9	364	73		2
Eddy	64	0	0	3	44	15		2
Grant	14	0	0	1	10	3		0
Guadalupe	7	0	0	0	6	1		0
Harding	0	0	0	0	0	0		0
Hidalgo	5	1	0	0	2	2		0
Lea	102	0	1	1	72	28		0
Lincoln	32	4	0	0	12	16		0
Los Alamos	6	0	1	0	2	3		0
Luna	51	1	0	0	45	5		0
McKinley	190	178	0	0	8	4		0
Mora	1	0	0	0	1	0		0
Otero	70	16	1	0	31	22		0
Quay	15	0	0	0	6	9		0
Rio Arriba	60	11	0	0	47	1		1
Roosevelt	29	0	0	1	13	13		2
Sandoval	105	45	0	3	35	20		2
San Juan	251	134	2	2	59	53		1
San Miguel	27	0	0	2	23	2		0
Santa Fe	56	6	0	0	33	17		0
Sierra	19	0	0	0	8	11		0
Socorro	33	14	0	0	15	3		1
Taos	43	10	1	0	26	6		0
Torrance	19	1	0	0	8	10		0
Union	5	0	0	0	1	4		0
Valencia	86	8	1	3	62	11		1
Unknown, Not Reported	1	0	0	0	0	0		1
Health Region								
Northwest	536	371	2	2	94	66		1
Northeast	216	27	2	2	145	39		1
Metro	912	113	24	28	522	208		17
Southeast	436	4	4	12	270	142		4
Southwest	658	38	8	10	475	124		3
Unknown, Not Reported	1	0	0	0	0	0		1

See Technical Appendix for information on health regions and the Modified Kessner Index.

Note: This table uses the Modified Kessner Index.

Natality Section

Table N-15(b) Percent of Births with Low or No Prenatal Care by Mother's Race/Ethnicity, County and Health Region, New Mexico, 2012

	Mother's Race/Ethnicity							Unknown or Not Reported
	All Races	American Indian or Alaska Native	Asian or Pacific Islander	Black or African American	Hispanic	White		
New Mexico	10.2	16.9	7.2	11.8	10.3	7.3	12.3	
County								
Bernalillo	8.4	12.1	7.9	11.8	8.7	6.6	11.7	
Catron	22.7	0.0	0.0	0.0	0.0	31.3	0.0	
Chaves	10.1	0.0	10.0	0.0	10.8	9.4	0.0	
Cibola	22.1	28.1	0.0	0.0	17.9	15.5	0.0	
Colfax	9.6	0.0	0.0	0.0	7.6	15.2	0.0	
Curry	10.6	0.0	8.7	10.9	14.2	7.7	0.0	
De Baca	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Dona Ana	15.0	26.1	15.6	20.0	16.9	9.3	16.7	
Eddy	8.0	0.0	0.0	33.3	9.8	4.8	100.0	
Grant	4.3	0.0	0.0	25.0	4.9	2.8	0.0	
Guadalupe	14.9	0.0	0.0	0.0	14.0	33.3	0.0	
Harding	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Hidalgo	8.5	50.0	0.0	0.0	5.3	12.5	0.0	
Lea	8.8	0.0	16.7	5.9	9.2	8.1	0.0	
Lincoln	16.3	30.8	0.0	0.0	12.9	18.4	0.0	
Los Alamos	3.4	0.0	3.7	0.0	5.6	2.7	0.0	
Luna	12.2	25.0	0.0	0.0	12.4	9.8	0.0	
McKinley	15.3	17.6	0.0	0.0	5.6	5.5	0.0	
Mora	2.4	0.0	0.0	0.0	3.0	0.0	0.0	
Otero	7.9	18.6	5.3	0.0	10.1	5.3	0.0	
Quay	12.5	0.0	0.0	0.0	9.1	18.4	0.0	
Rio Arriba	9.8	10.6	0.0	0.0	10.1	2.9	25.0	
Roosevelt	10.7	0.0	0.0	33.3	11.8	8.6	100.0	
Sandoval	7.2	15.0	0.0	11.5	5.9	4.1	14.3	
San Juan	13.5	16.1	10.5	28.6	15.6	8.6	9.1	
San Miguel	7.7	0.0	0.0	50.0	7.7	5.4	0.0	
Santa Fe	4.1	15.8	0.0	0.0	3.5	4.9	0.0	
Sierra	20.9	0.0	0.0	0.0	20.5	21.6	0.0	
Socorro	15.6	26.4	0.0	0.0	13.5	7.7	25.0	
Taos	13.5	55.6	20.0	0.0	11.9	8.0	0.0	
Torrance	11.9	25.0	0.0	0.0	10.0	14.3	0.0	
Union	9.4	0.0	0.0	0.0	6.3	10.8	0.0	
Valencia	9.9	23.5	20.0	33.3	10.7	4.8	8.3	
Unknown, Not Reported	50.0	0.0	0.0	0.0	0.0	0.0	100.0	
Health Region								
Northwest	15.2	18.0	6.5	12.5	14.0	8.9	8.3	
Northeast	7.0	15.7	3.2	13.3	6.8	5.7	7.7	
Metro	8.4	13.7	7.2	12.4	8.7	6.3	11.6	
Southeast	9.8	8.0	7.0	11.7	10.7	8.5	23.5	
Southwest	12.9	22.1	11.3	10.0	14.7	8.4	10.3	
Unknown, Not Reported	50.0	0.0	0.0	0.0	0.0	0.0	100.0	

See Technical Appendix for information on health regions and the Modified Kessner Index.

Note: This table uses the Modified Kessner Index.

Natality Section

**Table N-16(a) Number of Births with Low or No Prenatal Care by Mother's Age, County, and Health Region
New Mexico, 2012**

	Mother's age group											Unknown or Not Reported
	All Ages	10-14	15 - 17	18 - 19	20 - 24	25 - 29	30 - 34	35 - 39	40 - 44	45 - 49	50+	
New Mexico	2,759	8	141	309	902	678	460	201	54	3	1	2
County												
Bernalillo	702	2	28	63	198	171	160	60	17	1	1	1
Catron	5	0	0	1	0	3	1	0	0	0	0	0
Chaves	96	0	8	10	42	16	12	7	1	0	0	0
Cibola	95	0	3	7	32	23	16	12	2	0	0	0
Colfax	11	0	1	0	3	5	2	0	0	0	0	0
Curry	98	0	6	9	37	23	14	9	0	0	0	0
De Baca	0	0	0	0	0	0	0	0	0	0	0	0
Dona Ana	461	0	36	63	165	108	57	23	8	1	0	0
Eddy	64	1	2	7	32	16	5	1	0	0	0	0
Grant	14	0	0	2	3	7	0	1	1	0	0	0
Guadalupe	7	0	0	1	3	0	3	0	0	0	0	0
Harding	0	0	0	0	0	0	0	0	0	0	0	0
Hidalgo	5	0	0	0	2	1	1	1	0	0	0	0
Lea	102	0	4	18	36	25	11	7	1	0	0	0
Lincoln	32	0	2	6	9	6	4	3	1	1	0	0
Los Alamos	6	0	0	0	1	2	2	1	0	0	0	0
Luna	51	0	5	7	16	15	2	5	1	0	0	0
McKinley	190	1	6	29	58	39	34	15	8	0	0	0
Mora	1	0	0	0	1	0	0	0	0	0	0	0
Otero	70	0	0	8	14	26	15	5	2	0	0	0
Quay	15	0	1	0	10	2	2	0	0	0	0	0
Rio Arriba	60	0	4	6	17	19	9	4	1	0	0	0
Roosevelt	29	0	0	3	13	8	3	1	1	0	0	0
Sandoval	105	0	6	14	36	27	18	2	2	0	0	0
San Juan	251	3	9	26	95	54	39	21	4	0	0	0
San Miguel	27	0	2	2	7	12	4	0	0	0	0	0
Santa Fe	56	0	5	6	16	13	7	6	3	0	0	0
Sierra	19	0	0	4	8	3	4	0	0	0	0	0
Socorro	33	1	3	3	7	10	6	3	0	0	0	0
Taos	43	0	0	2	12	15	11	3	0	0	0	0
Torrance	19	0	4	2	7	2	2	2	0	0	0	0
Union	5	0	0	0	2	2	1	0	0	0	0	0
Valencia	86	0	6	10	20	25	15	9	1	0	0	0
Unknown, Not Reported	1	0	0	0	0	0	0	0	0	0	0	1
Health Region												
Northwest	536	4	18	62	185	116	89	48	14	0	0	0
Northeast	216	0	12	17	62	68	39	14	4	0	0	0
Metro	912	2	44	89	261	225	195	73	20	1	1	1
Southeast	436	1	23	53	179	96	51	28	4	1	0	0
Southwest	658	1	44	88	215	173	86	38	12	1	0	0
Unknown, Not Reported	1	0	0	0	0	0	0	0	0	0	0	1

See Technical Appendix for information on health regions and the Modified Kessner Index.

Note: This table uses the Modified Kessner Index.

Nativity Section

**Table N-16(b) Percent of Births with Low or No Prenatal Care by Mother's Age, County, and Health Region
New Mexico, 2012**

	Mother's Age Group										Unknown or Not Reported	
	All Ages	10-14	15 - 17	18 - 19	20 - 24	25 - 29	30 - 34	35 - 39	40 - 44	45 - 49		50+
New Mexico	10.2	16.0	14.1	13.6	11.5	9.0	8.6	8.5	9.9	10.3	20.0	15.4
County												
Bernalillo	8.4	20.0	11.5	11.8	9.6	7.1	8.0	6.5	8.6	8.3	100.0	25.0
Catron	22.7	0.0	0.0	50.0	0.0	75.0	12.5	0.0	0.0	0.0	0.0	0.0
Chaves	10.1	0.0	20.5	9.2	12.3	6.7	8.6	10.3	10.0	0.0	0.0	0.0
Cibola	22.1	0.0	23.1	17.1	21.2	19.7	23.5	36.4	33.3	0.0	0.0	0.0
Colfax	9.6	0.0	25.0	0.0	7.9	14.7	8.3	0.0	0.0	0.0	0.0	0.0
Curry	10.6	0.0	13.3	10.8	12.2	8.6	8.8	16.1	0.0	0.0	0.0	0.0
De Baca	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Dona Ana	15.0	0.0	27.9	20.2	16.6	13.5	10.1	11.3	13.8	33.3	0.0	0.0
Eddy	8.0	50.0	5.0	7.3	11.4	7.7	4.1	2.6	0.0	0.0	0.0	0.0
Grant	4.3	0.0	0.0	6.9	2.8	7.7	0.0	5.0	25.0	0.0	0.0	0.0
Guadalupe	14.9	0.0	0.0	25.0	15.8	0.0	37.5	0.0	0.0	0.0	0.0	0.0
Harding	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hidalgo	8.5	0.0	0.0	0.0	9.5	6.3	14.3	14.3	0.0	0.0	0.0	0.0
Lea	8.8	0.0	6.6	13.1	9.7	7.2	6.8	10.8	7.1	0.0	0.0	0.0
Lincoln	16.3	0.0	25.0	26.1	15.5	11.5	15.4	15.0	14.3	100.0	0.0	0.0
Los Alamos	3.4	0.0	0.0	0.0	3.8	4.5	2.9	4.0	0.0	0.0	0.0	0.0
Luna	12.2	0.0	12.2	13.7	11.0	12.9	5.7	23.8	14.3	0.0	0.0	0.0
McKinley	15.3	25.0	13.3	18.7	15.6	13.0	15.4	13.2	29.6	0.0	0.0	0.0
Mora	2.4	0.0	0.0	0.0	7.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Otero	7.9	0.0	0.0	11.1	5.2	9.3	8.9	8.9	14.3	0.0	0.0	0.0
Quay	12.5	0.0	25.0	0.0	20.0	6.3	14.3	0.0	0.0	0.0	0.0	0.0
Rio Arriba	9.8	0.0	15.4	11.8	7.9	11.7	9.5	7.8	10.0	0.0	0.0	0.0
Roosevelt	10.7	0.0	0.0	14.3	14.1	8.7	8.3	7.1	25.0	0.0	0.0	0.0
Sandoval	7.2	0.0	15.4	13.3	9.6	6.2	5.6	1.4	5.4	0.0	0.0	0.0
San Juan	13.5	33.3	15.8	16.5	15.4	10.7	11.7	13.7	14.3	0.0	0.0	0.0
San Miguel	7.7	0.0	12.5	6.1	6.0	11.5	7.1	0.0	0.0	0.0	0.0	0.0
Santa Fe	4.1	0.0	9.3	7.1	4.6	3.9	2.4	3.4	5.3	0.0	0.0	0.0
Sierra	20.9	0.0	0.0	36.4	32.0	11.5	23.5	0.0	0.0	0.0	0.0	0.0
Socorro	15.6	50.0	30.0	12.0	10.6	18.5	17.6	18.8	0.0	0.0	0.0	0.0
Taos	13.5	0.0	0.0	12.5	14.1	19.2	13.4	7.7	0.0	0.0	0.0	0.0
Torrance	11.9	0.0	40.0	12.5	13.0	4.7	9.1	15.4	0.0	0.0	0.0	0.0
Union	9.4	0.0	0.0	0.0	11.8	10.0	11.1	0.0	0.0	0.0	0.0	0.0
Valencia	9.9	0.0	14.3	15.9	8.5	9.7	7.7	14.8	5.9	0.0	0.0	0.0
Unknown, Not Reported	50.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0
Health Region												
Northwest	15.2	30.8	15.7	17.5	16.2	12.6	14.3	16.0	23.0	0.0	0.0	0.0
Northeast	7.0	0.0	10.3	8.2	7.1	8.5	6.0	4.3	4.4	0.0	0.0	0.0
Metro	8.4	18.2	13.2	12.4	9.6	7.2	7.7	6.4	7.9	6.3	100.0	20.0
Southeast	9.8	11.1	11.1	11.0	11.9	7.7	7.8	10.4	7.4	33.3	0.0	0.0
Southwest	12.9	8.3	19.6	17.4	13.2	12.4	9.6	11.4	13.3	33.3	0.0	0.0
Unknown, Not Reported	50.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0

See Technical Appendix for information on health regions and the Modified Kessner Index.

Note: This table uses the Modified Kessner Index.

Natality Section

**Table N-17(a) Number of Births by Trimester Prenatal Care Began, County, and Health Region
New Mexico, 2012**

	Trimester Prenatal Care Began					
	All Births	No Prenatal Care	1-3 Months (first trimester)	4-6 Months (second trimester)	7-9 Months (third trimester)	Unknown or not reported trimester
New Mexico	26,992	341	17,060	5,946	1,527	2,118
County						
Bernalillo	8,379	44	5,305	1,638	318	1074
Catron	22	0	8	8	4	2
Chaves	948	14	643	217	57	17
Cibola	429	7	215	116	67	24
Colfax	115	4	83	21	6	1
Curry	924	7	517	298	73	29
De Baca	12	0	10	2	0	0
Dona Ana	3,071	107	1,853	810	280	21
Eddy	796	5	458	157	42	134
Grant	322	6	250	51	5	10
Guadalupe	47	0	25	14	5	3
Harding	5	0	5	0	0	0
Hidalgo	59	3	34	17	2	3
Lea	1,161	25	637	276	60	163
Lincoln	196	5	116	53	16	6
Los Alamos	178	1	140	27	4	6
Luna	419	13	278	84	34	10
McKinley	1,239	22	684	363	112	58
Mora	41	0	26	13	1	1
Otero	888	15	558	180	29	106
Quay	120	5	68	36	7	4
Rio Arriba	610	8	433	121	32	16
Roosevelt	271	6	141	87	20	17
Sandoval	1,459	5	1,019	246	68	121
San Juan	1,862	21	1,202	464	108	67
San Miguel	352	3	258	59	17	15
Santa Fe	1,358	1	1,067	207	39	44
Sierra	91	2	51	23	13	2
Socorro	211	2	103	72	15	19
Taos	319	2	207	71	33	6
Torrance	160	0	87	34	11	28
Union	53	0	34	15	4	0
Valencia	873	7	545	165	45	111
Unknown, Not Reported	2	1	0	1	0	0
Health Region						
Northwest	3,530	50	2,101	943	287	149
Northeast	3,078	19	2,278	548	141	92
Metro	10,871	56	6,956	2,083	442	1,334
Southeast	4,428	67	2,590	1,126	275	370
Southwest	5,083	148	3,135	1,245	382	173
Unknown, Not Reported	2	1	0	1	0	0

See Technical Appendix for information on health regions and trimester prenatal care began.

Natality Section

**Table N-17(b) Percent of Births by Trimester Prenatal Care Began, County, and Health Region
New Mexico, 2012**

	Trimester Prenatal Care Began				
	No Prenatal Care	1-3 Months (first trimester)	4-6 Months (second trimester)	7-9 Months (third trimester)	Unknown or not reported trimester
New Mexico	1.3	63.2	22.0	5.7	7.8
County					
Bernalillo	0.5	63.3	19.5	3.8	12.8
Catron	0.0	36.4	36.4	18.2	9.1
Chaves	1.5	67.8	22.9	6.0	1.8
Cibola	1.6	50.1	27.0	15.6	5.6
Colfax	3.5	72.2	18.3	5.2	0.9
Curry	0.8	56.0	32.3	7.9	3.1
De Baca	0.0	83.3	16.7	0.0	0.0
Dona Ana	3.5	60.3	26.4	9.1	0.7
Eddy	0.6	57.5	19.7	5.3	16.8
Grant	1.9	77.6	15.8	1.6	3.1
Guadalupe	0.0	53.2	29.8	10.6	6.4
Harding	0.0	100.0	0.0	0.0	0.0
Hidalgo	5.1	57.6	28.8	3.4	5.1
Lea	2.2	54.9	23.8	5.2	14.0
Lincoln	2.6	59.2	27.0	8.2	3.1
Los Alamos	0.6	78.7	15.2	2.2	3.4
Luna	3.1	66.3	20.0	8.1	2.4
McKinley	1.8	55.2	29.3	9.0	4.7
Mora	0.0	63.4	31.7	2.4	2.4
Otero	1.7	62.8	20.3	3.3	11.9
Quay	4.2	56.7	30.0	5.8	3.3
Rio Arriba	1.3	71.0	19.8	5.2	2.6
Roosevelt	2.2	52.0	32.1	7.4	6.3
Sandoval	0.3	69.8	16.9	4.7	8.3
San Juan	1.1	64.6	24.9	5.8	3.6
San Miguel	0.9	73.3	16.8	4.8	4.3
Santa Fe	0.1	78.6	15.2	2.9	3.2
Sierra	2.2	56.0	25.3	14.3	2.2
Socorro	0.9	48.8	34.1	7.1	9.0
Taos	0.6	64.9	22.3	10.3	1.9
Torrance	0.0	54.4	21.3	6.9	17.5
Union	0.0	64.2	28.3	7.5	0.0
Valencia	0.8	62.4	18.9	5.2	12.7
Unknown, Not Reported	50.0	0.0	50.0	0.0	0.0
Health Region					
Northwest	1.4	59.5	26.7	8.1	4.2
Northeast	0.6	74.0	17.8	4.6	3.0
Metro	0.5	64.0	19.2	4.1	12.3
Southeast	1.5	58.5	25.4	6.2	8.4
Southwest	2.9	61.7	24.5	7.5	3.4
Unknown, Not Reported	50.0	0.0	50.0	0.0	0.0

See Technical Appendix for information on health regions and trimester prenatal care began.

Nativity Section

Table N-18 Number and Percent of Births that Occurred and Did Not Occur in New Mexico by County of Residence, 2012

	Total Resident Births	Births Occurring in New Mexico	%	Births Occurring Out- of-State	%
New Mexico	26,992	25,595	94.8	1,397	5.2
Residence County					
Bernalillo	8,379	8,359	99.8	20	0.2
Catron	22	20	90.9	2	9.1
Chaves	948	942	99.4	6	0.6
Cibola	429	429	100.0	0	0.0
Colfax	115	105	91.3	10	8.7
Curry	924	847	91.7	77	8.3
De Baca	12	11	91.7	1	8.3
Dona Ana	3,071	2,270	73.9	801	26.1
Eddy	796	785	98.6	11	1.4
Grant	322	321	99.7	1	0.3
Guadalupe	47	47	100.0	0	0.0
Harding	5	4	80.0	1	20.0
Hidalgo	59	54	91.5	5	8.5
Lea	1,161	1,052	90.6	109	9.4
Lincoln	196	189	96.4	7	3.6
Los Alamos	178	176	98.9	2	1.1
Luna	419	401	95.7	18	4.3
McKinley	1,239	1,207	97.4	32	2.6
Mora	41	41	100.0	0	0.0
Otero	888	842	94.8	46	5.2
Quay	120	109	90.8	11	9.2
Rio Arriba	610	607	99.5	3	0.5
Roosevelt	271	246	90.8	25	9.2
Sandoval	1,459	1,457	99.9	2	0.1
San Juan	1,862	1,687	90.6	175	9.4
San Miguel	352	352	100.0	0	0.0
Santa Fe	1,358	1,353	99.6	5	0.4
Sierra	91	89	97.8	2	2.2
Socorro	211	211	100.0	0	0.0
Taos	319	318	99.7	1	0.3
Torrance	160	159	99.4	1	0.6
Union	53	32	60.4	21	39.6
Valencia	873	871	99.8	2	0.2
Unknown, Not Reported	2	2	250.0	0	0.0

Nativity Section

**Table N-19 Number and Percent of Births by Facility Type and Attendant Type
New Mexico, 1990, 2000, and 2012**

Year/Facility Type	Total Births	Physician		Certified Nurse Midwife		Licensed Midwife		Other and Unknown Attendant Type	
		#	%	#	%	#	%	#	%
2012	26,992	17,889	66.3	7,279	27.0	350	1.3	1,474	5.5
Hospital	26,477	17,868	67.5	7,197	27.2	2	0.0	1,410	5.3
Freestanding Birthing Center	119	0	0.0	74	62.2	40	33.6	5	4.2
Mother's Residence	356	7	2.0	5	1.4	305	85.7	39	11.0
Other	40	14	35.0	3	7.5	3	7.5	20	50.0
2000	27,206	19,407	71.3	7,338	27.0	290	1.1	171	0.6
Hospital	26,780	19,391	72.4	7,304	27.3	6	0.0	79	0.3
Freestanding Birthing Center	101	6	5.9	22	21.8	73	72.3	0	0.0
Mother's Residence	263	5	1.9	9	3.4	195	74.1	54	20.5
Other	62	5	8.1	3	4.8	16	25.8	38	61.3
1990	27,317	24,026	88.0	2,884	10.6	244	0.9	163	0.6
Hospital	26,965	24,004	89.0	2,845	10.6	41	0.2	75	0.3
Freestanding Birthing Center	39	2	5.1	0	0.0	37	94.9	0	0.0
Mother's Residence	268	8	3.0	38	14.2	151	56.3	71	26.5
Other	45	12	26.7	1	2.2	15	33.3	17	37.8

Unknown and not reported facilities are included in Other.

Natality Section

**Table N-20(a) Number of Births by Birth Method, Mother's Age, and Race/Ethnicity
New Mexico, 2012**

Mother's age group	All Births	Vaginal birth after C-section	Primary C- section	Repeat C- section	Other and Unknown Birth Method
All Races					
10 to 14	50	44	0	6	0
15 to 17	998	859	1	130	5
18 to 19	2,267	1,879	7	331	49
20 to 24	7,867	6,132	69	1,200	453
25 to 29	7,503	5,619	111	1,189	574
30 to 34	5,346	3,758	99	1012	465
35 to 39	2,367	1,506	59	559	235
40 to 44	547	309	21	157	59
45 to 49	29	17	0	9	3
50+	5	2	0	2	1
Unknown or Not Reported Age	13	6	0	7	0
All Ages	26,992	20,131	367	4,602	1,844
American Indian or Alaska Native					
10 to 14	12	10	0	2	0
15 to 17	127	105	1	18	1
18 to 19	347	304	1	37	4
20 to 24	1,063	839	13	144	57
25 to 29	875	676	24	120	52
30 to 34	536	375	20	87	46
35 to 39	251	154	8	55	29
40 to 44	62	36	4	17	4
45 to 49	4	3	0	1	0
All Ages	3,277	2,502	71	481	193
Asian or Pacific Islander					
10 to 14	0	0	0	0	0
15 to 17	1	1	0	0	0
18 to 19	12	10	0	2	0
20 to 24	63	49	0	10	4
25 to 29	146	101	0	38	6
30 to 34	192	124	3	48	17
35 to 39	124	80	1	27	15
40 to 44	16	11	1	3	1
45 to 49	1	0	0	1	0
Unknown or Not Reported Age	1	1	0	0	0
All Ages	556	377	5	129	43
Black or African American					
10 to 14	1	1	0	0	0
15 to 17	9	8	0	1	0
18 to 19	40	32	0	7	1
20 to 24	148	107	0	30	11
25 to 29	119	87	2	22	8
30 to 34	98	66	0	21	11
35 to 39	34	19	1	10	4
40 to 44	9	2	0	5	2
45 to 49	1	0	0	1	0
All Ages	459	322	3	97	37
Hispanic					
10 to 14	35	31	0	4	0
15 to 17	717	624	0	89	3
18 to 19	1,430	1,173	4	219	34
20 to 24	4,629	3,600	41	704	282
25 to 29	3,937	2,936	50	618	331
30 to 34	2,473	1,723	42	467	240
35 to 39	1,084	685	26	260	113
40 to 44	261	147	9	69	36
45 to 49	7	3	0	1	3
50+	2	1	0	1	0
Unknown or Not Reported Age	5	1	0	4	0
All Ages	14,580	10,924	172	2,436	1042
White					
10 to 14	2	2	0	0	0
15 to 17	135	113	0	21	1
18 to 19	422	344	2	66	10
20 to 24	1,906	1,489	14	304	98
25 to 29	2,365	1,775	33	383	171
30 to 34	2,005	1,438	34	381	150
35 to 39	852	553	23	202	73
40 to 44	192	109	7	60	16
45 to 49	16	11	0	5	0
50+	2	1	0	0	1
Unknown or Not Reported Age	4	3	0	1	0
All Ages	7,901	5,838	113	1,423	520

See Technical Appendix for information on race/ethnicity and birth method.

Natality Section

**Table N-20(b) Percent of Births by Birth Method, Mother's Age, and Race/Ethnicity
New Mexico, 2012**

Mother's age group	Vaginal	Vaginal birth after C-section	Primary C- section	Repeat C- section	Other and Unknown Birth Method
All Races					
10 to 14	88.0	0.0	12.0	0.0	0.0
15 to 17	86.1	0.1	13.0	0.5	0.3
18 to 19	82.9	0.3	14.6	2.2	0.0
20 to 24	77.9	0.9	15.3	5.8	0.2
25 to 29	74.9	1.5	15.8	7.7	0.1
30 to 34	70.3	1.9	18.9	8.7	0.2
35 to 39	63.6	2.5	23.6	9.9	0.3
40 to 44	56.5	3.8	28.7	10.8	0.2
45 to 49	58.6	0.0	31.0	10.3	0.0
50+	40.0	0.0	40.0	20.0	0.0
Unknown or Not Reported Age	46.2	0.0	53.8	0.0	0.0
All Ages	74.6	1.4	17.0	6.8	0.2
American Indian or Alaska Native					
10 to 14	83.3	0.0	16.7	0.0	0.0
15 to 17	82.7	0.8	14.2	0.8	1.6
18 to 19	87.6	0.3	10.7	1.2	0.3
20 to 24	78.9	1.2	13.5	5.4	0.9
25 to 29	77.3	2.7	13.7	5.9	0.3
30 to 34	70.0	3.7	16.2	8.6	1.5
35 to 39	61.4	3.2	21.9	11.6	2.0
40 to 44	58.1	6.5	27.4	6.5	1.6
45 to 49	75.0	0.0	25.0	0.0	0.0
All Ages	76.4	2.2	14.7	5.9	0.9
Asian or Pacific Islander					
10 to 14	0.0	0.0	0.0	0.0	0.0
15 to 17	100.0	0.0	0.0	0.0	0.0
18 to 19	83.3	0.0	16.7	0.0	0.0
20 to 24	77.8	0.0	15.9	6.3	0.0
25 to 29	69.2	0.0	26.0	4.1	0.7
30 to 34	64.6	1.6	25.0	8.9	0.0
35 to 39	64.5	0.8	21.8	12.1	0.8
40 to 44	68.8	6.3	18.8	6.3	0.0
45 to 49	0.0	0.0	100.0	0.0	0.0
Unknown or Not Reported Age	100.0	0.0	0.0	0.0	0.0
All Ages	67.8	0.9	23.2	7.7	0.4
Black or African American					
10 to 14	100.0	0.0	0.0	0.0	0.0
15 to 17	88.9	0.0	11.1	0.0	0.0
18 to 19	80.0	0.0	17.5	2.5	0.0
20 to 24	72.3	0.0	20.3	7.4	0.0
25 to 29	73.1	1.7	18.5	6.7	0.0
30 to 34	67.3	0.0	21.4	11.2	0.0
35 to 39	55.9	2.9	29.4	11.8	0.0
40 to 44	22.2	0.0	55.6	22.2	0.0
45 to 49	0.0	0.0	100.0	0.0	0.0
All Ages	70.2	0.7	21.1	8.1	0.0
Hispanic					
10 to 14	88.6	0.0	11.4	0.0	0.0
15 to 17	87.0	0.0	12.4	0.4	0.1
18 to 19	82.0	0.3	15.3	2.4	0.0
20 to 24	77.8	0.9	15.2	6.1	0.0
25 to 29	74.6	1.3	15.7	8.4	0.1
30 to 34	69.7	1.7	18.9	9.7	0.0
35 to 39	63.2	2.4	24.0	10.4	0.0
40 to 44	56.3	3.4	26.4	13.8	0.0
45 to 49	42.9	0.0	14.3	42.9	0.0
50+	50.0	0.0	50.0	0.0	0.0
Unknown or Not Reported Age	20.0	0.0	80.0	0.0	0.0
All Ages	74.9	1.2	16.7	7.1	0.0
White					
10 to 14	100.0	0.0	0.0	0.0	0.0
15 to 17	83.7	0.0	15.6	0.1	0.0
18 to 19	81.5	0.5	15.6	0.2	0.0
20 to 24	78.1	0.7	15.9	0.5	0.1
25 to 29	75.1	1.4	16.2	0.7	0.1
30 to 34	71.7	1.7	19.0	0.7	0.1
35 to 39	64.9	2.7	23.7	0.9	0.1
40 to 44	56.8	3.6	31.3	0.8	0.0
45 to 49	68.8	0.0	31.3	0.0	0.0
50+	50.0	0.0	0.0	5.0	0.0
Unknown or Not Reported Age	75.0	0.0	25.0	0.0	0.0
All Ages	73.9	1.4	18.0	0.7	0.1

See Technical Appendix for information on race/ethnicity and birth method.

Nativity Section

**Table N-21 Number of Births by Birth Method, County, and Health Region
New Mexico, 2012**

	All Births	Vaginal birth after C-section	Primary C- section	Repeat C- section	Other and Unknown or Not Reported Birth Method	
New Mexico	26,992	20,131	367	4,602	1,844	48
County						
Bernalillo	8,379	6,414	154	1,558	250	3
Catron	22	13	2	2	3	2
Chaves	948	662	2	189	95	0
Cibola	429	306	10	83	29	1
Colfax	115	79	1	24	11	0
Curry	924	695	8	123	96	2
De Baca	12	10	0	1	1	0
Dona Ana	3,071	2,144	26	510	389	2
Eddy	796	532	5	153	106	0
Grant	322	245	1	35	41	0
Guadalupe	47	39	0	6	2	0
Harding	5	5	0	0	0	0
Hidalgo	59	36	0	12	9	2
Lea	1,161	885	6	179	89	2
Lincoln	196	143	2	34	17	0
Los Alamos	178	129	3	39	7	0
Luna	419	297	2	69	50	1
McKinley	1,239	959	28	159	62	31
Mora	41	38	0	1	2	0
Otero	888	650	11	148	78	1
Quay	120	95	0	12	13	0
Rio Arriba	610	470	13	92	35	0
Roosevelt	271	201	0	38	32	0
Sandoval	1,459	1,089	24	275	70	1
San Juan	1,862	1,311	31	313	207	0
San Miguel	352	282	2	49	19	0
Santa Fe	1,358	1,110	13	200	35	0
Sierra	91	71	0	12	8	0
Socorro	211	138	4	51	18	0
Taos	319	243	3	44	29	0
Torrance	160	122	1	32	5	0
Union	53	37	0	12	4	0
Valencia	873	680	15	146	32	0
Unknown, Not Reported	2	1	0	1	0	0
Health Region						
Northwest	3,530	2,576	69	555	298	32
Northeast	3,078	2,432	35	467	144	0
Metro	10,871	8,305	194	2,011	357	4
Southeast	4,428	3,223	23	729	449	4
Southwest	5,083	3,594	46	839	596	8
Unknown, Not Reported	2	1	0	1	0	0

See Technical Appendix for information on health regions and birth method.

Mortality Section




MORTALITY SECTION

ALL CAUSES OF DEATH

In 2012, 16,640 New Mexico resident deaths were reported. The age-adjusted death rate was 749.4 deaths per 100,000 standard population (Table M-1(b)).

The majority (68.2%) of the 2012 deaths were to individuals age 65 years or older (61.6% of all male deaths and 75.9% of all female deaths). A larger percentage of males died from ages 35 to 64 years (30.5%) than did females (20.2%). Females were more likely to have died at ages 85 years or older (36.1%) compared to males (19.7%). More than twice the percentage of males died from ages 15 to 34 years (6.0%) than did females in the same age group (2.6%). For residents under 15 years of age, the distribution of deaths for both males and females was similar (Figure M-1 and Table M-1(a)).

**Figure M-1, Percent Deaths by Age Group and Sex
New Mexico, 2012**




Population Note: Death rates for years 2001-2011 differ slightly from those shown in reports published in previous years. This reflects adjustments to 2001-2011 population estimates following the 2010 Census. See the *Technical Appendix* for more information.

Mortality Section

In 2012, the overall age-adjusted death rate for New Mexico males was 906.8 per 100,000 population compared to 612.8 for females. The top death rates among males exceeded 1 death per 100 persons in 2012 for American Indian or Alaska Natives (1197.7) and Black or African Americans (1048.6). The death rate for Asian or Pacific Islander males (738.6), White males (857.8), and Hispanic males (903.0) were lower than the overall rate for New Mexico males (Figure M-2).

**Figure M-2, Death Rates for All Causes by Sex and Race/Ethnicity
New Mexico, 2012**




There was little difference among the death rates for Black or African American females (596.3), Hispanic females (602.9), and White females (609.1). Among New Mexico females, the death rate for Asian or Pacific Islander females (395.4) was the lowest death rate for all racial/ethnic groups in the state (Figure M-2).

Mortality Section

American Indian or Alaska Natives died in New Mexico at earlier ages than other groups in 2012. Over 52% of American Indian or Alaska Natives died from ages 15 to 65 years while the percentage was 38% or less for other groups in that age range. The actual numbers of deaths by race/ethnicity are presented in Table M-1(a).

**Figure M-3, Death Rates for All Causes by Race/Ethnicity
New Mexico, 2004-2012**




The Asian or Pacific Islander population has had the lowest age-adjusted death rates for all causes of death since 2004. Due to the small number of deaths and the small population of Asian or Pacific Islanders, death rates for this group fluctuate greatly from year-to-year and should be interpreted with caution (Figure M-3).

Note on Race/Ethnicity: Death rates by Race or Ethnicity differ from those shown in reports published in previous years. This reflects a revision to the Department of Health’s standard definition for presentation of race/ethnicity as a single construct. See the *Technical Appendix* for more information.

From 2003 to 2010 New Mexico’s overall age-adjusted death rates were 4.1% to 7.2% less than comparable rates for the United States (Figure M-4).

Mortality Section

**Figure M-4, Death Rates
New Mexico, 2003-2012, and United States, 2003-2010**



Population Note: In 2013 the National Center for Health Statistics updated its intercensal population estimates and released revised death rates for the U.S. that differ from rates published in previous reports.

LEADING CAUSES OF DEATH (RANKED BY NUMBERS OF DEATHS)

For New Mexico, the 2012 leading causes of death were:


- | | |
|---------------------------------------|--|
| 1) Malignant Neoplasms (Cancer) | 6) Diabetes Mellitus |
| 2) Heart Disease | 7) Chronic Liver Disease and Cirrhosis |
| 3) Accidents (Unintentional Injuries) | 8) Intentional Self-harm (Suicide) |
| 4) Chronic Lower Respiratory Diseases | 9) Alzheimer's Disease |
| 5) Cerebrovascular Diseases (Stroke) | 10) Influenza and Pneumonia |

For the United States, the 2010 leading causes of death were:

- | | |
|---------------------------------------|--|
| 1) Heart Disease | 7) Diabetes Mellitus |
| 2) Malignant Neoplasms (Cancer) | 8) Nephritis, Nephrotic Syndrome, and Nephrosis (Kidney Disease) |
| 3) Chronic Lower Respiratory Diseases | 9) Influenza and Pneumonia |
| 4) Cerebrovascular Diseases (Stroke) | 10) Intentional Self-harm (Suicide) |
| 5) Accidents (Unintentional Injuries) | |
| 6) Alzheimer's Disease | |

Mortality Section

**Figure M-5, Leading Causes of Death
New Mexico, 2012, and United States, 2010**




In 2010 and 2012 Malignant Neoplasms (Cancer) was the leading cause of death in New Mexico based on numbers of deaths. In 2012, 40.5% of all deaths were attributed to Cancer and Heart Disease. New Mexico's share of deaths from Accidents (Unintentional Injuries) (8.0%) and Chronic Liver Disease (2.6%) was notably higher compared to that for United States, 4.9% and 1.3%. This was also the case for Diabetes Mellitus and Suicide, while Alzheimer's Disease and Cerebrovascular Disease made up a larger proportion of deaths for the United States (Figure M-5).

Compared with 2003, New Mexico 2012 death rates for Heart Disease and Malignant Neoplasms declined while the death rate for Accidents (Unintentional Injuries) increased. In 2012, the Cerebrovascular Disease death rate (30.4) was at the lowest point in ten years (Figure M-6).

Mortality Section


**Figure M-6, Death Rates by Cause
New Mexico, 2003-2012**




The death rates for the top three causes of death for New Mexicans have varied among the racial/ethnic groups in the state. Black or African Americans, Whites, and Hispanics have generally had the highest rates for Cancer and Heart Disease. American Indian or Alaska Natives have had the highest death rates for Unintentional Injuries (Accidents) (Figures M-7(a-c)).

Mortality Section

**Figure M-7(a), Malignant Neoplasm Death Rates by Race/Ethnicity
New Mexico, 2004-2006, 2007-2009, and 2010-2012**




**Figure M-7(b), Heart Disease Death Rates by Race/Ethnicity,
New Mexico, 2004-2006, 2007-2009, and 2010-2012**



Mortality Section

**Figure M-7(c), Unintentional Injury Death Rates by Race/Ethnicity
New Mexico, 2004-2006, 2007-2009, and 2010-2012**



SELECTED CAUSES

Among the 39 selected causes of death, more women than men died of Alzheimer’s Disease. More men than women died of Major Cardiovascular Diseases. More than twice as many men died of Chronic Liver Disease and Cirrhosis than women, and male HIV deaths occurred at 7 times the rate for women. More males died of Cancer than females in 2012. 283 Influenza and Pneumonia deaths were reported in 2012 (Table M-16).

INJURY DEATHS AND ALCOHOL- AND DRUG-INDUCED DEATHS

The death rate for Total Injuries in New Mexico was 65% higher than the United States rate. The New Mexico Motor Vehicle Accident rate was 60% higher than the U.S. rate. The Homicide rate was 25% higher and the Suicide rate was almost 75% higher than the U.S. rate (Table M-20).


In New Mexico death rates for Suicide, Homicide, and Motor Vehicle Accidents were higher for males than females. For Homicide the male-female rate ratio was 5 to 1, for Suicide the ratio was 3.5 to 1; and for Motor Vehicle Accidents the ratio was 2.5 to 1 (Table M-20).

Mortality Section

Among deaths from Unintentional Injuries (Figure M-8), the 2012 death rate for Motor Vehicle Accidents (18.1) interrupted the downward trend since 2006 with a 26% increase over 2011. The generally increasing rates for Poisonings deaths since 2004 saw a slight decline of 8% between 2011 and 2012. Much of the increase in the death rate from 2003 to 2012 for Unintentional Poisoning is the result of prescription drug overdose. (<http://nmhealth.org/ERD/pdf/ER%20prescription%20drug%20overdose%20112009.pdf>).


Falls death rates have remained relatively stable since 2010 (Figure M-8).

Figure M-8, Death Rates for Unintentional Injuries by Type, New Mexico, 2003-2012



Mortality Section


**Figure M-9, Death Rates for Intentional Injuries by Type,
New Mexico, 2003-2012**



Intentional firearm injury deaths (suicide, homicide, and legal intervention) accounted for 54.9% of intentional injury deaths, while all firearm injury deaths made up 16.2% of New Mexico injury deaths (intentional and unintentional) in 2012. The actual numbers of deaths due to intentional and unintentional injuries are presented in Table M-19, and deaths due to firearms in Table M-21 and Figure M-10. In 2012 suicide deaths accounted for 71.3% of all firearm injury deaths, followed by homicide deaths (26.8%), and legal intervention and accidents (each 0.3%). For another 1.3% of firearm injury deaths, the intent could not be determined.

Mortality Section

**Figure M-10, Percent of Firearm Injury Deaths by Manner
New Mexico, 2012**



Approximately 2.1% of all injury deaths in the state occurred at work, compared to 2.3% for the United States. The total number of deaths caused by injury at work in New Mexico was 41 in 2012, with 38 of these occurring among males (Table M-19 and Table M-21).

New Mexico's 2012 age-adjusted death rate for alcohol-induced deaths (21.8) was nearly 3 times higher than the 2010 national rate (7.6). New Mexico's rate for drug-induced deaths (24.6) was almost twice as high as the U.S. rate (12.9) (Table M-22).

MATERNAL MORTALITY

Maternal mortality includes deaths that were the result of, or aggravated by, pregnancy or pregnancy management, and occurred within 42 days of pregnancy and excludes all external injury deaths. (See the *Technical Appendix*.)

For the time period 2005-2008, the maternal death rate was 10.9 per 100,000 live births in New Mexico. The maternal mortality rate for 2009-2012 was 17.1 per 100,000 live births (Table M-23). The numbers of maternal deaths in New Mexico are so few each year that even with reporting 4-year totals for death counts, the rates presented in Table M-23 should be interpreted with caution.


Mortality Section

INFANT MORTALITY

In 2012, 186 infant deaths to New Mexico residents were reported (Table M-24(a)). Infant deaths are defined as deaths before the age of one year. The 2012 infant mortality rate was 6.9 infant deaths per 1,000 live births. This rate was a 32.7% increase over the 2011 New Mexico rate (5.2). In 2012 the infant mortality rate in New Mexico was higher than the 2010 U.S. rate of 6.2 for the first time in more than a decade.


A historical perspective is provided in Figure M-11. In 1930, New Mexico's infant mortality rate of 145.4 was 125.1% higher than the national rate. By the early 1970's, however, New Mexico had achieved parity with the United States and by the 1980's the rate had dropped below the United States rate. With the exception of the 1994 rate, the state's infant mortality rate has remained below the national rate every year since 1980 until the 2012 rate interrupted that relationship. The historical decline in infant mortality is due to the role of public health outreach, improved education about pregnancy and childbirth, and expanded medical services, including advancements in antibiotics.

**Figure M-11, Infant Mortality Rates
New Mexico, 1930-2012, and United States, 1930-2010**



Mortality Section

**Figure M-12, Infant Mortality Rates by Race/Ethnicity
New Mexico, 2012**




Infant deaths can be broken down into two age categories: (1) neonatal deaths -- less than 28 days old and (2) postneonatal deaths -- at least 28 days but less than one year old. In 2012, 67.8% of infant deaths occurred during the neonatal period, and 32.2% occurred during the postneonatal period (Table M-24(a)).

Asian or Pacific Islander infants had the highest neonatal mortality rate (5.4) and Black infants had the highest postneonatal mortality rate (6.5). Due to small numbers, the Asian or Pacific Islander and Black or African American rates may not be statistically reliable and should be interpreted with caution (Figure M-12 and Table M-24(b)).


From 1996 until 2011, the state's neonatal infant mortality rate, like the overall infant mortality rate, remained below the United States rate, but rose above the 2010 national rate in 2012. State and national postneonatal rates fluctuated but remained close to one another (Figure 13(a,b)).

Mortality Section

**Figure M-13(a), Neonatal Mortality Rates
New Mexico, 1996-2012, and United States, 1996-2010**



**Figure M-13(b), Postneonatal Mortality Rates
New Mexico, 1996-2012, and United States, 1996-2010**




Of the 186 New Mexico infant deaths in 2012, 48 were caused by congenital malformations (birth defects), and 36 were caused by disorders related to length of gestation or birthweight (Table M-25).

DISPOSITION OF REMAINS

Since 2006, the percentage of persons who died in the state and then were cremated has exceeded the percentage of persons who were buried. The percent of burials declined from 49.5% in 2003 to 36.6% in 2012; whereas, cremations increased from 44.9% in 2003 to 54.9% in 2012 (Table M-26).

**Figure M-14, Percent Distribution by Method of Disposition
New Mexico Occurrence, 2012**



Mortality Section

**Table M-1(a) Number of Deaths by Age, Race/Ethnicity, and Sex
New Mexico, 2012, and United States, 2010**

	All Ages	Less than 1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	Unknown or Not Stated
All Races													
Male	8,967	112	24	26	210	331	412	869	1,452	1,636	2,129	1,764	2
Female	7,673	74	15	10	71	133	216	501	830	1,180	1,875	2,767	1
Both Sexes	16,640	186	39	36	281	464	628	1,370	2,282	2,816	4,004	4,531	3
American Indian or Alaska Native													
Male	735	5	1	2	47	74	76	110	131	114	115	60	0
Female	506	11	2	1	17	31	45	57	55	85	97	105	0
Both Sexes	1,241	16	3	3	64	105	121	167	186	199	212	165	0
Asian or Pacific Islander													
Male	58	3	0	1	0	2	4	9	8	4	14	13	0
Female	43	0	0	1	1	2	0	2	4	6	10	17	0
Both Sexes	101	3	0	2	1	4	4	11	12	10	24	30	0
Black or African American													
Male	174	5	1	0	3	10	7	18	39	32	41	18	0
Female	104	0	2	0	0	0	3	7	20	15	27	30	0
Both Sexes	278	5	3	0	3	10	10	25	59	47	68	48	0
Hispanic													
Male	3,153	61	14	16	110	152	206	388	540	550	657	459	0
Female	2,486	46	10	6	42	66	94	198	271	387	636	730	0
Both Sexes	5,639	107	24	22	152	218	300	586	811	937	1,293	1,189	0
White													
Male	4,799	38	8	7	49	91	116	340	718	925	1295	1211	1
Female	4,523	17	1	1	11	33	74	236	478	684	1,103	1,885	0
Both Sexes	9,322	55	9	8	60	124	190	576	1196	1609	2,398	3,096	1
Unknown Race/Ethnicity													
Male	48	0	0	0	1	2	3	4	16	11	7	3	1
Female	11	0	0	1	0	1	0	1	2	3	2	0	1
Both Sexes	59	0	0	1	1	3	3	5	18	14	9	3	2
United States													
All Races													
Male	1,232,432	13,702	2,460	3,054	21,790	29,192	43,434	112,018	189,295	229,704	311,830	275,866	87
Female	1,236,003	10,884	1,856	2,225	7,761	13,067	26,599	71,189	121,507	177,447	313,821	489,608	39
Both Sexes	2,468,435	24,586	4,316	5,279	29,551	42,259	70,033	183,207	310,802	407,151	625,651	765,474	126

See Technical Appendix for information on race/ethnicity.

Mortality Section

**Table M-1(b) Death Rates by Age, Race/Ethnicity, and Sex
New Mexico, 2012, and United States, 2010**

	All Ages	Less										
		than 1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
All Races												
Male	906.8	753.0	40.3	17.9	140.2	240.2	330.4	616.6	1,136.7	2,059.2	5,232.4	15,021.4
Female	612.8	520.0	26.3	7.1	49.7	100.4	172.4	340.4	602.1	1,321.1	3,700.9	13,090.2
Both Sexes	749.4	639.1	33.5	12.6	96.0	171.7	251.2	475.5	859.2	1,668.5	4,383.1	13,779.9
American Indian or Alaska Native												
Male	1197.7	318.1	15.9	12.7	285.9	588.9	730	942.8	1,780.4	2,887.5	6,506.4	12,090.2
Female	678.8	706.7	32.1	6.4	104.2	239.1	399.7	442.1	612.5	1,581.9	3,638.4	11,205.4
Both Sexes	908.4	511.4	24.0	9.5	195.4	411.2	558.4	680.0	1,138.5	2,134.9	4,781.8	11,511.8
Asian or Pacific Islander												
Male	738.6	1,652.70	0.0	53.4	0.0	75.3	202.1	490.1	573.6	540.1	5088.1	18375.4
Female	395.4	0.0	0.0	52.7	41.1	65.3	0.0	86.4	188.7	583.1	1851.8	14412.9
Both Sexes	527.6	857.2	0.0	53.1	20.4	69.9	88.3	265.0	341.5	565.1	2944.2	15898.6
Black or African American												
Male	1048.6	1,450.20	72.5	0.0	71.2	277.1	227.9	574.7	1,556.7	2,529.7	7,949.1	13,104.2
Female	596.3	0.0	162.3	0.0	0.0	0.0	134.4	254.9	1,018.5	1,345.2	3,755.6	11,929.6
Both Sexes	817.1	765.9	114.9	0.0	41.8	165	188.5	425.3	1,320.2	1,974.8	5,507.4	12,344.5
Hispanic												
Male	903.0	689.7	39.6	19.6	145.7	223.4	363.2	554.5	1,255.5	2,194.3	5,180.8	14,013.5
Female	602.9	539.6	29.3	7.6	56.9	98.9	165.7	273.9	588.2	1,322.4	3,843.2	12,673.2
Both Sexes	742.8	616.0	34.5	13.6	101.8	161.8	264.4	411.9	910.4	1,724.6	4,423.5	13,159.1
White												
Male	857.8	966.7	50.9	16.4	95.7	178.7	220.8	625.9	977.2	1,909.9	5,088.7	15,598.7
Female	609.1	462.9	6.8	2.5	23.2	69.8	140.9	414.5	607.2	1,301.9	3,653.6	13,396.0
Both Sexes	724.8	723.4	29.6	9.6	60.9	126.3	180.8	517.7	785.8	1,593.6	4,310.0	14,179.2
United States												
All Races												
Male	812.0	680.2	29.6	14.6	97.6	141.5	212.5	505.9	1,075.5	2,275.1	5,693.7	15,414.3
Female	787.4	564.0	23.3	11.1	36.4	64.0	128.9	311.4	643.5	1,527.5	4,137.7	13,219.2
Both Sexes	799.5	623.4	26.5	12.9	67.7	102.9	170.5	407.1	851.9	1,875.1	4,790.2	13,934.3

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

Age-adjusted death rates (all ages) are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates and race/ethnicity. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution. See numbers in Table M-1(a).

Mortality Section

**Table M-2(a) Number of Deaths by Age, Sex, County, and Health Region
New Mexico, 2012, and United States, 2010
Both Sexes**

	All Ages	Less than											Unknown or Not Stated
		1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	
United States	2,468,435	24,586	4,316	5,279	29,551	42,259	70,033	183,207	310,802	407,151	625,651	765,474	126
New Mexico	16,640	186	39	36	281	464	628	1,370	2,282	2,816	4,004	4,531	3
County													
Bernalillo	5,097	56	12	7	72	124	169	400	712	794	1,219	1,532	0
Catron	53	0	0	0	0	2	5	3	10	11	13	9	0
Chaves	612	6	1	1	7	11	18	45	70	100	163	190	0
Cibola	249	2	1	3	3	7	14	26	31	32	71	58	1
Colfax	154	1	1	1	2	0	2	9	15	34	42	47	0
Curry	384	7	0	0	3	11	10	28	50	77	94	104	0
De Baca	38	0	0	0	0	0	1	3	5	9	7	13	0
Dona Ana	1,488	16	6	5	18	38	49	114	193	283	384	382	0
Eddy	534	2	2	0	9	16	20	39	67	100	124	155	0
Grant	343	2	1	0	7	3	6	23	39	70	84	108	0
Guadalupe	33	0	0	0	1	0	2	3	6	8	9	4	0
Harding	7	0	0	0	0	0	0	1	0	2	4	0	0
Hidalgo	44	0	0	0	1	1	0	5	2	5	18	12	0
Lea	533	12	2	2	5	16	15	41	65	100	126	149	0
Lincoln	181	5	1	0	1	5	5	19	29	36	44	36	0
Los Alamos	127	0	0	1	1	2	2	5	9	14	26	67	0
Luna	273	2	0	0	3	8	3	14	38	56	75	74	0
McKinley	553	7	2	2	30	34	47	64	94	71	108	94	0
Mora	56	0	0	0	3	1	3	6	8	9	10	16	0
Otero	547	3	1	2	5	9	20	41	95	98	158	115	0
Quay	116	4	0	0	2	2	1	14	14	18	30	31	0
Rio Arriba	400	3	2	1	17	20	38	52	50	64	75	78	0
Roosevelt	171	3	0	0	2	2	6	6	25	32	46	49	0
Sandoval	907	12	2	4	14	26	42	74	122	146	199	266	0
San Juan	915	14	1	1	31	53	50	81	125	152	195	212	0
San Miguel	278	2	0	1	4	7	9	36	46	48	65	60	0
Santa Fe	1,102	11	2	2	20	30	44	100	157	163	240	333	0
Sierra	227	2	0	1	0	0	2	10	35	48	67	62	0
Socorro	165	3	0	0	4	2	5	12	21	42	36	40	0
Taos	290	4	1	0	4	15	16	28	41	47	67	67	0
Torrance	151	0	0	0	1	4	3	14	28	36	33	32	0
Union	41	0	0	1	0	1	0	4	3	9	10	13	0
Valencia	568	7	1	1	11	14	21	51	74	104	164	119	1
Unknown County	3	0	0	0	0	0	0	0	2	0	0	0	1
Health Region													
Northwest	1,717	23	4	6	64	94	111	171	250	255	374	364	1
Northeast	2,488	21	6	7	52	76	116	243	336	396	546	689	0
Metro	6,723	75	15	12	98	168	235	539	936	1,080	1,615	1,949	1
Southeast	2,569	39	6	3	29	63	76	195	325	472	634	727	0
Southwest	3,140	28	8	8	38	63	90	222	433	613	835	802	0
Unknown Region	3	0	0	0	0	0	0	0	2	0	0	0	1

See Technical Appendix for information on health regions.

Mortality Section

**Table M-2(b) Number of Deaths by Age, Sex, County, and Health Region
New Mexico, 2012, and United States, 2010**

Females

	Less than												Unknown or Not Stated
	All Ages	1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	
United States	1,236,003	10,884	1,856	2,225	7,761	13,067	26,599	71,189	121,507	177,447	313,821	489,608	39
New Mexico	7,673	74	15	10	71	133	216	501	830	1,180	1,875	2,767	1
County													
Bernalillo	2,406	15	5	0	18	35	70	133	266	330	570	964	0
Catron	25	0	0	0	0	2	4	1	2	4	5	7	0
Chaves	308	2	1	0	2	3	10	17	28	41	86	118	0
Cibola	104	1	1	0	1	2	8	7	7	12	26	38	1
Colfax	68	0	0	0	0	0	1	1	3	17	20	26	0
Curry	207	3	0	0	0	2	4	13	27	37	48	73	0
De Baca	17	0	0	0	0	0	1	2	2	2	3	7	0
Dona Ana	703	6	1	4	4	12	15	48	75	112	198	228	0
Eddy	258	1	0	0	1	6	6	15	30	46	56	97	0
Grant	160	1	0	0	3	1	2	5	16	26	39	67	0
Guadalupe	14	0	0	0	0	0	2	2	1	2	6	1	0
Harding	2	0	0	0	0	0	0	0	0	0	0	2	0
Hidalgo	21	0	0	0	1	0	0	2	0	1	6	11	0
Lea	250	6	2	2	1	4	5	16	21	46	63	84	0
Lincoln	73	4	0	0	1	1	2	13	6	8	21	17	0
Los Alamos	72	0	0	0	0	1	0	3	4	6	14	44	0
Luna	111	0	0	0	2	2	1	3	15	24	28	36	0
McKinley	213	5	0	1	10	6	9	24	26	28	52	52	0
Mora	27	0	0	0	2	0	1	2	3	4	6	9	0
Otero	236	3	0	0	0	1	6	15	33	36	70	72	0
Quay	57	2	0	0	0	0	1	9	4	6	12	23	0
Rio Arriba	176	2	1	0	3	8	12	17	16	29	34	54	0
Roosevelt	77	1	0	0	1	1	2	2	9	19	19	23	0
Sandoval	423	3	1	2	1	9	16	24	47	62	103	155	0
San Juan	421	7	0	0	8	12	19	35	48	70	100	122	0
San Miguel	116	2	0	0	1	3	3	13	14	17	27	36	0
Santa Fe	517	2	1	1	3	9	8	41	57	73	101	221	0
Sierra	91	0	0	0	0	0	1	4	11	20	25	30	0
Socorro	78	1	0	0	1	1	0	0	9	26	16	24	0
Taos	117	2	1	0	1	3	2	6	14	22	27	39	0
Torrance	61	0	0	0	0	2	1	4	9	16	15	14	0
Union	21	0	0	0	0	0	0	2	1	5	6	7	0
Valencia	243	5	1	0	6	7	4	22	26	33	73	66	0
Health Region													
Northwest	738	13	1	1	19	20	36	66	81	110	178	212	1
Northeast	1,130	8	3	1	10	24	29	87	113	175	241	439	0
Metro	3,133	23	7	2	25	53	91	183	348	441	761	1,199	0
Southeast	1,247	19	3	2	6	17	31	87	127	205	308	442	0
Southwest	1,425	11	1	4	11	19	29	78	161	249	387	475	0

See Technical Appendix for information on health regions.

Mortality Section

**Table M-2(c) Number of Deaths by Age, Sex, County, and Health Region
New Mexico, 2012, and United States, 2010
Males**

	All Ages	Less than										Unknown or Not Stated	
		1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84		85+
United States	1,232,432	13,702	2,460	3,054	21,790	29,192	43,434	112,018	189,295	229,704	311,830	275,866	87
New Mexico	8,967	112	24	26	210	331	412	869	1452	1,636	2,129	1,764	2
County													
Bernalillo	2,691	41	7	7	54	89	99	267	446	464	649	568	0
Catron	28	0	0	0	0	0	1	2	8	7	8	2	0
Chaves	304	4	0	1	5	8	8	28	42	59	77	72	0
Cibola	145	1	0	3	2	5	6	19	24	20	45	20	0
Colfax	86	1	1	1	2	0	1	8	12	17	22	21	0
Curry	177	4	0	0	3	9	6	15	23	40	46	31	0
De Baca	21	0	0	0	0	0	0	1	3	7	4	6	0
Dona Ana	785	10	5	1	14	26	34	66	118	171	186	154	0
Eddy	276	1	2	0	8	10	14	24	37	54	68	58	0
Grant	183	1	1	0	4	2	4	18	23	44	45	41	0
Guadalupe	19	0	0	0	1	0	0	1	5	6	3	3	0
Harding	5	0	0	0	0	0	0	0	1	0	2	2	0
Hidalgo	23	0	0	0	0	1	0	3	2	4	12	1	0
Lea	283	6	0	0	4	12	10	25	44	54	63	65	0
Lincoln	108	1	1	0	0	4	3	6	23	28	23	19	0
Los Alamos	55	0	0	1	1	1	2	2	5	8	12	23	0
Luna	162	2	0	0	1	6	2	11	23	32	47	38	0
McKinley	340	2	2	1	20	28	38	40	68	43	56	42	0
Mora	29	0	0	0	1	1	2	4	5	5	4	7	0
Otero	311	0	1	2	5	8	14	26	62	62	88	43	0
Quay	59	2	0	0	2	2	0	5	10	12	18	8	0
Rio Arriba	224	1	1	1	14	12	26	35	34	35	41	24	0
Roosevelt	94	2	0	0	1	1	4	4	16	13	27	26	0
Sandoval	484	9	1	2	13	17	26	50	75	84	96	111	0
San Juan	494	7	1	1	23	41	31	46	77	82	95	90	0
San Miguel	162	0	0	1	3	4	6	23	32	31	38	24	0
Santa Fe	585	9	1	1	17	21	36	59	100	90	139	112	0
Sierra	136	2	0	1	0	0	1	6	24	28	42	32	0
Socorro	87	2	0	0	3	1	5	12	12	16	20	16	0
Taos	173	2	0	0	3	12	14	22	27	25	40	28	0
Torrance	90	0	0	0	1	2	2	10	19	20	18	18	0
Union	20	0	0	1	0	1	0	2	2	4	4	6	0
Valencia	325	2	0	1	5	7	17	29	48	71	91	53	1
Unknown County	3	0	0	0	0	0	0	0	2	0	0	0	1
Health Region													
Northwest	979	10	3	5	45	74	75	105	169	145	196	152	0
Northeast	1,358	13	3	6	42	52	87	156	223	221	305	250	0
Metro	3,590	52	8	10	73	115	144	356	588	639	854	750	1
Southeast	1,322	20	3	1	23	46	45	108	198	267	326	285	0
Southwest	1,715	17	7	4	27	44	61	144	272	364	448	327	0
Unknown Region	3	0	0	0	0	0	0	0	2	0	0	0	2

See Technical Appendix for information on health regions.

Mortality Section

**Table M-3(a) Death Rates by Age, Sex, County, and Health Region
New Mexico, 2012, and United States, 2010
Both Sexes**

	All ages	Less										
		than 1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
United States	747.0	623.4	26.5	12.9	67.7	102.9	170.5	407.1	851.9	1,875.1	4,790.2	13,934.3
New Mexico	749.4	639.1	33.5	12.6	96.0	171.7	251.2	475.5	859.2	1,668.5	4,383.1	13,779.9
County												
Bernalillo	730.5	610.4	32.7	7.9	74.4	127.0	198.8	428.5	864.7	1,660.4	4,516.9	13,787.4
Catron	1,071.6	0.0	0.0	0.0	0.0	745.4	2,078.5	549.8	1,138.4	1,524.4	4,316.2	10,636.2
Chaves	815.0	567.8	23.7	10.0	71.4	136.8	242.2	531.7	925.7	2,022.6	5,071.3	14,391.7
Cibola	929.9	533.7	66.7	81.9	80.8	189.5	396.7	660.1	934.9	1,512.9	6,102.4	16,856.8
Colfax	769.2	713.1	178.3	65.6	140.0	0.0	143.7	437.1	668.5	2,108.9	4,795.0	14,546.4
Curry	833.6	800.8	0.0	0.0	39.5	147.7	168.7	457.1	1,033.1	2,465.8	4,925.6	15,098.9
De Baca	1,124.7	0.0	0.0	0.0	0.0	0.0	569.2	1,018.0	1,517.3	3,579.1	4,615.4	20,582.0
Dona Ana	666.2	519.9	48.7	16.5	49.4	139.9	207.6	435.5	819.8	1,502.9	3,708.9	12,803.7
Eddy	857.7	257.9	64.5	0.0	125.5	230.7	309.0	503.9	965.4	2,403.0	4,781.2	14,600.4
Grant	768.3	572.8	71.6	0.0	211.2	105.6	216.0	601.7	808.6	1,899.7	3,922.5	14,489.8
Guadalupe	599.0	0.0	0.0	0.0	180.4	0.0	363.2	403.8	1,021.5	1,922.2	3,554.9	5,480.6
Harding	345.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	629.2	0.0	2,469.6	11,581.1
Hidalgo	733.9	0.0	0.0	0.0	154.6	216.2	0.0	723.0	304.0	1,097.7	6,475.3	12,655.7
Lea	919.9	991.1	41.3	19.1	51.8	165.4	184.6	484.2	973.2	2,525.5	5,173.9	18,863.2
Lincoln	649.6	2,591.5	129.6	0.0	51.7	281.6	247.9	597.1	754.3	1,219.6	3,174.0	9,787.4
Los Alamos	539.7	0.0	0.0	39.4	61.0	114.6	85.8	150.1	314.5	924.9	2,936.7	16,480.5
Luna	801.9	553.5	0.0	0.0	89.9	303.2	113.5	469.3	1,205.6	2,040.3	4,303.3	14,575.3
McKinley	892.7	558.9	39.9	16.4	240.9	371.1	542.5	656.9	1,238.9	1,644.4	4,669.2	12,497.3
Mora	939.7	0.0	0.0	0.0	598.3	227.7	547.8	837.2	946.6	1,635.0	3,205.1	18,675.6
Otero	760.6	313.6	26.1	23.3	54.2	105.9	277.6	477.4	1,218.3	1,683.3	4,760.6	12,192.6
Quay	957.0	3,967.5	0.0	0.0	215.7	232.5	104.5	1,097.7	996.7	1,659.1	5,037.8	16,337.3
Rio Arriba	947.3	559.3	93.2	18.5	326.6	432.5	777.8	874.9	901.6	1,814.7	4,105.5	12,947.0
Roosevelt	839.4	921.6	0.0	0.0	47.0	72.1	273.0	261.0	1,235.8	2,348.0	5,650.4	13,836.2
Sandoval	673.1	650.8	27.1	19.8	84.5	161.0	241.4	371.3	662.7	1,315.8	3,926.2	13,934.9
San Juan	786.9	651.5	11.6	4.9	165.1	299.3	331.0	466.4	865.0	1,870.3	4,437.2	12,796.4
San Miguel	841.4	634.3	0.0	29.4	91.3	221.7	284.4	810.6	1,059.3	1,660.9	4,752.6	13,855.4
Santa Fe	657.6	670.0	30.4	11.9	126.3	179.4	241.9	453.8	634.8	1,092.5	3,543.4	13,940.0
Sierra	915.0	1,803.2	0.0	100.6	0.0	0.0	201.1	624.9	1,569.5	2,312.8	5,349.4	14,762.7
Socorro	868.3	1,272.0	0.0	0.0	138.4	95.5	261.1	487.8	900.0	2,674.3	4,703.6	15,754.1
Taos	726.9	1,115.3	69.7	0.0	117.6	448.0	409.7	557.2	700.5	1,265.0	3,895.9	10,927.8
Torrance	833.1	0.0	0.0	0.0	51.0	240.8	157.8	555.2	1,078.1	2,292.4	4,888.3	14,714.0
Union	660.7	0.0	0.0	208.9	0.0	149.9	0.0	576.6	546.1	2,072.9	3,757.6	11,552.4
Valencia	745.0	665.5	23.8	8.9	106.9	154.8	219.6	445.8	730.0	1,648.8	5,230.7	12,537.1
Health Region												
Northwest	839.6	609.1	26.5	16.5	183.2	307.6	406.6	550.7	986.0	1,751.4	4,751.5	13,222.1
Northeast	723.4	629.3	44.9	20.0	155.2	232.0	325.8	538.8	704.1	1,334.5	3,802.0	13,582.3
Metro	725.1	612.3	30.6	9.8	78.0	135.0	206.4	423.6	824.8	1,616.9	4,502.8	13,737.8
Southeast	830.8	855.7	32.9	7.1	69.9	167.1	228.2	515.2	966.6	2,161.2	4,840.1	15,035.0
Southwest	728.6	540.1	38.6	15.9	66.5	140.8	225.4	473.7	953.4	1,707.3	4,143.1	13,293.4

Age-specific death rates are the numbers of deaths per 100,000 in the specified age group.

Age-adjusted death rates (all ages) are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates and health regions. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

See numbers in Table M-2(a).

Mortality Section

**Table M-3(b) Death Rates by Age, Sex, County, and Health Region
New Mexico, 2012, and United States, 2010**

Females

	Less											
	All ages	than 1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
United States	634.9	564.0	23.3	11.1	36.4	64.0	128.9	311.4	643.5	1,527.5	4,137.7	13,219.2
New Mexico	612.8	520.0	26.3	7.1	49.7	100.4	172.4	340.4	602.1	1,321.1	3,700.9	13,090.2
County												
Bernalillo	591.2	336.3	28.0	0.0	37.5	72.3	164.3	277.4	615.0	1,274.7	3,651.3	13,175.8
Catron	1,375.3	0.0	0.0	0.0	0.0	1,615.5	3,736.9	351.2	469.2	1,246.0	3,611.6	13,932.0
Chaves	719.4	390.2	48.8	0.0	42.8	75.0	261.1	401.5	708.9	1,583.8	4,706.9	14,254.8
Cibola	733.1	533.1	133.3	0.0	53.1	118.6	511.4	363.3	419.4	1,033.2	4,080.3	17,730.8
Colfax	591.9	0.0	0.0	0.0	0.0	0.0	155.2	97.9	270.8	2,152.5	4,323.7	12,497.8
Curry	780.7	694.5	0.0	0.0	0.0	56.6	134.6	422.5	1,099.6	2,220.6	4,361.0	15,888.6
De Baca	1,004.6	0.0	0.0	0.0	0.0	0.0	1,110.2	1,386.0	1,281.9	1,641.1	3,596.3	16,490.1
Dona Ana	572.7	395.6	16.5	26.6	22.2	89.3	120.9	347.6	607.5	1,100.8	3,597.1	12,494.6
Eddy	714.8	272.1	0.0	0.0	29.0	180.2	188.7	386.8	867.8	2,111.1	3,884.1	13,567.9
Grant	624.1	604.2	0.0	0.0	186.2	69.0	138.2	250.3	631.2	1,409.4	3,490.0	14,169.6
Guadalupe	593.8	0.0	0.0	0.0	0.0	0.0	969.0	611.9	402.2	913.0	4,859.9	2,609.3
Harding	172.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	11,131.1
Hidalgo	617.9	0.0	0.0	0.0	308.5	0.0	0.0	569.9	0.0	450.2	4,309.1	17,753.1
Lea	789.0	1,001.4	83.4	39.1	21.2	90.4	128.2	397.5	636.2	2,199.1	4,636.8	17,188.0
Lincoln	555.7	4,404.4	0.0	0.0	107.4	115.6	191.9	775.3	293.8	528.2	3,063.8	8,421.1
Los Alamos	529.9	0.0	0.0	0.0	0.0	120.3	0.0	179.0	289.1	793.2	2,982.3	17,932.3
Luna	611.6	0.0	0.0	0.0	120.2	157.3	75.0	192.6	928.3	1,749.4	3,152.0	12,737.7
McKinley	616.4	819.6	0.0	16.5	162.4	129.6	203.2	464.3	634.0	1,147.8	3,768.4	10,760.4
Mora	934.9	0.0	0.0	0.0	859.4	0.0	357.9	581.8	709.5	1,540.6	3,665.7	22,638.1
Otero	600.7	646.4	0.0	0.0	0.0	24.9	170.8	343.2	834.9	1,200.5	3,933.4	11,985.6
Quay	858.4	4,262.8	0.0	0.0	0.0	0.0	207.4	1,368.0	544.2	1,097.7	3,776.1	18,828.5
Rio Arriba	746.6	727.7	91.0	0.0	118.6	352.5	494.1	574.4	561.0	1,588.6	3,319.0	13,337.7
Roosevelt	683.8	660.6	0.0	0.0	48.0	75.2	189.3	171.5	872.1	2,709.2	4,248.7	10,214.1
Sandoval	554.0	336.3	28.0	20.2	12.3	109.2	179.8	234.3	481.7	1,064.6	3,569.0	12,582.9
San Juan	651.3	655.4	0.0	0.0	87.1	138.6	251.1	401.8	652.1	1,608.3	4,056.3	11,052.1
San Miguel	647.0	1,245.0	0.0	0.0	47.6	197.4	186.3	592.4	632.0	1,172.2	3,563.9	12,502.6
Santa Fe	525.0	243.8	30.5	12.0	39.2	110.9	88.6	357.7	433.2	914.2	2,709.4	13,892.4
Sierra	704.8	0.0	0.0	0.0	0.0	0.0	197.7	499.8	968.8	1,978.2	4,110.0	13,272.0
Socorro	740.9	845.2	0.0	0.0	78.8	102.3	0.0	0.0	805.0	3,365.9	4,124.1	14,635.5
Taos	496.5	1,133.4	141.7	0.0	62.2	177.5	104.1	229.6	461.9	1,138.5	2,788.1	9,797.3
Torrance	652.4	0.0	0.0	0.0	0.0	260.0	115.7	311.5	707.0	2,116.8	4,377.5	10,226.7
Union	606.6	0.0	0.0	0.0	0.0	0.0	0.0	734.3	386.8	2,224.8	3,868.2	9,895.8
Valencia	607.8	1,001.3	50.1	0.0	121.2	158.4	84.0	386.2	509.8	1,005.3	4,482.6	11,304.5
Health Region												
Northwest	649.8	696.8	13.4	5.6	110.3	133.5	265.5	417.5	616.9	1,383.0	3,971.0	11,768.4
Northeast	575.0	480.0	44.9	5.8	62.4	152.5	165.5	379.6	456.5	1,129.6	3,055.7	13,295.5
Metro	588.4	386.8	29.4	3.3	40.3	85.7	159.3	280.8	586.0	1,233.6	3,718.0	12,935.6
Southeast	720.6	859.0	33.9	9.6	30.1	94.2	187.3	461.3	741.4	1,796.7	4,241.2	14,338.8
Southwest	604.2	433.0	9.8	16.2	39.7	86.8	141.1	319.2	686.4	1,330.4	3,663.5	12,896.4

Age-specific death rates are the numbers of deaths per 100,000 in the specified age group.

Age-adjusted death rates (all ages) are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates and health regions. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Mortality Section

**Table M-3(c) Death Rates by Age, Sex, County, and Health Region
New Mexico, 2012, and United States, 2010**

Males

	Less than											
	All ages	1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
United States	887.1	680.2	29.6	14.6	97.6	141.5	212.5	509.9	1,075.5	2,275.1	5,693.7	15,414.3
New Mexico	906.8	753.0	40.3	17.9	140.2	240.2	330.4	616.6	1,136.7	2,059.2	5,232.4	15,021.4
County												
Bernalillo	900.6	869.8	37.1	15.6	110.7	180.8	233.3	587.8	1,141.1	2,115.7	5,704.5	14,966.6
Catron	805.5	0.0	0.0	0.0	0.0	0.0	749.0	766.3	1,769.2	1,747.4	4,915.5	5,818.6
Chaves	921.3	735.2	0.0	19.5	97.5	198.0	222.1	662.0	1,162.7	2,504.6	5,551.2	14,621.8
Cibola	1,144.5	534.4	0.0	158.7	109.2	249.1	305.4	944.4	1,457.3	2,097.0	8,550.6	15,413.1
Colfax	968.6	1,366.2	341.6	128.8	259.5	0.0	133.8	771.0	1,056.2	2,067.0	5,322.5	18,250.2
Curry	883.8	904.7	0.0	0.0	74.0	229.9	202.9	492.1	964.6	2,746.2	5,695.0	13,516.9
De Baca	1,309.3	0.0	0.0	0.0	0.0	0.0	0.0	664.9	1,729.1	5,401.6	5,861.1	28,968.2
Dona Ana	772.3	640.7	80.1	6.5	76.0	189.4	303.7	533.8	1,053.7	1,975.4	3,835.8	13,290.4
Eddy	1,027.3	245.0	122.5	0.0	215.2	277.2	425.3	621.5	1,062.3	2,723.7	5,904.1	16,729.7
Grant	924.9	544.5	136.1	0.0	234.7	143.8	300.6	986.1	1,005.2	2,391.2	4,394.6	15,045.2
Guadalupe	644.6	0.0	0.0	0.0	321.1	0.0	0.0	240.3	1,475.9	3,043.9	2,312.8	8,655.5
Harding	486.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1,220.5	0.0	4,296.1	12,069.1
Hidalgo	776.3	0.0	0.0	0.0	0.0	416.4	0.0	880.7	606.0	1,714.1	8,649.3	3,043.4
Lea	1,071.8	981.0	0.0	0.0	81.1	228.8	236.5	562.7	1,302.5	2,890.9	5,851.7	21,581.4
Lincoln	755.2	979.2	244.8	0.0	0.0	439.4	307.8	398.6	1,276.3	1,948.4	3,281.8	11,449.6
Los Alamos	536.8	0.0	0.0	76.8	117.0	109.5	173.9	120.9	338.3	1,056.5	2,885.4	14,270.3
Luna	1,004.7	1,084.6	0.0	0.0	59.7	438.9	152.8	771.4	1,497.4	2,330.9	5,500.1	16,882.8
McKinley	1,235.3	311.3	77.8	16.4	317.7	617.5	897.7	874.6	1,950.4	2,289.4	6,001.3	15,618.6
Mora	954.3	0.0	0.0	0.0	372.1	462.3	745.5	1,072.6	1,184.0	1,719.2	2,696.8	15,244.8
Otero	929.0	0.0	50.8	45.8	98.9	178.7	379.2	616.6	1,612.4	2,196.1	5,716.8	12,555.6
Quay	1,034.4	3,710.6	0.0	0.0	407.2	485.3	0.0	809.8	1,493.5	2,229.2	6,481.6	11,835.2
Rio Arriba	1,151.3	382.3	95.6	36.9	523.5	509.6	1,058.1	1,172.9	1,262.2	2,057.4	5,109.4	12,146.3
Roosevelt	1,049.5	1,148.4	0.0	0.0	46.0	69.2	350.3	353.1	1,614.5	1,965.0	7,358.7	20,160.8
Sandoval	819.6	945.6	26.3	19.4	154.6	215.0	305.8	516.2	866.8	1,593.4	4,398.5	16,394.9
San Juan	956.3	647.7	23.1	9.6	239.8	453.3	411.3	531.4	1,085.8	2,172.4	4,924.0	16,279.1
San Miguel	1,069.8	0.0	0.0	57.3	131.6	244.4	385.9	1,023.8	1,504.2	2,153.3	6,228.9	16,539.8
Santa Fe	803.0	1,095.6	30.4	11.9	207.4	244.2	393.0	558.0	864.2	1,297.9	4,564.2	14,035.0
Sierra	1,124.3	3,532.5	0.0	191.3	0.0	0.0	204.6	750.0	2,192.5	2,630.6	6,519.8	16,500.0
Socorro	1,014.0	1,701.8	0.0	0.0	185.0	89.6	524.9	1,002.6	987.5	2,004.9	5,299.2	17,794.1
Taos	990.9	1,097.8	0.0	0.0	167.1	724.0	705.6	912.2	956.8	1,402.0	5,323.7	13,020.6
Torrance	1,060.0	0.0	0.0	0.0	91.9	224.3	192.8	808.3	1,434.8	2,455.2	5,414.9	22,337.2
Union	718.4	0.0	0.0	382.8	0.0	208.9	0.0	474.7	687.7	1,909.9	3,603.0	14,356.2
Valencia	900.6	362.0	0.0	17.4	93.6	151.4	354.1	504.8	953.1	2,347.3	6,039.2	14,506.8
Health Region												
Northwest	1,067.3	523.5	39.3	27.2	254.0	474.8	546.0	688.9	1,382.5	2,194.8	5,783.9	15,974.3
Northeast	883.8	778.4	44.8	33.7	240.4	305.3	481.2	703.4	971.1	1,558.2	4,711.1	14,116.9
Metro	890.8	824.9	31.7	16.1	114.8	183.6	253.7	573.4	1,086.9	2,058.4	5,545.9	15,249.8
Southeast	957.5	852.6	32.0	4.6	106.6	233.8	268.5	568.8	1,200.4	2,560.1	5,585.2	16,259.3
Southwest	864.6	643.0	66.2	15.6	91.7	192.4	314.9	642.0	1,238.6	2,117.6	4,671.5	13,915.6

Age-specific death rates are the numbers of deaths per 100,000 in the specified age group.

Age-adjusted death rates (all ages) are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates and health regions. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution. See numbers in Table M-2(c).

Mortality Section

**Table M-4(a) Number of Deaths Ranked by Leading Causes of Death by County and Health Region
New Mexico, 2012, and United States, 2010**

	1	2	3	4	5	6	7	8	9	10	
	Neoplasm, malignant (ICD10: C00-C97)	Circulatory, heart disease (ICD10: I00-I09, I11, I13, I20-I51)	Injury, unintentional injuries (ICD10: V01-X59, Y85-Y86)	Respiratory, chronic lower respiratory diseases (ICD10: J40-J47)	Circulatory, cardiovascular diseases (ICD10: I60-I69)	Diabetes mellitus (ICD10: E10-E14)	Chronic liver disease and cirrhosis (ICD10: K70, K73-K74)	Injury, intentional self-harm (suicide) (ICD10: X84, X87, X90, X93)	Alzheimer's disease (ICD10: G30)	Respiratory, influenza and pneumonia (ICD10: J09-J18)	
United States	2,468,435	574,743	597,689	120,859	138,080	129,476	69,071	31,903	38,364	83,494	50,097
New Mexico	16,640	3,422	3,306	1,331	961	665	648	436	435	369	283
County											
Bernalillo	5,097	1,059	1,008	390	269	213	197	113	132	132	78
Catron	53	17	11	5	1	1	1	0	8	2	0
Chaves	612	114	146	41	58	18	23	8	13	18	11
Cibola	249	35	34	21	15	9	13	20	6	2	6
Colfax	154	30	36	12	9	7	7	4	4	4	0
Curry	384	71	86	22	32	14	27	5	6	15	5
De Baca	38	9	10	2	0	3	3	1	1	0	1
Dona Ana	1,488	328	247	91	79	69	69	24	35	32	19
Eddy	534	104	128	35	51	22	26	10	14	7	5
Grant	343	70	81	20	14	17	9	9	10	7	7
Guadalupe	33	6	5	2	1	1	3	2	2	0	1
Harding	7	4	2	0	0	0	0	0	0	0	0
Hidalgo	44	6	8	2	4	3	3	2	0	1	1
Lea	533	95	120	47	46	19	15	9	8	5	8
Lincoln	181	40	40	14	13	8	7	7	8	5	2
Los Alamos	127	22	32	7	7	5	3	2	0	4	4
Luna	273	69	70	12	21	12	6	5	4	7	3
McKinley	553	88	72	73	12	21	40	37	4	6	14
Mora	56	8	14	10	0	5	1	2	31	1	1
Otero	547	126	110	43	36	26	13	11	14	10	4
Quay	116	18	25	7	10	5	1	3	4	4	3
Rio Arriba	400	77	58	63	10	13	18	24	10	4	4
Roosevelt	171	29	44	15	18	3	3	1	1	3	4
Sandoval	907	184	177	72	46	36	30	27	25	30	22
San Juan	915	178	151	102	49	30	38	34	27	25	26
San Miguel	278	66	69	26	13	3	14	6	4	2	3
Santa Fe	1,102	233	228	96	46	36	30	32	37	21	15
Sierra	227	49	64	11	16	12	5	3	1	0	7
Socorro	165	37	34	9	17	9	5	7	3	0	4
Taos	290	72	47	32	17	11	10	8	12	5	4
Torrance	151	33	33	11	8	8	2	2	1	5	1
Union	41	10	8	1	3	0	1	1	1	1	1
Valencia	568	137	108	37	40	25	20	14	11	12	12
Unknown County	3	0	0	0	0	1	0	0	1	0	0
Health Region											
Northwest	1,717	301	257	196	76	60	91	33	64	91	46
Northeast	2,488	528	499	249	106	81	87	41	71	82	40
Metro	6,723	1,411	1,326	510	363	282	249	179	169	156	113
Southeast	2,569	480	599	183	228	92	103	57	55	46	39
Southwest	3,140	702	625	193	188	149	118	59	75	61	45
Unknown Region	3	0	0	0	0	1	0	0	1	0	0

See Technical Appendix for information on health regions. Table is ranked by number of deaths for New Mexico.

Mortality Section

**Table M-4(b) Death Rates for Ranked Leading Causes by County and Health Region
New Mexico, 2012, and United States, 2010**

	1	2	3	4	5	6	7	8	9	10	
	Neoplasm, malignant (ICD10: C00- C97)	Circulatory, Heart disease (ICD10: I00-I09, I11, I13, I20-I51)	Injury, Unintentional Injuries (ICD10: V01-X59, Y85-Y86)	Respiratory, Chronic lower respiratory diseases (ICD10: J40-J47)	Circulatory, Cerebrovascula r diseases (ICD10: I60-I69)	Diabetes mellitus (ICD10: E10- E14)	Betetes E10- E14	Chronic liver disease and cirrhosis (ICD10: K70, K73-K74)	Injury, self-harm (suicide) (ICD10: X60- X84, Y87.0, *U09)	Alzheimer's disease (ICD10: G30)	Respiratory, Influenza and pneumonia (ICD10: J09-J18)
United States	799.5	1862.2	193.6	39.1	44.7	41.9	22.4	10.3	12.4	27.0	16.2
New Mexico	749.4	147.3	148.5	63.9	42.8	30.4	28.3	18.9	20.9	17.4	12.8
County											
Bernalillo	730.5	148.5	144.5	56.4	39.3	30.9	28.3	15.5	19.0	19.3	11.4
Caron	1071.6	321.8	162.5	251.1	14.9	14.9	18.3	0.0	144.7	27.5	0.0
Chaves	815.0	150.7	189.2	59.6	76.1	22.5	29.5	11.4	19.8	23.7	13.8
Cibola	929.9	124.0	122.5	76.6	56.2	35.2	49.9	70.2	22.0	8.4	25.7
Colfax	769.2	134.3	170.1	74.9	42.4	34.3	31.9	23.8	31.8	19.4	0.0
Curry	833.6	150.5	189.3	47.9	70.1	31.6	58.7	14.4	12.0	34.1	11.2
De Baca	1124.7	328.7	263.0	54.1	0.0	83.7	75.4	46.8	26.5	0.0	26.5
Dona Ana	666.2	137.9	107.6	44.5	34.5	32.3	30.5	10.9	17.1	15.3	8.7
Eddy	857.7	163.6	199.2	66.2	78.9	35.4	39.6	16.8	26.4	11.3	6.9
Grant	768.3	141.2	180.2	56.0	28.3	38.0	17.7	19.7	39.6	14.6	14.6
Guadalupe	599.0	98.7	86.6	43.2	17.7	2.12	48.4	30.7	59.0	0.0	17.7
Harding	345.2	210.5	89.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hidalgo	733.9	100.4	126.4	37.6	63.1	48.8	60.2	34.0	0.0	16.4	16.4
Lea	919.9	161.0	214.6	76.7	83.3	34.5	24.4	13.3	14.0	9.6	13.4
Lincoln	649.6	127.5	137.0	58.5	39.1	28.8	15.2	20.7	34.3	19.1	8.4
Los Alamos	539.7	87.8	129.5	41.9	29.8	20.9	11.8	11.0	0.0	19.0	15.3
Luna	801.9	205.4	190.3	46.3	55.9	34.8	17.9	12.8	17.2	20.7	8.9
McKinley	892.7	140.2	121.3	105.2	20.2	41.6	64.2	54.0	46.2	12.1	24.0
Mora	939.7	114.4	220.9	215.5	0.0	86.8	14.4	22.3	29.7	0.0	18.8
Otero	760.6	162.0	153.7	64.6	48.5	37.0	18.7	13.9	20.6	14.2	5.3
Rio Arriba	947.3	172.9	188.2	77.4	23.1	40.2	6.1	18.5	51.9	32.7	22.6
Quay	957.0	128.8	131.6	131.6	89.7	15.2	14.2	5.5	7.4	10.3	19.9
Roosevelt	839.4	137.5	215.8	78.6	89.7	28.2	19.8	18.3	7.4	13.1	20.2
Sandoval	673.1	126.4	132.0	55.5	34.0	19.8	24.5	19.8	18.3	24.5	17.5
San Juan	786.9	151.4	134.5	82.0	43.6	26.9	31.4	26.5	22.2	23.8	22.4
San Miguel	841.4	183.0	204.5	88.0	35.8	10.4	46.2	19.8	12.5	6.9	9.4
Santa Fe	915.0	124.3	135.5	63.7	28.6	20.8	17.8	17.0	24.8	13.2	9.2
Sierra	868.3	203.4	249.5	57.3	57.0	40.8	17.0	11.0	3.6	0.0	25.0
Socorro	726.9	157.3	122.3	92.0	40.4	49.9	62.0	33.6	14.5	0.0	22.5
Taos	833.1	154.4	186.3	65.2	41.4	46.9	13.8	6.7	3.4	31.3	6.6
Torrance	660.7	162.6	118.0	15.2	45.8	0.0	16.8	19.4	20.3	13.8	15.2
Union	745.0	169.6	144.8	51.1	51.2	34.3	25.3	17.2	15.4	19.4	17.2
Health Region											
Northwest	839.6	144.7	128.7	88.8	38.3	32.3	43.7	40.8	29.7	18.5	23.2
Northeast	723.4	139.3	141.8	84.0	30.1	23.2	24.0	22.6	24.3	12.4	12.0
Metro	725.1	147.1	143.8	55.7	39.8	31.0	26.6	16.1	18.3	20.2	12.6
Southwest	830.8	151.2	192.8	64.3	72.3	29.9	32.2	14.6	19.4	18.8	12.1
Southwest	728.6	153.5	141.8	51.4	41.7	34.8	27.1	14.0	20.2	14.0	10.4

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.
See Technical Appendix for information on rates and health regions. Table is ranked by number of deaths for New Mexico. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.
See numbers in Table M-4(a).

Mortality Section

**Table M-5 Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death
New Mexico, 2012, All Ages**

	Male	Female	Both Sexes
All Races			
All Causes	906.8	612.8	749.4
Neoplasm, malignant (C00-C97)	180.6	121.9	147.3
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	180.0	121.4	148.5
Injury, Unintentional injuries (V01-X59, Y85-Y86)	86.7	42.4	63.9
Respiratory, Chronic lower respiratory diseases (J40-J47)	50.2	37.4	42.8
Circulatory, Cerebrovascular diseases (I60-I69)	30.3	29.8	30.4
American Indian and Alaska Native			
All Causes	1,197.7	678.8	908.4
Injury, Unintentional injuries (V01-X59, Y85-Y86)	171.9	57.2	110.7
Neoplasm, malignant (C00-C97)	166.9	113.3	135.8
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	162.0	69.0	107.9
Diabetes mellitus (E10-E14)	102.1	57.4	77.1
Chronic liver disease and cirrhosis (ICD10: K70, K73-K74)	72.6	43.4	57.0
Asian or Pacific Islander			
All Causes	738.6	395.4	527.6
Neoplasm, malignant (C00-C97)	183.3	62.7	109.3
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	132.1	90.6	108.7
Circulatory, Cerebrovascular diseases (I60-I69)	60.1	38.4	45.6
Diabetes mellitus (E10-E14)	71.6	8.3	32.0
Injury, Unintentional injuries (V01-X59, Y85-Y86)	46.4	13.2	27.0
Injury, Intentional self-harm (suicide) (ICD10: X60-X84, Y87.0, *U03)	7.3	18.7	13.3
Black or African American			
All Causes	1048.6	596.3	817.1
Neoplasm, malignant (C00-C97)	299.9	123.6	200.8
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	213.3	138.2	176.4
Injury, Unintentional injuries (V01-X59, Y85-Y86)	72.7	31.5	53.6
Diabetes mellitus (E10-E14)	65.2	37.0	53.9
Respiratory, Chronic lower respiratory diseases (J40-J47)	25.2	44.1	34.3
Hispanic			
All Causes	903	602.9	742.8
Neoplasm, malignant (C00-C97)	181.1	117.2	144.9
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	164.8	120.5	141.7
Injury, Unintentional injuries (V01-X59, Y85-Y86)	85.7	43.0	64.1
Diabetes mellitus (E10-E14)	44.0	34.6	39.3
Respiratory, Chronic lower respiratory diseases (J40-J47)	39.2	25.2	31.2
White			
All Causes	857.8	609.1	724.8
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	187.8	127.1	155.0
Neoplasm, malignant (C00-C97)	178.1	128.4	149.7
Respiratory, Chronic lower respiratory diseases (J40-J47)	58.2	47.2	51.8
Injury, Unintentional injuries (V01-X59, Y85-Y86)	71.7	38.8	54.6
Circulatory, Cerebrovascular diseases (I60-I69)	27.2	30.6	29.6

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

See Technical Appendix for information on rates and race/ethnicity. Causes are the five leading causes of death ranked by number of deaths for both sexes combined within each racial/ethnic category. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Mortality Section

**Table M-6 Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death
New Mexico, 2012, Ages 1 to 4 Years**

	Male	Female	Both Sexes
All Races			
All Causes	40.3	26.3	33.5
Injury, Unintentional injuries (V01-X59, Y85-Y86)	20.2	8.8	14.6
Neoplasm, malignant (ICD10: C00-C97)	5.0	5.3	5.2
Congenital malformations, deformations and chromosomal abnormalities (ICD10: Q00-Q99)	3.4	1.8	2.6
Respiratory, Chronic lower respiratory diseases (ICD10: J40-J47)	3.4	1.8	2.6
American Indian and Alaska Native			
All Causes	15.9	32.1	24.0
Injury, Unintentional injuries (V01-X59, Y85-Y86)	15.9	32.1	24.0
Asian or Pacific Islander			
All Causes	0.0	0.0	0.0
Black or African American			
All Causes	72.5	162.3	114.9
Hispanic			
All Causes	39.6	29.3	34.5
Injury, Unintentional injuries (V01-X59, Y85-Y86)	17.0	8.8	13.0
Neoplasm, malignant (ICD10: C00-C97)	5.6	5.9	5.8
Respiratory, Chronic lower respiratory diseases (ICD10: J40-J47)	5.6	2.9	4.3
White			
All Causes	50.9	6.8	29.6
Injury, Unintentional injuries (V01-X59, Y85-Y86)	25.4	0.0	13.2

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

See Technical Appendix for information on rates and race/ethnicity. Causes are the five leading causes of death ranked by number of deaths for both sexes combined within each racial/ethnic category. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Mortality Section

**Table M-7 Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death
New Mexico, 2012, Ages 5 to 14 Years**

	Male	Female	Both Sexes
All Races			
All Causes	17.9	6.4	12.2
Injury, Unintentional injuries (V01-X59, Y85-Y86)	6.2	0.7	3.5
Neoplasm, malignant (C00-C97)	4.8	2.1	3.5
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1.4	0.7	1.0
American Indian and Alaska Native			
All Causes	12.7	6.4	9.5
Asian or Pacific Islander			
All Causes	53.4	52.7	53.1
Black or African American			
All Causes	0.0	0.0	0.0
Hispanic			
All Causes	19.6	7.6	13.6
Injury, Unintentional injuries (V01-X59, Y85-Y86)	7.4	0.0	3.7
Neoplasm, malignant (C00-C97)	4.9	2.5	3.7
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2.4	1.3	1.9
White			
All Causes	16.4	2.5	9.6
Neoplasm, malignant (C00-C97)	4.9	2.5	3.6

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

See Technical Appendix for information on rates and race/ethnicity. Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Mortality Section

**Table M-8 Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death
New Mexico, 2012, Ages 15 to 24 Years**

	Male	Female	Both Sexes
All Races			
All Causes	139.5	49.7	95.7
Injury, Unintentional injuries (V01-X59, Y85-Y86)	60.1	21.0	41.0
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	32.7	9.8	21.5
Injury, Assault (homicide) (X85-Y09, Y87.1, U01-U02)	20.0	3.5	12.0
Neoplasm, malignant (C00-C97)	6.0	2.8	4.4
Circulatory, Heart disease (ICD10: I00-I09, I11, I13, I20-I51)	2.0	0.0	1.0
American Indian and Alaska Native			
All Causes	285.9	104.2	195.4
Injury, Unintentional injuries (V01-X59, Y85-Y86)	127.8	49.0	88.5
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	54.8	24.5	39.7
Injury, Assault (homicide) (X85-Y09, Y87.1, U01-U02)	24.3	6.1	15.3
Neoplasm, malignant (C00-C97)	18.2	0.0	9.2
Asian or Pacific Islander			
All Causes	0.0	41.1	20.4
Black or African American			
All Causes	71.2	0.0	41.8
Injury, Assault (homicide) (X85-Y09, Y87.1, U01-U02)	71.2	0.0	41.8
Hispanic			
All Causes	145.7	56.9	101.8
Injury, Unintentional injuries (V01-X59, Y85-Y86)	55.6	25.8	40.9
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	34.4	6.8	20.8
Injury, Assault (homicide) (X85-Y09, Y87.1, U01-U02)	27.8	5.4	16.8
Neoplasm, malignant (C00-C97)	6.6	4.1	5.4
White			
All Causes	95.7	23.2	60.9
Injury, Unintentional injuries (V01-X59, Y85-Y86)	52.7	6.3	30.4
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	27.3	8.4	18.3

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

See Technical Appendix for information on rates and race/ethnicity. Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Due to small numbers, causes are only shown if there are 3 or more deaths.

Mortality Section

**Table M-9 Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death
New Mexico, 2012, Ages 25 to 44 Years**

All Races	Male	Female	Both Sexes
All Causes	281.1	135.0	208.7
Injury, Unintentional injuries (V01-X59, Y85-Y86)	99.0	44.2	71.9
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	43.4	11.2	27.5
Neoplasm, malignant (C00-C97)	15.2	14.4	14.8
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	18.3	9.3	13.8
Chronic liver disease and cirrhosis (K70, K73-K74)	14.9	6.6	10.8
American Indian and Alaska Native			
All Causes	652.8	313.7	478.8
Injury, Unintentional injuries (V01-X59, Y85-Y86)	217.6	82.6	148.3
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	108.8	16.5	61.4
Chronic liver disease and cirrhosis (K70, K73-K74)	69.6	37.2	53.0
Injury, Assault (homicide) (X85-Y09, Y87.1, U01-U02)	30.5	8.3	19.1
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	21.8	16.5	19.1
Asian or Pacific Islander			
All Causes	129.4	35.6	78.0
Black or African American			
All Causes	254.5	64.1	176.0
Injury, Unintentional injuries (V01-X59, Y85-Y86)	59.9	42.7	52.8
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	44.9	0.0	26.4
Hispanic			
All Causes	287.0	129.6	208.7
Injury, Unintentional injuries (V01-X59, Y85-Y86)	109.8	44.6	77.4
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	28.9	8.1	18.5
Neoplasm, malignant (C00-C97)	16.0	17.0	16.5
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	21.6	6.5	14.1
Injury, Assault (homicide) (X85-Y09, Y87.1, U01-U02)	20.8	4.9	12.9
White			
All Causes	200.1	107.2	154.5
Injury, Unintentional injuries (V01-X59, Y85-Y86)	63.8	36.1	50.2
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	48.3	13.0	31.0
Neoplasm, malignant (C00-C97)	13.5	12.0	12.8
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	10.6	12.0	11.3
Chronic liver disease and cirrhosis (K70, K73-K74)	7.7	2.0	4.9

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

See Technical Appendix for information on rates and race/ethnicity. Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Mortality Section

**Table M-10 Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death
New Mexico, 2012, Ages 45 to 64 Years**

	Male	Female	Both Sexes
All Races			
All Causes	856.4	465.9	655.4
Neoplasm, malignant (C00-C97)	191.7	150.5	170.5
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	144.4	56.1	99.0
Injury, Unintentional injuries (V01-X59, Y85-Y86)	103.8	43.2	72.6
Chronic liver disease and cirrhosis (K70, K73-K74)	70.4	23.9	46.4
Diabetes mellitus (E10-E14)	39.1	20.7	29.6
American Indian and Alaska Native			
All Causes	1266.8	512.1	863.2
Neoplasm, malignant (ICD10: C00-C97)	184.0	123.4	151.6
Injury, Unintentional injuries (ICD10: V01-X59, Y85-Y86)	262.8	50.3	149.2
Chronic liver disease and cirrhosis (ICD10: K70, K73-K74)	168.2	86.9	124.7
Circulatory, Heart disease (ICD10: I00-I09, I11, I13, I20-I51)	126.2	22.9	70.9
Diabetes mellitus (ICD10: E10-E14)	94.6	27.4	58.7
Asian or Pacific Islander			
All Causes	526.1	135.3	300.0
Neoplasm, malignant (C00-C97)	278.5	45.1	143.5
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	154.8	0.0	65.2
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	31.0	22.6	26.1
Black or African American			
All Causes	1011.1	573.3	811.8
Circulatory, Heart disease (ICD10: I00-I09, I11, I13, I20-I51)	159.6	127.4	145.0
Neoplasm, malignant (ICD10: C00-C97)	159.6	127.4	145.0
Injury, Unintentional injuries (ICD10: V01-X59, Y85-Y86)	124.2	21.2	77.3
Diabetes mellitus (ICD10: E10-E14)	88.7	0.0	48.3
Injury, Assault (homicide) (ICD10: X85-Y09, Y87.1, U01-U02)	71.0	0.0	38.7
Circulatory, Cerebrovascular diseases (ICD10: I60-I69)	53.2	21.2	38.7
Chronic liver disease and cirrhosis (K70, K73-K74)	53.2	0.0	29.0
Respiratory, Chronic lower respiratory diseases (ICD10: J40-J47)	0.0	63.7	29.0
Human immunodeficiency virus (HIV) disease (ICD10: B20-B24)	53.2	0.0	29.0
Hispanic			
All Causes	821.4	396.2	603.9
Neoplasm, malignant (C00-C97)	169.9	116.6	142.6
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	112.4	53.2	82.1
Injury, Unintentional injuries (V01-X59, Y85-Y86)	101.8	39.7	70.0
Chronic liver disease and cirrhosis (K70, K73-K74)	86.7	24.5	54.9
Diabetes mellitus (E10-E14)	46.0	24.5	35.0
White			
All Causes	827.9	526.3	672.6
Neoplasm, malignant (C00-C97)	210.5	188.0	198.9
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	171.4	63.4	115.8
Injury, Unintentional injuries (V01-X59, Y85-Y86)	79.0	47.2	62.6
Injury, Intentional self-harm (suicide) (X60-X84, Y87.0, U03)	58.7	17.7	37.6
Respiratory, Chronic lower respiratory diseases (J40-J47)	36.0	33.9	34.9

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

See Technical Appendix for information on rates and race/ethnicity. Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Mortality Section

**Table M-11 Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death
New Mexico, 2012, Ages 65 to 84 Years**

	Male	Female	Both Sexes
All Races			
All Causes	3,118.9	2,178.8	2,613.0
Neoplasm, malignant (C00-C97)	848.2	575.1	701.2
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	695.0	417.9	545.9
Respiratory, Chronic lower respiratory diseases (J40-J47)	243.9	183.6	211.4
Diabetes mellitus (E10-E14)	148.2	100.7	122.6
Circulatory, Cerebrovascular diseases (I60-I69)	116.5	112.2	114.2
American Indian and Alaska Native			
All Causes	4,006.6	2,263.9	2,988.0
Neoplasm, malignant (C00-C97)	787.3	547.3	647.0
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	752.3	248.8	458.0
Diabetes mellitus (E10-E14)	507.4	323.4	399.9
Injury, Unintentional injuries (V01-X59, Y85-Y86)	210.0	74.6	130.9
Nephritis, nephrotic syndrome and nephrosis (ICD10: N00-N07, N17-N19, N25-N27)	192.5	74.6	123.6
Asian or Pacific Islander			
All Causes	1,772.1	1,019.7	1,315.4
Neoplasm, malignant (C00-C97)	689.2	318.7	464.2
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	295.4	191.2	232.1
Nephritis, nephrotic syndrome and nephrosis (ICD10: N00-N07, N17-N19, N25-N27)	98.4	191.2	154.8
Diabetes mellitus (E10-E14)	295.4	63.7	154.8
Black or African American			
All Causes	4,099.4	2,290.1	3,181.4
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	1,235.4	545.3	885.3
Neoplasm, malignant (C00-C97)	1,403.9	327.2	857.6
Diabetes mellitus (E10-E14)	280.8	218.1	249.0
Injury, Unintentional injuries (V01-X59, Y85-Y86)	168.5	109.0	138.3
Respiratory, Chronic lower respiratory diseases (J40-J47)	112.3	163.6	138.3
Circulatory, Cerebrovascular diseases (I60-I69)	56.2	109.0	83.0
Hispanic			
All Causes	3,197.6	2,232.9	2,668.7
Neoplasm, malignant (C00-C97)	876.9	578.4	713.2
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	680.8	421.3	538.5
Diabetes mellitus (E10-E14)	198.7	148.4	171.1
Respiratory, Chronic lower respiratory diseases (J40-J47)	182.8	115.7	146.0
Circulatory, Cerebrovascular diseases (I60-I69)	153.7	122.2	136.4
White			
All Causes	3,004.9	2,160.1	2,558.6
Neoplasm, malignant (C00-C97)	824.3	585.0	697.9
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	687.6	431.5	552.3
Respiratory, Chronic lower respiratory diseases (J40-J47)	289.7	239.3	263.1
Circulatory, Cerebrovascular diseases (I60-I69)	97.5	105.2	101.5
Injury, Unintentional injuries (V01-X59, Y85-Y86)	101.5	72.5	86.2

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

See Technical Appendix for information on rates and race/ethnicity. Causes are the five leading causes of death for both sexes combined within each racial/ethnic category. Due to small numbers, causes are only shown if there are 3 or more deaths.

Mortality Section

**Table M-12 Death Rates by Race/Ethnicity and Sex for Ranked Causes of Death
New Mexico, 2012, Ages 85 Plus Years**

	Male	Female	Both Sexes
All Races			
All Causes	14,995.8	13,090.2	13,770.7
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	3,951.2	3,737.3	3,813.7
Neoplasm, malignant (C00-C97)	2,341.8	1,282.0	1,660.5
Circulatory, Cerebrovascular diseases (I60-I69)	749.4	908.3	851.6
Respiratory, Chronic lower respiratory diseases (J40-J47)	1,021.9	733.3	836.3
Alzheimer's disease (G30)	459.8	846.8	708.6
American Indian and Alaska Native			
All Causes	12,090.2	11,205.4	11,511.8
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	2,216.5	1,814.2	1,953.5
Neoplasm, malignant (C00-C97)	1,612.0	1,067.2	1,255.8
Respiratory, Influenza and pneumonia (J10-J18)	1,209.0	1,173.9	1,186.1
Circulatory, Cerebrovascular diseases (I60-I69)	1,007.5	853.8	907.0
Diabetes mellitus (ICD10: E10-E14)	1,007.5	853.8	907.0
Parkinson's disease (ICD10: G20-G21)	806.0	213.4	418.6
Asian or Pacific Islander			
All Causes	18,375.4	14,412.9	15,898.6
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	2,827.0	4,239.1	3,709.7
Circulatory, Cerebrovascular diseases (I60-I69)	2,827.0	847.8	1,589.9
Black or African American			
All Causes	13,104.2	11,929.6	12,344.5
Neoplasm, malignant (C00-C97)	5,096.1	3,578.9	4,114.8
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	728.0	3,181.2	2,314.6
Respiratory, Chronic lower respiratory diseases (J40-J47)	728.0	795.3	771.5
Diabetes mellitus (ICD10: E10-E14)	728.0	795.3	771.5
Hispanic			
All Causes	14,013.5	12,673.2	13,159.1
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	3,266.8	3,663.1	3,519.4
Neoplasm, malignant (C00-C97)	2,198.2	1,180.5	1,549.4
Circulatory, Cerebrovascular diseases (I60-I69)	732.7	798.6	774.7
Respiratory, Chronic lower respiratory diseases (J40-J47)	854.8	572.9	675.1
Alzheimer's disease (G30)	488.5	763.9	664.0
White			
All Causes	15,598.7	13,396.0	14,179.2
Circulatory, Heart disease (I00-I09, I11, I13, I20-I51)	4,418.1	3,901.5	4,085.2
Neoplasm, malignant (C00-C97)	2,395.8	1,300.5	1,690.0
Respiratory, Chronic lower respiratory diseases (J40-J47)	1,120.6	838.6	938.9
Circulatory, Cerebrovascular diseases (I60-I69)	721.3	973.6	883.9
Alzheimer's disease (G30)	476.6	923.9	764.8

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

See Technical Appendix for information on rates and race/ethnicity. Causes are the five leading causes of death for both sexes combined

within each racial/ethnic category. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Mortality Section

**Table M-13(a) Number of Deaths by Race/Ethnicity and Sex
New Mexico, 2008-2012, and United States, 2008-2010**

	2008	2009	2010	2011	2012
All Races					
Male	8,008	8,094	8,299	8,510	8,967
Female	7,392	7,298	7,567	7,735	7,673
Both Sexes	15,400	15,392	15,866	16,245	16,640
American Indian or Alaska Native					
Male	629	609	627	638	735
Female	557	496	521	507	506
Both Sexes	1,186	1,105	1,148	1,145	1,241
Asian or Pacific Islander					
Male	29	38	48	44	58
Female	43	31	60	57	43
Both Sexes	72	69	108	101	101
Black or African American					
Male	148	136	149	150	174
Female	109	113	132	103	104
Both Sexes	257	249	281	253	278
Hispanic					
Male	2,690	2,777	2,793	2,994	3,153
Female	2,209	2,226	2,234	2,415	2,486
Both Sexes	4,899	5,003	5,027	5,409	5,639
White					
Male	4,476	4,487	4,640	4,639	4,799
Female	4,459	4,416	4,601	4,636	4,523
Both Sexes	8,935	8,903	9,241	9,275	9,322
Other and Unknown Race/Ethnicity					
Male	36	47	42	45	48
Female	15	16	19	17	11
Both Sexes	51	63	61	62	59
United States					
All Races					
Male	1,226,197	1,217,379	1,232,432	N/A	N/A
Female	1,245,787	1,219,786	1,236,003	N/A	N/A
Both Sexes	2,471,984	2,437,163	2,468,435	N/A	N/A

See Technical Appendix for information on race/ethnicity.

Mortality Section

**Table M-13(b) Death Rates by Race/Ethnicity and Sex
New Mexico, 2008-2012, and United States, 2008-2010**

	2008	2009	2010	2011	2012
All Races					
Male	883.0	873.5	870.6	868.5	906.8
Female	641.9	622.9	626.5	622.2	612.8
Both Sexes	756.0	738.5	741.9	735.4	749.4
American Indian or Alaska Native					
Male	1084.3	1047.4	1088.5	1062.1	1197.7
Female	799.0	702.8	734.5	686.2	678.8
Both Sexes	930.8	855.6	891.5	852.4	908.4
Asian or Pacific Islander					
Male	380.9	466.1	551.9	454.7	738.6
Female	419.2	260.7	494.6	467.2	395.4
Both Sexes	403.2	342.9	520.3	467.3	527.6
Black or African American					
Male	999.8	894.3	870.3	870.9	1048.6
Female	690.0	683.9	773.4	590.4	596.3
Both Sexes	845.1	784.9	835.4	726.8	817.1
Hispanic					
Male	855.2	857.4	838.6	873.6	903.0
Female	601.4	582.4	564.1	586.0	602.9
Both Sexes	721.4	709.0	691.8	718.1	742.8
White					
Male	858.1	848.6	858.6	832.7	857.8
Female	643.5	635.1	648.7	637.3	609.1
Both Sexes	744.9	733.6	749.0	726.4	724.8
United States					
All Races					
Male	820.3	807.2	812.0	NA	NA
Female	805.8	782.1	787.4	NA	NA
Both Sexes	812.9	794.5	799.5	NA	NA

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates and race/ethnicity. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

See numbers in Table M-13(a).

Mortality Section

**Table M-14(a) Number of Deaths for Leading Causes of Death
New Mexico, 2003-2012**

Rank in 2003	Rank in 2012	Cause of death	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
2	1	Malignant neoplasms (cancer)	3,045	3,020	3,145	3,136	3,210	3,245	3,141	3,328	3,260	3,422
1	2	Diseases of heart	3,145	3,161	3,376	3,331	3,263	3,174	3,152	3,186	3,275	3,306
3	3	Accidents (unintentional injuries)	1,127	1,138	1,192	1,267	1,277	1,298	1,227	1,208	1,296	1,331
4	4	Chronic lower respiratory diseases	957	759	866	911	897	965	970	1,015	1,022	961
5	5	Cerebrovascular diseases (stroke)	724	609	625	632	649	714	714	798	704	665
6	6	Diabetes mellitus	639	637	623	560	642	562	592	636	594	648
10	7	Chronic liver disease and cirrhosis	307	278	286	322	383	367	360	372	461	436
8	8	Intentional self-harm (suicide)	343	359	341	347	400	406	372	410	419	435
7	9	Alzheimer's disease	383	343	345	358	324	359	369	339	417	369
9	10	Influenza and pneumonia	325	299	351	357	349	341	337	288	350	283
12	11	Nephritis, nephrotic syndrome and nephrosis	200	244	248	266	249	276	270	276	271	282
16	12	Septicemia	109	117	130	174	177	155	197	184	205	212
13	13	Parkinson's disease	166	145	153	173	164	158	161	171	169	173
11	14	Essential primary hypertension and hypertensive renal disease	227	200	169	232	200	125	115	125	127	141
14	15	Assault (homicide)	157	175	151	137	164	144	174	149	146	133
ALL CAUSES			14,489	14,197	14,866	15,231	15,400	15,400	15,392	15,866	16,245	16,640

Table is ranked by number of deaths for New Mexico.

Mortality Section

**Table M-14(b) Death Rates for Leading Causes of Death
New Mexico, 2003-2012**

Rank in 2003	Rank in 2012	Cause of death	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
2	1	Malignant neoplasms (cancer)	163.5	157.9	160.0	156.5	155.9	154.6	146.4	150.2	141.7	147.3
1	2	Diseases of heart	174.4	170.9	178.1	171.9	163.6	155.4	150.7	148.7	147.3	148.5
3	3	Accidents (unintentional injuries)	61.0	60.6	62.6	66.0	65.1	65.7	61.1	59.3	62.5	63.9
4	4	Chronic lower respiratory diseases	53.0	41.0	45.6	47.0	45.4	47.7	46.5	47.1	45.6	42.8
5	5	Cerebrovascular diseases (stroke)	40.7	33.5	33.4	33.0	33.0	35.8	34.7	37.8	32.5	30.4
6	6	Diabetes mellitus	34.6	33.6	32.2	28.0	31.5	27.2	28.0	29.2	26.2	28.3
10	7	Chronic liver disease and cirrhosis	16.2	14.4	14.5	15.6	18.6	17.7	16.8	16.7	20.4	18.9
8	8	Intentional self-harm (suicide)	18.4	18.9	17.8	17.7	20.3	20.3	18.1	19.9	20.3	20.9
7	9	Alzheimer's disease	22.1	19.3	19.0	19.2	17.0	18.4	18.4	16.6	19.7	17.4
9	10	Influenza and pneumonia	18.3	16.4	18.8	18.8	17.9	16.9	16.4	13.8	16.3	12.8
12	11	Nephritis, nephrotic syndrome and nephrosis	11.0	13.1	13.2	13.4	12.6	13.5	13.0	13.0	12.1	12.6
16	12	Septicemia	6.0	6.2	6.7	8.9	8.8	7.6	9.3	8.3	9.3	9.3
13	13	Parkinson's disease	9.5	8.1	8.3	9.2	8.6	8.1	8.1	8.3	7.8	8.0
11	14	Essential primary hypertension and hypertensive renal disease	12.9	11.0	9.0	12.1	10.1	6.3	5.4	5.7	5.8	6.4
14	15	Assault (homicide)	8.4	9.3	7.9	7.2	8.5	7.4	8.8	7.4	7.4	6.6
ALL CAUSES			794.7	759.8	777.7	779.8	770.4	756.0	738.5	741.9	735.4	749.4

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates. Table is ranked by number of deaths for New Mexico. See numbers in Table M-14(a).

Mortality Section

Table M-15(a) Number of Deaths for 39 Selected Causes by Age

New Mexico, 2012

Cause of Death -- (ICD-10 Code)	0-4 Years	5-24 Years	25-64 Years	65+ Years	Unknown Age	All Ages
Tuberculosis (A16-A19)	0	0	1	4	0	5
Syphilis (A50-A53)	0	0	1	1	0	2
Human immunodeficiency virus (HIV) disease (B20-B24)	0	0	25	1	0	26
Malignant neoplasms (C00-C97)	7	23	1,021	2,370	1	3,422
Malignant neoplasm of stomach (C16)	0	0	30	54	0	84
Malignant neoplasms of colon, rectum and anus (C18-C21)	0	0	90	231	0	321
Malignant neoplasm of pancreas (C25)	0	0	64	150	0	214
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	0	0	174	580	1	755
Malignant neoplasm of breast (C50)	0	0	111	123	0	234
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	0	0	78	89	0	167
Malignant neoplasm of prostate (C61)	0	0	26	169	0	195
Malignant neoplasms of urinary tract (C64-C68)	0	2	45	148	0	195
Non-Hodgkin's lymphoma (C82-C85)	0	1	41	86	0	128
Leukemia (C91-C95)	1	4	38	101	0	144
Other malignant neoplasms (C00-C15,C17,C22-C24,C26-C32,C37-C49,C51-C52,C57-C60,C62-C63,C69-C81,C88,C90,C96-C97)	6	16	324	639	0	985
Diabetes mellitus (E10-E14)	0	2	190	456	0	648
Alzheimer's disease (G30)	0	0	6	363	0	369
Major cardiovascular diseases (I00-I78)	8	6	749	3,502	1	4,266
Diseases of heart (I00-I09, I11, I13, I20-I51)	8	3	620	2,674	1	3,306
Hypertensive heart disease with or without renal disease (I11,I13)	0	1	71	153	0	225
Ischemic heart diseases (I20-I25)	0	0	431	1,758	1	2,190
Other diseases of heart (I00-I09,I26-I51)	8	2	118	763	0	891
Essential (primary) hypertension and hypertensive renal disease (I10,I12)	0	0	22	119	0	141
Cerebrovascular diseases (I60-I69)	0	2	86	577	0	665
Atherosclerosis (I70)	0	0	5	39	0	44
Other diseases of circulatory system (I71-I78)	0	1	16	92	0	109
Influenza and pneumonia (J10-J18)	2	1	49	231	0	283
Chronic lower respiratory diseases (J40-J47)	3	1	132	825	0	961
Peptic ulcer (K25-K28)	0	0	14	27	0	41
Chronic liver disease and cirrhosis (K70,K73-K74)	0	2	313	121	0	436
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0	59	222	0	282
Pregnancy, childbirth and the puerperium (O00-O99)	0	0	1	0	0	1
Certain conditions originating in the perinatal period (P00-P96)	83	0	1	1	0	85
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	4	25	11	0	91
Sudden infant death syndrome (R95)	0	0	0	0	0	0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R94,R96-R99)	24	13	98	117	0	252
All other diseases (Residual) (A00-A09,A20-A49,A54-B19,B25-B99,D00-E07,E15-G25,G31-H93,I80-J06,J20-J39,J60-K22,K29-K66,K71-K72,K75-M99,N10-N15,N20-N23,N28-N98)	13	32	856	2,608	1	3,510
Motor vehicle accidents (V02-V04,V090,V092,V12-V14,V190-V192,V194-V196,V20-V79,V803-V805,V810-V811,V820-V821,V83-V86,V870-V878,V880-V888,V890,V892)	10	70	230	66	0	376
All other and unspecified accidents and adverse effects (V01,V05-V06,V091,V093-V099,V10-V11,V15-V18,V193,V198-V199,V800-V802,V806-V809,V812-V819,V822-V829,V879,V889,V891,V893,V899,V90-X59,Y40-Y86,Y88)	18	60	552	346	0	976
Intentional self-harm (suicide) (X60-X84,Y870)	0	64	303	68	0	435
Assault (homicide) (X85-Y09,Y871)	3	36	87	7	0	133
All other external (injury) causes (Y10-Y36,Y872,Y89)	2	3	31	4	0	40
Residual - Diseases/injuries other than NCHS 39 leading causes	0	0	0	1	0	1

Mortality Section

Table M-15(b) Death Rates for 39 Selected Causes by Age
New Mexico, 2012

Cause of Death -- (ICD-10 Code)	0-4 Years	5-24 Years	25-64 Years	65+ Years	All Ages
Tuberculosis (A16-A19)	0.0	0.0	0.1	1.4	0.2
Syphilis (A50-A53)	0.0	0.0	0.1	0.3	0.1
Human immunodeficiency virus (HIV) disease (B20-B24)	0.0	0.0	2.3	0.3	1.2
Malignant neoplasms (C00-C97)	4.8	4.0	95.1	808.9	147.3
Malignant neoplasm of stomach (C16)	0.0	0.0	2.8	18.4	3.6
Malignant neoplasms of colon, rectum and anus (C18-C21)	0.0	0.0	8.4	78.8	13.8
Malignant neoplasm of pancreas (C25)	0.0	0.0	6.0	51.2	9.1
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	0.0	0.0	16.2	198.0	32.3
Malignant neoplasm of breast (C50)	0.0	0.0	10.3	42.0	9.9
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	0.0	0.0	7.3	30.4	7.2
Malignant neoplasm of prostate (C61)	0.0	0.0	2.4	57.7	8.8
Malignant neoplasms of urinary tract (C64-C68)	0.0	0.4	4.2	50.5	8.5
Non-Hodgkin's lymphoma (C82-C85)	0.0	0.2	3.8	29.4	5.5
Leukemia (C91-C95)	0.7	0.7	3.5	34.5	6.3
Other malignant neoplasms (C00-C15,C17,C22-C24,C26-C32,C37-C49,C51-C52,C57-C60,C62-C63,C69-C81,C88,C90,C96-C97)	4.1	2.8	30.2	218.1	42.3
Diabetes mellitus (E10-E14)	0.0	0.4	17.7	155.6	28.3
Alzheimer's disease (G30)	0.0	0.0	0.6	123.9	17.4
Major cardiovascular diseases (I00-I78)	5.5	1.0	69.7	1,195.2	192.3
Diseases of heart (I00-I09, I11, I13, I20-I51)	5.5	0.5	57.7	912.6	148.5
Hypertensive heart disease with or without renal disease (I11,I13)	0.0	0.2	6.6	52.2	10.3
Ischemic heart diseases (I20-I25)	0.0	0.0	40.1	600.0	97.2
Other diseases of heart (I00-I09,I26-I51)	5.5	0.4	11.0	260.4	41.0
Essential (primary) hypertension and hypertensive renal disease (I10,I12)	0.0	0.0	2.0	40.6	6.4
Cerebrovascular diseases (I60-I69)	0.0	0.4	8.0	196.9	30.4
Atherosclerosis (I70)	0.0	0.0	0.5	13.3	2.0
Other diseases of circulatory system (I71-I78)	0.0	0.2	1.5	31.4	5.0
Influenza and pneumonia (J10-J18)	1.4	0.2	4.6	78.8	12.8
Chronic lower respiratory diseases (J40-J47)	2.1	0.2	12.3	281.6	42.8
Peptic ulcer (K25-K28)	0.0	0.0	1.3	9.2	1.8
Chronic liver disease and cirrhosis (K70,K73-K74)	0.0	0.4	29.1	41.3	18.9
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	0.7	0.0	5.5	75.8	12.6
Pregnancy, childbirth and the puerperium (O00-O99)	0.0	0.0	0.1	0.0	0.1
Certain conditions originating in the perinatal period (P00-P96)	57.0	0.0	0.1	0.3	4.0
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35.0	0.7	2.3	3.8	4.3
Sudden infant death syndrome (R95)	0.0	0.0	0.0	0.0	0.0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R94,R96-R99)	16.5	2.2	9.1	39.9	11.5
All other diseases (Residual) (A00-A09,A20-A49,A54-B19,B25-B99,D00-E07,E15-G25,G31-H93,I80-J06,J20-J39,J60-K22,K29-K66,K71-K72,K75-M99,N10-N15,N20-N23,N28-N98)	8.9	5.5	79.7	890.1	159.7
Motor vehicle accidents (V02-V04,V090,V092,V12-V14,V190-V192,V194-V196,V20-V79,V803-V805,V810-V811,V820-V821,V83-V86,V870-V878,V880-V888,V890,V892)	6.9	12.1	21.4	22.5	18.1
All other and unspecified accidents and adverse effects (V01,V05-V06,V091,V093-V099,V10-V11,V15-V18,V193,V198-V199,V800-V802,V806-V809,V812-V819,V822-V829,V879,V889,V891,V893,V899,V90-X59,Y40-Y86,Y88)	12.4	10.4	51.4	118.1	46.7
Intentional self-harm (suicide) (X60-X84,Y870)		11.1	28.2	23.2	20.9
Assault (homicide) (X85-Y09,Y871)	2.1	6.2	8.1	2.4	6.6
All other external (injury) causes (Y10-Y36,Y872,Y89)	1.4	0.5	2.9	1.4	1.8
Residual - Diseases/injuries other than NCHS 39 leading causes	0.0	0.0	0.0	0.3	0.0

Age-specific death rates are the numbers of deaths per 100,000 in specified age group.

Age-adjusted death rates (all ages) are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

See numbers in Table M-15(a).

Mortality Section

**Table M-16(a) Number of Deaths for 39 Selected Causes by Sex
New Mexico, 2012**

Cause of Death -- (ICD-10 Code)	Male	Female	Both Sexes
Tuberculosis (A16-A19)	4	1	5
Syphilis (A50-A53)	2	0	2
Human immunodeficiency virus (HIV) disease (B20-B24)	23	3	26
Malignant neoplasms (C00-C97)	1,870	1,552	3,422
Malignant neoplasm of stomach (C16)	45	39	84
Malignant neoplasms of colon, rectum and anus (C18-C21)	173	148	321
Malignant neoplasm of pancreas (C25)	117	97	214
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	407	348	755
Malignant neoplasm of breast (C50)	4	230	234
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	0	167	167
Malignant neoplasm of prostate (C61)	195	0	195
Malignant neoplasms of urinary tract (C64-C68)	128	67	195
Non-Hodgkin's lymphoma (C82-C85)	67	61	128
Leukemia (C91-C95)	89	55	144
Other malignant neoplasms (C00-C15,C17,C22-C24,C26-C32,C37-C49,C51-C52,C57-C60,C62-C63,C69-C81,C88,C90,C96-C97)	645	340	985
Diabetes mellitus (E10-E14)	352	296	648
Alzheimer's disease (G30)	110	259	369
Major cardiovascular diseases (I00-I78)	2,165	2,101	4,266
Diseases of heart (I00-I09, I11, I13, I20-I51)	1,744	1,562	3,306
Hypertensive heart disease with or without renal disease (I11,I13)	113	112	225
Ischemic heart diseases (I20-I25)	1,228	962	2,190
Other diseases of heart (I00-I09,I26-I51)	403	488	891
Essential (primary) hypertension and hypertensive renal disease (I10,I12)	55	86	141
Cerebrovascular diseases (I60-I69)	282	383	665
Atherosclerosis (I70)	17	27	44
Other diseases of circulatory system (I71-I78)	67	42	109
Influenza and pneumonia (J10-J18)	162	121	283
Chronic lower respiratory diseases (J40-J47)	482	479	961
Peptic ulcer (K25-K28)	22	19	41
Chronic liver disease and cirrhosis (K70,K73-K74)	300	136	436
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	125	282
Pregnancy, childbirth and the puerperium (O00-O99)	0	1	1
Certain conditions originating in the perinatal period (P00-P96)	51	34	85
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	41	91
Sudden infant death syndrome (R95)	0	0	0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R94,R96-R99)	140	112	252
All other diseases (Residual) (A00-A09,A20-A49,A54-B19,B25-B99,D00-E07,E15-G25,G31-H93,I80-J06,J20-J39,J60-K22,K29-K66,K71-K72,K75-M99,N10-N15,N20-N23,N28-N98)	1,736	1,774	3,510
Motor vehicle accidents (V02-V04,V090,V092,V12-V14,V190-V192,V194-V196,V20-V79,V803-V805,V810-V811,V820-V821,V83-V86,V870-V878,V880-V888,V890,V892)	266	110	376
All other and unspecified accidents and adverse effects (V01,V05-V06,V091,V093-V099,V10-V11,V15-V18,V193,V198-V199,V800-V802,V806-V809,V812-V819,V822-V829,V879,V889,V891,V893,V899,V90-X59,Y40-Y86,Y88)	605	371	976
Intentional self-harm (suicide) (X60-X84,Y870)	338	97	435
Assault (homicide) (X85-Y09,Y871)	110	23	133
All other external (injury) causes (Y10-Y36,Y872,Y89)	22	18	40
Residual - Diseases/injuries other than NCHS 39 leading causes	0	1	1

Mortality Section

Table M-16(b) Death Rates for 39 Selected Causes by Sex

New Mexico, 2012

Cause of Death -- (ICD-10 Code)	Male	Female	Both Sexes
Tuberculosis (A16-A19)	0.4	0.1	0.2
Syphilis (A50-A53)	0.2	0.0	0.1
Human immunodeficiency virus (HIV) disease (B20-B24)	2.2	0.3	1.2
Malignant neoplasms (C00-C97)	180.6	121.9	147.3
Malignant neoplasm of stomach (C16)	4.2	3.0	3.6
Malignant neoplasms of colon, rectum and anus (C18-C21)	16.6	11.6	13.8
Malignant neoplasm of pancreas (C25)	10.7	7.6	9.1
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	39.3	27.0	32.3
Malignant neoplasm of breast (C50)	0.3	17.8	9.9
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	0.0	13.3	7.2
Malignant neoplasm of prostate (C61)	21.2		8.8
Malignant neoplasms of urinary tract (C64-C68)	12.5	5.3	8.5
Non-Hodgkin's lymphoma (C82-C85)	6.4	4.8	5.5
Leukemia (C91-C95)	8.8	4.3	6.3
Other malignant neoplasms (C00-C15,C17,C22-C24,C26-C32,C37-C49,C51-C52,C57-C60,C62-C63,C69-C81,C88,C90,C96-C97)	60.5	27.1	42.3
Diabetes mellitus (E10-E14)	34.2	22.9	28.3
Alzheimer's disease (G30)	13.0	20.0	17.4
Major cardiovascular diseases (I00-I78)	225.3	163.4	192.3
Diseases of heart (I00-I09, I11, I13, I20-I51)	180.0	121.4	148.5
Hypertensive heart disease with or without renal disease (I11,I13)	11.4	9.0	10.3
Ischemic heart diseases (I20-I25)	124.7	74.2	97.2
Other diseases of heart (I00-I09,I26-I51)	44.0	38.2	41.0
Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5.9	6.6	6.4
Cerebrovascular diseases (I60-I69)	30.3	29.8	30.4
Atherosclerosis (I70)	1.8	2.1	2.0
Other diseases of circulatory system (I71-I78)	7.3	3.4	5.0
Influenza and pneumonia (J10-J18)	17.7	9.6	12.8
Chronic lower respiratory diseases (J40-J47)	50.2	37.4	42.8
Peptic ulcer (K25-K28)	2.2	1.5	1.8
Chronic liver disease and cirrhosis (K70,K73-K74)	26.6	11.7	18.9
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16.3	9.9	12.6
Pregnancy, childbirth and the puerperium (O00-O99)	0.0	0.1	0.1
Certain conditions originating in the perinatal period (P00-P96)	4.7	3.3	4.0
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4.7	3.9	4.3
Sudden infant death syndrome (R95)	0.0	0.0	0.0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R94,R96-R99)	13.7	9.2	11.5
All other diseases (Residual) (A00-A09,A20-A49,A54-B19,B25-B99,D00-E07,E15-G25,G31-H93,I80-J06,J20-J39,J60-K22,K29-K66,K71-K72,K75-M99,N10-N15,N20-N23,N28-N98)	181.1	141.2	159.7
Motor vehicle accidents (V02-V04,V090,V092,V12-V14,V190-V192,V194-V196,V20-V79,V803-V805,V810-V811,V820-V821,V83-V86,V870-V878,V880-V888,V890,V892)	26.2	10.5	18.1
All other and unspecified accidents and adverse effects (V01,V05-V06,V091,V093-V099,V10-V11,V15-V18,V193,V198-V199,V800-V802,V806-V809,V812-V819,V822-V829,V879,V889,V891,V893,V899,V90-X59,Y40-Y86,Y88)	61.5	32.8	46.7
Intentional self-harm (suicide) (U03,X60-X84,Y870)	33.0	9.4	20.9
Assault (homicide) (U01-U02,X85-Y09,Y871)	11.0	2.2	6.6
All other external (injury) causes (Y10-Y36,Y872,Y89)	2.1	1.6	1.8
Residual - Diseases/injuries other than NCHS 39 leading causes	0.0	0.1	0.0

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

See numbers in Table M-16(a).

Mortality Section

**Table M-17 (a) Number of Deaths for 39 Selected Causes by Health Region
New Mexico, 2012**

Cause of Death – (ICD-10 Code)	Northwest	Northeast	Metro	Southeast	Southwest	Unknown	
						Region	All Regions
Tuberculosis (A16-A19)	2	1	0	0	2	0	5
Syphilis (A50-A53)	0	0	0	1	1	0	2
Human immunodeficiency virus (HIV) disease (B20-B24)	4	1	13	3	5	0	26
Malignant neoplasms (C00-C97)	301	528	1,411	480	702	0	3,422
Malignant neoplasm of stomach (C16)	15	13	36	12	8	0	84
Malignant neoplasms of colon, rectum and anus (C18-C21)	24	54	135	54	54	0	321
Malignant neoplasm of pancreas (C25)	12	37	81	32	52	0	214
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	48	109	293	134	171	0	755
Malignant neoplasm of breast (C50)	25	25	109	26	49	0	234
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	20	34	55	24	34	0	167
Malignant neoplasm of prostate (C61)	20	32	75	26	42	0	195
Malignant neoplasms of urinary tract (C64-C68)	21	29	75	30	40	0	195
Non-Hodgkin's lymphoma (C82-C85)	10	16	61	16	25	0	128
Leukemia (C91-C95)	10	23	56	22	33	0	144
Other malignant neoplasms (C00-C15,C17,C22-C24,C26-C32,C37-C49,C51-C52,C57-C60,C62-C63,C69-C81,C88,C90,C96-C97)	96	156	435	104	194	0	985
Diabetes mellitus (E10-E14)	91	87	249	103	118	0	648
Alzheimer's disease (G30)	33	41	179	57	59	0	369
Major cardiovascular diseases (I00-I78)	346	632	1,724	728	835	1	4,266
Diseases of heart (I00-I09, I11, I13, I20-I51)	257	499	1,326	599	625	0	3,306
Hypertensive heart disease with or without renal disease (I11,I13)	19	47	116	15	28	0	225
Ischemic heart diseases (I20-I25)	153	307	862	430	438	0	2,190
Other diseases of heart (I00-I09,I26-I51)	85	145	348	154	159	0	891
Essential (primary) hypertension and hypertensive renal disease (I10,I12)	14	20	66	24	17	0	141
Cerebrovascular diseases (I60-I69)	60	81	282	92	149	1	665
Atherosclerosis (I70)	4	14	6	2	18	0	44
Other diseases of circulatory system (I71-I78)	10	18	44	11	26	0	109
Influenza and pneumonia (J10-J18)	46	40	113	39	45	0	283
Chronic lower respiratory diseases (J40-J47)	76	106	363	228	188	0	961
Peptic ulcer (K25-K28)	7	4	13	2	15	0	41
Chronic liver disease and cirrhosis (K70,K73-K74)	91	82	156	46	61	0	436
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	41	104	49	66	0	282
Pregnancy, childbirth and the puerperium (O00-O99)	0	0	0	1	0	0	1
Certain conditions originating in the perinatal period (P00-P96)	8	10	39	16	12	0	85
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	14	35	14	16	0	91
Sudden infant death syndrome (R95)	0	0	0	0	0	0	
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R94,R96-R99)	14	31	47	99	61	0	252
All other diseases (Residual) (A00-A09,A20-A49,A54-B19,B25-B99,D00-E07,E15-G25,G31-H93,I80-J06,J20-J39,J60-K22,K29-K66,K71-K72,K75-M99,N10-N15,N20-N23,N28-N98)	368	517	1,529	432	663	1	3,510
Motor vehicle accidents (V02-V04,V090,V092,V12-V14,V190-V192,V194-V196,V20-V79,V803-V805,V810-V811,V820-V821,V83-V86,V870-V878,V880-V888,V890,V892)	79	63	118	69	47	0	376
All other and unspecified accidents and adverse effects (V01,V05-V06,V091,V093-V099,V10-V11,V15-V18,V193,V198-V199,V800-V802,V806-V809,V812-V819,V822-V829,V879,V889,V891,V893,V899,V90-X59,Y40-Y86,Y88)	119	190	397	116	154	0	976
Intentional self-harm (suicide) (X60-X84,Y870)	64	71	169	55	75	1	435
Assault (homicide) (X85-Y09,Y871)	26	24	52	24	7	0	133
All other external (injury) causes (Y10-Y36,Y872,Y89)	8	5	12	7	8	0	40
Residual - Diseases/injuries other than NCHS 39 leading causes	1	0	0	0	0	0	1

See Technical Appendix for information on health regions.

Mortality Section

**Table M-17(b) Death Rates for 39 Selected Causes by Health Region
New Mexico, 2012**

Cause of Death – (ICD-10 Code)	Northwest	Northeast	Metro	Southeast	Southwest	All Regions
Tuberculosis (A16-A19)	1.1	0.3	0.0	0.0	0.5	0.2
Syphilis (A50-A53)	0.0	0.0	0.0	0.3	0.2	0.1
Human immunodeficiency virus (HIV) disease (B20-B24)	1.7	0.5	1.4	1.1	1.6	1.2
Malignant neoplasms (C00-C97)	144.7	139.3	147.1	151.2	153.5	147.3
Malignant neoplasm of stomach (C16)	7.4	3.3	3.8	3.8	1.6	3.6
Malignant neoplasms of colon, rectum and anus (C18-C21)	11.5	14.5	14.2	16.8	11.9	13.8
Malignant neoplasm of pancreas (C25)	5.6	9.7	8.3	10.1	11.2	9.1
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	23.6	28.4	30.8	41.7	36.6	32.3
Malignant neoplasm of breast (C50)	11.5	6.3	11	8.6	10.6	9.9
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	9.1	8.9	5.7	7.6	8	7.2
Malignant neoplasm of prostate (C61)	10.8	8.9	8.2	8.2	9.5	8.8
Malignant neoplasms of urinary tract (C64-C68)	9.5	7.4	8.2	9.5	8.6	8.5
Non-Hodgkin's lymphoma (C82-C85)	4.6	3.8	6.4	4.9	5.9	5.5
Leukemia (C91-C95)	4.5	6.3	6	7	7.1	6.3
Other malignant neoplasms (C00-C15,C17,C22-C24,C26-C32,C37-C49,C51-C52,C57-C60,C62-C63,C69-C81,C88,C90,C96-C97)	46.6	41.9	44.6	33	42.5	42.3
Diabetes mellitus (E10-E14)	43.7	24	26.6	32.2	27.1	28.3
Alzheimer's disease (G30)	18.5	12.4	20.2	18.8	14	17.4
Major cardiovascular diseases (I00-I78)	175.3	180.4	187.7	234.7	190.9	192.3
Diseases of heart (I00-I09, I11, I13, I20-I51)	128.7	141.8	143.8	192.8	141.8	148.5
Hypertensive heart disease with or without renal disease (I11,I13)	9.1	13.9	12.8	4.8	6.2	10.3
Ischemic heart diseases (I20-I25)	76.1	84.8	92.6	137.7	97.6	97.2
Other diseases of heart (I00-I09,I26-I51)	43.5	43.2	38.3	50.2	38	41
Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6.8	5.8	7.4	7.7	3.8	6.4
Cerebrovascular diseases (I60-I69)	32.3	23.2	31	29.9	34.8	30.4
Atherosclerosis (I70)	2	4.1	0.7	0.7	4.3	2
Other diseases of circulatory system (I71-I78)	5	5.4	4.8	3.7	6.2	5
Influenza and pneumonia (J10-J18)	23.2	12	12.6	12.1	10.4	12.8
Chronic lower respiratory diseases (J40-J47)	38.3	30.1	39.8	72.3	41.7	42.8
Peptic ulcer (K25-K28)	3.4	1	1.3	0.6	3.3	1.8
Chronic liver disease and cirrhosis (K70,K73-K74)	40.8	22.6	16.1	14.6	14	18.9
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11.1	11.6	11.3	15.9	14.6	12.6
Pregnancy, childbirth and the puerperium (O00-O99)	0.0	0.0	0.0	0.5	0.0	0.1
Certain conditions originating in the perinatal period (P00-P96)	2.9	4.1	4.4	4.8	3.1	4
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4.7	5.2	3.9	4.6	4.3	4.3
Sudden infant death syndrome (R95)	0.0	0.0	0.0	0.0	0.0	0.0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R94,R96-R99)	6.8	10.1	5	31.5	14.8	11.5
All other diseases (Residual) (A00-A09,A20-A49,A54-B19,B25-B99,D00-E07,E15-G25,G31-H93,I80-J06,J20-J39,J60-K22,K29-K66,K71-K72,K75-M99,N10-N15,N20-N23,N28-N98)	188.2	149.5	166.3	140.2	157.2	159.7
Motor vehicle accidents (V02-V04,V090,V092,V12-V14,V190-V192,V194-V196,V20-V79,V803-V805,V810-V811,V820-V821,V83-V86,V870-V878,V880-V888,V890,V892)	35.2	21.6	12.7	24.8	12.8	18.1
All other and unspecified accidents and adverse effects (V01,V05-V06,V091,V093-V099,V10-V11,V15-V18,V193,V198-V199,V800-V802,V806-V809,V812-V819,V822-V829,V879,V889,V891,V893,V899,V90-X59,Y40-Y86,Y88)	54.8	63.3	43.5	40.1	40.4	46.7
Intentional self-harm (suicide) (X60-X84,Y870)	29.7	24.3	18.3	19.4	20.2	20.9
Assault (homicide) (X85-Y09,Y871)	11.7	9	5.7	8.8	2	6.6
All other external (injury) causes (Y10-Y36,Y872,Y89)	3.6	1.9	1.2	2.2	2.2	1.8
Residual - Diseases/injuries other than NCHS 39 leading causes	0.6	0.0	0.0	0.0	0.0	0.0

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates and health regions. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution. See numbers in Table M-17(a).

Mortality Section

**Table M-18(a) Number of Deaths for 39 Selected Causes by Race/Ethnicity
New Mexico, 2012**

Cause of Death -- (ICD-10 Code)	American	Asian or	Black or			Unknown	All
	Indian or Alaska Native	Pacific Islander	African American	Hispanic	White	Race/ Ethnicity	
Tuberculosis (A16-A19)	3	0	0	1	1	0	5
Syphilis (A50-A53)	0	0	0	2	0	0	2
Human immunodeficiency virus (HIV) disease (B20-B24)	4	0	3	11	8	0	26
Malignant neoplasms (C00-C97)	180	26	64	1,125	2,019	8	3,422
Malignant neoplasm of stomach (C16)	17	2	2	33	30	0	84
Malignant neoplasms of colon, rectum and anus (C18-C21)	20	3	7	118	173	0	321
Malignant neoplasm of pancreas (C25)	7	1	4	70	131	1	214
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	19	6	8	215	505	2	755
Malignant neoplasm of breast (C50)	10	0	6	72	145	1	234
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	18	2	2	55	90	0	167
Malignant neoplasm of prostate (C61)	10	1	9	65	110	0	195
Malignant neoplasms of urinary tract (C64-C68)	18	0	6	53	117	1	195
Non-Hodgkin's lymphoma (C82-C85)	7	1	3	36	80	1	128
Leukemia (C91-C95)	4	1	1	57	81	0	144
Other malignant neoplasms (C00-C15, C17, C22-C24, C26-C32, C37-C49, C51-C52, C57-C60, C62-C63, C69-C81, C88, C90, C96-C97)	50	9	16	351	557	2	985
Diabetes mellitus (E10-E14)	97	6	18	298	227	2	648
Alzheimer's disease (G30)	9	1	2	96	261	0	369
Major cardiovascular diseases (I00-I78)	182	31	73	1,297	2,671	10	4,264
Diseases of heart (I00-I09, I11, I13, I20-I51)	131	20	58	999	2,088	9	3,305
Hypertensive heart disease with or without renal disease (I11, I13)	6	2	7	89	120	1	225
Ischemic heart diseases (I20-I25)	93	15	40	648	1,386	8	2,190
Other diseases of heart (I00-I09, I26-I51)	32	3	11	262	582	1	891
Essential (primary) hypertension and hypertensive renal disease (I10, I12)	10	1	4	37	89	0	141
Cerebrovascular diseases (I60-I69)	33	8	8	216	398	2	665
Atherosclerosis (I70)	3	0	0	13	28	0	44
Other diseases of circulatory system (I71-I78)	5	2	3	32	67	0	109
Influenza and pneumonia (J09-J18)	41	0	5	84	153	0	283
Chronic lower respiratory diseases (J40-J47)	18	1	11	218	712	1	961
Peptic ulcer (K25-K28)	4	0	0	18	19	0	41
Chronic liver disease and cirrhosis (K70, K73-K74)	92	0	3	211	127	3	436
Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19, N25-N27)	26	1	4	128	123	0	282
Pregnancy, childbirth and the puerperium (O00-O99)	0	0	0	1	0	0	1
Certain conditions originating in the perinatal period (P00-P96)	6	2	1	48	28	0	85
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0	2	49	32	0	91
Sudden infant death syndrome (R95)	0	0	0	0	0	0	0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R94, R96-R99)	17	0	6	88	138	3	252
All other diseases (Residual) (A00-A09, A20-A49, A54-B19, B25-B99, D00-E07, E15-G25, G31-H93, I80-J06, J20-J39, J60-K22, K29-K66, K71-K72, K75-M99, N10-N15, N20-N23, N28-N98)	283	23	49	1,180	1,961	14	3,510
Motor vehicle accidents (V02-V04, V090, V092, V12-V14, V190-V192, V194-V196, V20-V79, V803-V805, V810-V811, V820-V821, V83-V86, V870-V878, V880-V888, V890, V892)	82	3	7	152	130	2	376
All other and unspecified accidents and adverse effects (V01, V05-V06, V091, V093-V099, V10-V11, V15-V18, V193, V198-V199, V800-V802, V806-V809, V812-V819, V822-V829, V879, V889, V891, V893, V899, V90-X59, Y40-Y86, Y88)	106	2	14	402	444	8	976
Intentional self-harm (suicide) (*U03, X60-X84, Y870)	55	5	4	134	233	4	435
Assault (homicide) (*U01-*U02, X85-Y09, Y871)	19	0	10	80	23	1	133
All other external (injury) causes (Y10-Y36, Y872, Y89)	9	0	2	16	12	1	40
Residual - Diseases/injuries other than NCHS 39 leading causes					1		1

See Technical Appendix for information on race/ethnicity.

Mortality Section

**Table M-18(b) Death Rates for 39 Selected Causes by Race/Ethnicity
New Mexico, 2012**

Cause of Death -- (ICD-10 Code)	American Indian or Alaska Native	Asian or Pacific Islander	Black or African American	Hispanic	White	All Races
Tuberculosis (A16-A19)	3.2	0.0	0.0	0.2	0.1	0.2
Syphilis (A50-A53)	0.0	0.0	0.0	0.2	0.0	0.1
Human immunodeficiency virus (HIV) disease (B20-B24)	2.4	0.0	6.5	1.3	0.9	1.2
Malignant neoplasms (C00-C97)	135.8	109.3	200.8	144.9	149.7	147.3
Malignant neoplasm of stomach (C16)	13.3	12.0	5.9	4.1	2.4	3.6
Malignant neoplasms of colon, rectum and anus (C18-C21)	15.1	13.5	26.8	15.6	12.8	13.8
Malignant neoplasm of pancreas (C25)	5.2	5.5	11.1	9.0	9.7	9.0
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	15.3	30.0	23.8	28.5	36.7	32.3
Malignant neoplasm of breast (C50)	5.7	0.0	17.0	8.7	11.1	9.8
Malignant neoplasms of cervix uteri, corpus uteri and ovary (C53-C56)	12.2	5.7	4.7	6.7	7.1	7.2
Malignant neoplasm of prostate (C61)	8.4	3.7	31.6	9.4	7.9	8.8
Malignant neoplasms of urinary tract (C64-C68)	13.8	0.0	18.4	7.3	8.7	8.4
Non-Hodgkin's lymphoma (C82-C85)	5.3	3.2	8.7	4.5	5.8	5.5
Leukemia (C91-C95)	3.0	5.5	3.6	7.2	6.0	6.3
Other malignant neoplasms (C00-C15, C17, C22-C24, C26-C32, C37-C49, C51-C52, C57-C60, C62-C63, C69-C81, C88, C90, C96-C97)	38.4	30.1	49.0	43.9	41.6	42.2
Diabetes mellitus (E10-E14)	77.1	32.0	53.9	39.3	17.0	28.2
Alzheimer's disease (G30)	9.5	8.2	6.4	15.6	19.2	17.4
Major cardiovascular diseases (I00-I78)	152.0	167.3	218.2	185.2	198.0	192.3
Diseases of heart (I00-I09, I11, I13, I20-I51)	107.9	108.7	176.4	141.7	155.0	148.5
Hypertensive heart disease with or without renal disease (I11, I13)	4.7	4.8	21.8	12.2	9.4	10.2
Ischemic heart diseases (I20-I25)	77.3	84.9	121.3	91.0	101.8	96.9
Other diseases of heart (I00-I09, I26-I51)	25.9	18.9	33.4	38.4	43.8	41.0
Essential (primary) hypertension and hypertensive renal disease (I10, I12)	8.2	8.2	10.3	5.4	6.5	6.4
Cerebrovascular diseases (I60-I69)	29.4	45.6	23.0	31.5	29.6	30.3
Atherosclerosis (I70)	2.5	0.0	0.0	1.9	2.0	2.0
Other diseases of circulatory system (I71-I78)	4.1	4.8	8.4	4.7	5.0	5.0
Influenza and pneumonia (J09-J18)	36.0	0.0	15.6	12.1	11.4	12.8
Chronic lower respiratory diseases (J40-J47)	16.4	8.2	34.3	31.2	51.8	42.8
Peptic ulcer (K25-K28)	3.0	0.0	0.0	2.3	1.4	1.8
Chronic liver disease and cirrhosis (K70, K73-K74)	57.0	0.0	6.2	23.5	10.9	18.8
Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19, N25-N27)	20.1	8.2	13.2	17.5	9.3	12.6
Pregnancy, childbirth and the puerperium (O00-O99)	0.0	0.0	0.0	0.1	0.0	0.1
Certain conditions originating in the perinatal period (P00-P96)	2.6	7.9	2.1	3.8	5.0	4.0
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.5	0.0	4.4	4.5	4.0	4.3
Sudden infant death syndrome (R95)	0.0	0.0	0.0	0.0	0.0	0.0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R94, R96-R99)	11.4	0.0	16.6	10.0	12.0	11.3
All other diseases (Residual) (A00-A09, A20-A49, A54-B19, B25-B99, D00-E07, E15-G25, G31-H93, I80-J06, J20-J39, J60-K22, K29-K66, K71-K72, K75-M99, N10-N15, N20-N23, N28-N98)	219.6	146.1	150.8	161.8	149.3	159.1
V79, V803-V805, V810-V811, V820-V821, V83-V86, V870-V878, V880-V888, V890, V892)	45.9	15.2	18.1	16.3	13.4	18.0
All other and unspecified accidents and adverse effects (V01, V05-V06, V091, V093-V099, V10-V11, V15-V18, V193, V198-V199, V800-V802, V806-V809, V812-V819, V822-V829, V879, V889, V891, V893, V899, V90-X59, Y40-Y86, Y88)	65.5	11.8	35.4	48.4	42.3	46.4
Intentional self-harm (suicide) (*U03, X60-X84, Y870)	31.8	13.3	9.0	14.4	25.1	20.7
Assault (homicide) (*U01-*U02, X85-Y09, Y871)	10.3	0.0	21.7	8.2	2.8	6.5
All other external (injury) causes (Y10-Y36, Y872, Y89)	5.3	0.0	3.9	1.7	1.2	1.8

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates and race/ethnicity. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

See numbers in Table M-18(a).

Mortality Section

**Table M-19 Number of Injury Deaths by Sex, County, and Health Region
New Mexico, 2012, and United States, 2010**

County	Total Injuries				Accidents (Unintentional Injuries)										Intentional Injuries				Other Injury			
	(Unintentional and Intentional)		Motor Vehicle Accidents		All other accidents		Intentional self-harm (suicide)		Assault (homicide)		Other Injury		Assault (homicide)		Other Injury		Other Injury					
	Male	Female	Both	Sexes	Male	Female	Both	Sexes	Male	Female	Both	Sexes	Male	Female	Both	Sexes	Male	Female	Both	Sexes		
United States	123,477	59,824	183,301	24,723	10,609	35,332	51,198	34,329	85,527	30,277	8,087	38,364	12,774	3,485	16,259	4,505	3,314	7,819	5	3		
New Mexico	1,341	619	1,960	266	110	376	595	360	955	338	97	435	110	23	133	32	29	61	5	6		
Bernalillo	389	189	578	61	20	81	191	118	309	93	39	132	35	7	42	9	5	14	8	10		
Catron	8	5	13	0	2	2	1	2	3	7	1	8	0	0	0	0	0	0	0	0		
Chaves	37	26	63	8	5	13	11	17	28	12	1	13	6	0	6	0	0	3	0	3		
Cibola	24	7	31	4	4	8	10	3	13	6	0	6	3	0	3	1	0	1	0	1		
Colfax	13	4	17	3	3	6	5	1	6	4	0	4	1	0	1	0	0	0	0	0		
Curry	19	13	32	4	1	5	7	10	17	4	2	6	3	0	3	1	0	1	0	1		
De Baca	3	0	3	0	0	1	1	0	1	1	0	1	0	0	0	0	0	0	0	0		
Doña Ana	88	52	140	14	7	21	39	31	70	29	6	35	4	0	4	2	8	10	8	10		
Eddy	37	19	56	10	4	14	9	12	21	13	1	14	5	1	6	0	1	1	1	1		
Grant	23	11	34	3	0	3	11	6	17	7	3	10	1	0	1	0	2	3	2	3		
Guadalupe	2	2	4	2	0	2	0	0	0	0	2	2	0	0	0	0	0	0	0	0		
Harding	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Hidalgo	1	1	2	0	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0	0		
Lea	40	21	61	18	8	26	11	10	21	5	3	8	4	0	4	2	0	2	0	2		
Lincoln	15	9	24	2	0	2	4	8	12	7	7	8	2	0	2	0	0	0	0	0		
Los Alamos	6	4	10	1	0	1	5	1	6	0	0	0	0	0	0	0	0	3	3	3		
Luna	9	7	16	0	2	2	5	5	10	4	0	4	0	0	0	0	0	0	0	0		
McKinley	102	18	120	27	9	36	33	4	47	29	2	31	10	2	12	3	1	4	1	4		
Mora	9	2	11	4	1	5	4	1	5	1	0	1	0	0	0	0	0	0	0	0		
Otero	39	21	60	11	1	12	13	18	31	13	1	14	1	0	1	1	1	2	1	2		
Quay	4	11	15	1	2	3	0	4	4	3	1	4	0	3	3	0	1	1	1	1		
Rio Arriba	62	19	81	14	3	17	33	13	46	7	3	10	7	0	7	1	0	1	1	1		
Roosevelt	14	3	17	5	0	5	7	3	10	1	0	1	0	0	0	1	1	0	1	1		
Sandoval	77	26	103	10	9	19	40	13	53	22	3	25	3	0	3	2	1	3	2	3		
San Juan	100	45	145	25	10	35	43	24	67	20	7	27	9	2	11	3	2	5	2	5		
San Miguel	23	9	32	7	1	8	11	7	18	3	1	4	0	0	0	2	0	2	0	2		
Santa Fe	100	46	146	11	4	15	54	27	81	25	12	37	10	2	12	0	1	1	1	1		
Sierra	10	4	14	1	2	3	6	2	8	1	0	1	1	0	1	1	0	1	0	1		
Socorro	8	4	12	3	0	3	3	3	6	2	1	3	0	0	0	0	0	0	0	0		
Taos	33	17	50	2	6	8	17	7	24	9	3	12	3	1	4	2	0	2	0	2		
Torrance	8	4	12	5	1	6	3	2	5	0	1	1	0	0	0	0	0	0	0	0		
Union	2	0	2	1	1	1	0	0	0	1	0	1	0	0	0	0	0	0	0	0		
Valencia	35	20	55	8	4	12	17	8	25	8	3	11	2	5	7	0	0	0	0	0		
Unknown County	1	0	1	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0		
Health Region																						
Northwest	226	70	296	56	23	79	86	31	117	55	9	64	22	4	26	7	3	10	3	10		
Northeast	250	103	353	45	18	63	129	57	186	50	21	71	21	3	24	5	4	9	4	9		
Metrol	509	239	748	84	34	118	251	141	392	123	46	169	40	12	52	11	6	17	6	17		
Southeast	169	102	271	49	20	69	50	64	114	46	9	55	20	4	24	4	4	5	9	9		
Southwest	186	105	291	32	15	47	79	67	146	63	12	75	7	0	7	5	11	16	11	16		
Unknown Region	1	0	1	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0		

See Technical Appendix for information on health regions. Total injuries include unintentional, intentional, and other injuries. Other injury includes legal intervention; operations of war; complications of medical and surgical care; and events of undetermined intent.

Mortality Section

**Table M-20 Death Rates for Injury by Sex, County, and Health Region
New Mexico, 2012, and United States, 2010**

	Total Injuries (Unintentional and Intentional)						Accidents (Unintentional Injuries) (Motor Vehicle Accidents)						All other accidents						Intentional Injuries (suicide)						Assault (homicide)						Other Injury						
	Male		Female		Sexes		Male		Female		Sexes		Male		Female		Sexes		Male		Female		Sexes		Male		Female		Sexes		Male		Female		Sexes		
United States	N/A	N/A	57.1	16.2	6.5	11.3	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A		
New Mexico	133.7	56.4	94.1	26.2	10.5	18.1	60.5	31.9	45.8	33.0	19.8	5.0	9.4	20.9	11.0	2.2	6.6	3.1	2.7	6.2	2.7	6.6	3.1	2.2	6.6	3.1	2.7	6.2	2.7	6.6	3.1	2.2	6.6	3.1	2.7		
County																																					
Bernalillo	118.5	51.8	83.5	17.9	5.6	11.4	59.7	31.6	44.9	27.7	11.2	19.0	10.4	2.1	6.2	2.7	6.2	2.7	6.2	2.7	6.2	2.7	6.2	2.7	6.2	2.7	6.2	2.7	6.2	2.7	6.2	2.7	6.2	2.7	6.2		
Carton	276.0	543.4	395.8	0.0	0.0	261.4	118.1	27.6	261.4	133.0	248.5	20.6	144.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Cheaves	116.3	71.7	93.7	23.6	14.7	18.9	46.5	48.6	48.6	42.7	0.0	24.7	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	
Chibola	172.3	57.8	114.5	24.0	33.2	28.1	71.6	48.6	48.6	42.7	0.0	24.7	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	0.0	11.4	12.0	
Colfax	181.6	50.3	113.3	49.7	25.1	38.1	58.5	25.2	36.8	60.4	0.0	31.8	13.0	0.0	6.6	0.0	6.6	0.0	6.6	0.0	6.6	0.0	6.6	0.0	6.6	0.0	6.6	0.0	6.6	0.0	6.6	0.0	6.6	0.0	6.6	0.0	
Curry	90.4	49.2	69.4	17.9	3.8	11.0	69.4	36.6	37.0	15.1	8.8	12.0	14.5	0.0	7.3	4.4	0.0	7.3	4.4	0.0	7.3	4.4	0.0	7.3	4.4	0.0	7.3	4.4	0.0	7.3	4.4	0.0	7.3	4.4	0.0	7.3	4.4
De Baca	190.9	0.0	80.6	65.7	0.0	29.6	74.9	0.0	24.6	50.3	0.0	26.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Dona Ana	89.4	46.4	67.6	13.0	7.2	10.0	42.3	27.0	34.4	28.8	5.9	17.1	3.5	0.0	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8		
Eddy	142.5	67.4	105.2	38.7	16.7	27.0	38.0	39.1	39.2	47.6	5.1	26.4	18.2	4.1	11.3	0.0	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5		
Grant	160.3	62.9	110.2	16.1	0.0	8.1	69.9	27.2	48.0	58.0	23.8	39.6	8.1	0.0	4.2	8.1	11.9	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	
Guadalupe	76.9	157.6	102.2	76.9	0.0	43.2	0.0	0.0	0.0	0.0	157.6	59.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Harding	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Hidalgo	32.3	42.8	37.6	0.0	42.8	0.0	21.4	32.3	0.0	16.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Lea	127.7	68.9	100.5	55.7	26.9	41.7	35.9	31.3	35.0	16.8	10.7	14.0	12.5	0.0	6.5	6.8	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2	3.2		
Lincoln	128.6	87.6	107.5	13.6	0.0	6.5	31.0	72.8	52.0	55.7	14.9	34.3	28.4	0.0	14.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Los Alamos	75.3	38.5	57.7	11.2	0.0	5.7	64.1	6.3	36.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Luna	73.9	52.6	63.5	0.0	13.4	6.7	39.8	39.2	39.6	34.1	0.0	17.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
McKinley	311.4	49.5	174.7	82.2	24.4	52.0	98.5	12.8	53.1	91.0	4.5	46.2	30.0	5.2	17.2	9.8	2.6	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1		
Mora	371.9	117.8	245.2	182.2	59.6	119.0	129.1	58.2	96.4	60.6	0.0	29.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Otero	117.3	62.2	90.8	32.5	3.4	17.4	39.2	51.7	47.1	39.0	2.5	20.6	4.4	0.0	2.3	2.3	4.6	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4	3.4		
Quay	104.8	215.3	162.7	21.8	41.0	31.7	88.8	45.7	45.7	83.0	20.5	51.9	0.0	53.1	27.2	0.0	11.9	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2		
Rio Arriba	335.8	98.2	217.4	77.9	13.6	45.2	181.2	70.4	126.3	37.9	14.1	26.2	34.2	0.0	17.3	4.5	0.0	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.3		
Roosevelt	162.4	26.9	90.4	60.7	0.0	27.6	78.7	26.9	51.0	14.2	0.0	7.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Sandoval	122.6	37.1	77.8	17.2	13.6	15.2	65.6	17.8	40.3	32.6	4.8	18.3	5.0	0.0	2.4	2.3	0.9	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5		
San Juan	164.4	71.5	116.8	39.4	16.0	27.2	72.8	36.8	54.8	31.9	12.6	22.2	14.9	3.1	8.6	5.4	3.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1		
San Miguel	159.3	58.3	105.9	53.5	4.6	29.7	80.5	43.6	58.3	14.7	10.1	12.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Santa Fe	146.0	56.2	98.3	15.6	5.4	10.5	80.6	31.8	53.2	34.0	15.5	24.8	15.8	2.4	9.2	0.0	1.2	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6		
Sierra	117.0	29.8	73.0	16.8	0.0	14.7	15.6	68.3	15.1	7.0	0.0	3.6	16.8	0.0	8.4	8.0	0.0	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7		
Socorro	93.3	45.7	69.6	42.7	0.0	21.8	31.1	34.8	33.3	19.5	10.9	14.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Taos	208.9	92.2	151.7	11.2	28.0	20.3	106.4	35.7	71.8	55.8	20.5	37.7	24.1	0.0	16.3	11.3	0.0	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6	5.6		
Torrance	91.0	43.1	68.5	58.3	8.7	35.6	32.7	27.6	29.6	0.0	6.8	3.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Union	59.8	0.0	35.5	31.5	0.0	15.2	0.0	0.0	0.0	28.3	0.0	20.3	0.0																								

Mortality Section

**Table M-21 Number of Deaths due to Firearms, Alcohol, Drugs, and Injury at Work
by Sex, County, and Health Region
New Mexico, 2012, and United States, 2010**

	Firearms			Alcohol-Induced			Drug-Induced			Injury at Work		
	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes
United States	27,356	4,316	31,672	19,038	6,654	25,692	24,376	16,017	40,393	3,829	328	4,157
New Mexico	280	37	317	383	107	490	299	198	497	38	3	41
County												
Bernalillo	89	9	98	110	21	131	115	79	194	11	1	12
Catron	7	1	8	0	0	0	0	2	2	0	0	0
Chaves	17	0	17	7	0	7	3	11	14	2	0	2
Cibola	3	0	3	8	7	15	2	0	2	0	0	0
Colfax	3	0	3		1	1	1	1	2	0	0	0
Curry	4	1	5	6	1	7	1	1	2	2	0	2
De Baca	1	0	1	0	0	0	0	0	0	0	0	0
Dona Ana	21	1	22	23	1	24	22	17	39	2	1	3
Eddy	13	2	15	2	2	4	4	5	9	3	1	4
Grant	3	2	5	7	0	7	2	3	5	0	0	0
Guadalupe	0	0	0	3	0	3	0	1	1	0	0	0
Harding	0	0	0	0	0	0	0	0	0	0	0	0
Hidalgo	0	0	0	3	0	3	0	0	0	1	0	1
Lea	5	1	6	9	1	10	3	3	6	4	0	4
Lincoln	5	0	5	4	0	4	1	5	6	0	0	0
Los Alamos	0	0	0	2	0	2	3		3	0	0	0
Luna	2	0	2	4	1	5	1	2	3	1	0	1
McKinley	11	0	11	35	17	52	7	2	9	4	0	4
Mora	1	0	1	1	0	1	2	1	3	0	0	0
Otero	7	1	8	9	5	14	9	5	14	0	0	0
Quay	1	1	2	3	0	3	0	4	4	0	0	0
Rio Arriba	10	1	11	24	4	28	25	8	33	0	0	0
Roosevelt	1	0	1	0	0	0	4	2	6	0	0	0
Sandoval	16	2	18	20	13	33	21	4	25	0	0	0
San Juan	18	3	21	36	13	49	16	14	30	4	0	4
San Miguel	0	0	0	4	1	5	8	4	12	1	0	1
Santa Fe	24	5	29	29	9	38	23	15	38	2	0	2
Sierra	1	0	1	5	0	5	2	1	3	0	0	0
Socorro	2	1	3	5	1	6	2	0	2	0	0	0
Taos	8	1	9	10	3	13	10	4	14	0	0	0
Torrance	0	0	0	5	1	6	1	2	3	1	0	1
Union	1	0	1	2	0	2	0	0	0	0	0	0
Valencia	6	5	11	7	5	12	11	2	13	0	0	0
Health Region												
Northwest	32	3	35	79	37	116	25	16	41	8	0	8
Northeast	47	7	54	75	18	93	72	34	106	3	0	3
Metro	111	16	127	142	40	182	148	87	235	12	1	13
Southeast	47	5	52	31	4	35	16	31	47	11	1	12
Southwest	43	6	49	56	8	64	38	30	68	4	1	5

See Technical Appendix for information on health regions, firearm deaths, alcohol-induced deaths, drug-induced deaths, and injury at work. The categories presented are not mutually exclusive. Injury at work deaths include all deaths (ages 15 and over) due to external causes that occurred while the decedent was working.

Mortality Section

**Table M-22 Death Rates due to Firearms, Alcohol, Drugs, and Injury at Work
by Sex, County, and Health Region
New Mexico, 2012, and United States, 2010**

	Firearms			Alcohol-Induced			Drug-Induced			Injury at Work		
	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes
United States	17.9	2.7	10.1	11.7	3.9	7.6	15.9	10.0	12.9	3.1	0.3	1.7
New Mexico	27.2	3.6	15.1	34.3	10.1	21.8	29.9	19.4	24.6	3.6	0.3	1.9
County												
Bernalillo	27.0	2.6	14.3	31.5	5.9	18.2	34.2	23.4	28.7	3.1	0.3	1.7
Catron	248.5	20.6	144.7	0.0	0.0	0.0	0.0	261.4	118.1	6.5	0.0	3.3
Chaves	54.0	0.0	26.2	19.6	0.0	9.6	9.0	37.3	23.5	0.0	0.0	0.0
Cibola	21.0	0.0	10.6	56.1	52.1	53.0	13.5	0.0	7.1	0.0	0.0	0.0
Colfax	38.7	0.0	20.1	0.0	7.9	3.9	13.0	25.2	18.2	0.0	0.0	0.0
Curry	16.9	4.4	10.7	24.4	4.4	14.1	3.5	3.6	3.6	6.9	0.0	3.6
De Baca	50.3	0.0	26.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Dona Ana	21.0	1.0	10.7	23.3	1.0	11.5	24.6	16.0	20.2	2.4	0.8	1.7
Eddy	46.4	9.2	27.7	6.0	7.0	6.5	18.6	18.7	18.7	11.2	4.1	7.8
Grant	27.7	17.2	21.9	35.3	0.0	16.9	13.3	23.8	18.7	0.0	0.0	0.0
Guadalupe	0.0	0.0	0.0	85.0	0.0	45.6	0.0	78.8	29.5	0.0	0.0	0.0
Harding	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hidalgo	0.0	0.0	0.0	96.2	0.0	48.5	0.0	0.0	0.0	28.3	0.0	14.5
Lea	15.8	3.2	9.9	25.3	3.4	14.3	8.2	10.2	9.3	10.8	0.0	5.7
Lincoln	35.9	0.0	17.2	19.1	0.0	9.0	4.8	55.4	30.7	0.0	0.0	0.0
Los Alamos	0.0	0.0	0.0	22.3	0.0	11.0	45.2	0.0	23.2	0.0	0.0	0.0
Luna	14.3	0.0	6.9	21.0	4.8	12.8	9.5	22.8	15.8	9.9	0.0	5.1
McKinley	31.8	0.0	15.4	104.6	46.8	74.7	21.7	6.9	14.3	10.2	0.0	4.9
Mora	60.6	0.0	29.7	36.2	0.0	18.8	72.3	58.2	67.3	0.0	0.0	0.0
Otero	20.2	2.5	11.6	26.2	16.1	21.1	27.8	17.6	22.7	0.0	0.0	0.0
Quay	21.8	12.1	16.7	47.1	0.0	22.9	0.0	86.6	44.4	0.0	0.0	0.0
Rio Arriba	52.0	4.4	28.5	105.0	21.1	62.2	139.8	47.3	93.8	0.0	0.0	0.0
Roosevelt	8.8	0.0	4.3	0.0	0.0	0.0	42.2	20.0	31.2	0.0	0.0	0.0
Sandoval	22.7	3.1	12.7	28.7	19.0	23.3	33.2	5.3	19.1	0.0	0.0	0.0
San Juan	29.8	5.8	17.1	57.2	21.8	39.3	25.1	22.8	24.1	6.8	0.0	3.4
San Miguel	0.0	0.0	0.0	32.9	6.1	19.4	46.8	29.0	37.8	4.6	0.0	2.3
Santa Fe	33.7	6.4	20.2	32.8	10.0	21.0	34.4	18.2	26.5	3.0	0.0	1.5
Sierra	7.0	0.0	3.6	38.1	0.0	18.8	41.2	7.7	24.2	0.0	0.0	0.0
Socorro	19.5	10.9	14.5	48.2	8.6	28.1	19.8	0.0	10.3	0.0	0.0	0.0
Taos	52.4	8.0	30.4	39.7	18.8	28.8	61.4	22.8	41.8	0.0	0.0	0.0
Torrance	0.0	0.0	0.0	38.8	17.6	27.8	10.9	25.7	17.3	15.2	0.0	8.2
Union	28.3	0.0	20.3	63.5	0.0	34.6	0.0	0.0	0.0	0.0	0.0	0.0
Valencia	16.5	12.8	14.6	15.7	13.9	14.7	29.6	5.1	17.5	0.0	0.0	0.0
Health Region												
Northwest	29.3	3.2	15.8	71.7	33.5	52.3	22.7	14.9	19.0	6.9	0.0	3.4
Northeast	33.0	4.6	19.1	41.8	11.0	26.1	51.9	24.4	38.3	2.0	0.0	1.0
Metro	25.0	3.5	13.8	29.7	8.8	18.7	33.1	19.2	26.0	2.5	0.2	1.4
Southeast	33.1	3.6	18.1	18.7	2.9	10.6	11.9	22.9	17.4	7.3	0.8	4.1
Southwest	23.2	2.8	12.8	28.4	4.1	15.8	23.1	17.1	20.0	2.4	0.4	1.4

Age-adjusted death rates are the numbers of deaths per 100,000 U.S. standard population.

See Technical Appendix for information on rates, health regions, firearm deaths, alcohol-induced deaths, drug-induced deaths, and injury at work. The categories presented are not mutually exclusive. Injury at work deaths include all deaths (ages 15 and over) due to external causes that occurred while the decedent was working. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution. See numbers in Table M-21.

Mortality Section

Table M-23 Maternal Mortality Deaths and Death Rates
New Mexico, 2005-2008 and 2009-2012

Maternal Cause of Death (ICD-10 Code) <i>10th Revision International Classification of Diseases</i>	2005-2008		2009-2012	
	(4-year) Total	Average Rate	(4-year) Total	Average Rate
Total Maternal Causes (A34, O00-O95, O98-O99)	13	10.9	19	17.1
Pregnancy with abortive outcomes (O00-O08)	1	0.8	1	0.9
Ectopic pregnancy (O00)	0	0.0	0	0.0
Spontaneous abortion (O03)	0	0.0	0	0.0
Medical abortion (O04)	1	0.8	0	0.0
Other abortion (O05)	0	0.0	0	0.0
Other & unspecified pregnancy with abortive outcomes (O01-O02, O06-O07)	0	0.0	1	0.9
Complications following abortion and ectopic and molar pregnancy (O08)	0	0.0	0	0.0
Other direct obstetric causes (A34, O10-O92)	10	8.4	14	12.6
Eclampsia and pre-eclampsia (O11, O13-O16)	0	0.0	4	3.6
Hemorrhage of pregnancy and childbirth and placenta previa (O20, O44-O46, O67, O72)	2	1.7	0	0.0
Complications predominantly related to puerperium (A34, O85-O92)	3	2.5	3	2.7
Obstetrical tetanus (A34)	0	0.0	0	0.0
Obstetrical embolism (O88)	2	1.7	3	2.7
Other complications predominantly related to the puerperium (O85-O87, O89-O92)	1	0.8	0	0.0
All other direct obstetric causes (O10, O12, O21-O43, O47-O66, O68-O71, O73-O75)	5	4.2	7	6.3
Obstetric death of unspecified cause (O95)	1	0.8	1	0.9
Indirect obstetric causes (O98-O99)	1	0.8	3	2.7

Maternal mortality rates are the numbers of maternal deaths per 100,000 live births.

See Technical Appendix for information on maternal mortality. Rate numerators are total maternal deaths for the specified 4-year period divided by 4. Rate denominators are total births for the specified 4-year period divided by 4. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Mortality Section

**Table M-24(a) Number of Infant Deaths by Sex and Race/Ethnicity
New Mexico, 2012**

Race/Ethnicity	Both Sexes	2012 Male	Female
All Races			
Infant Deaths	186	112	74
Neonatal Deaths	126	77	49
Postneonatal Deaths	60	35	25
American Indian or Alaska Native			
Infant Deaths	16	5	11
Neonatal Deaths	12	4	8
Postneonatal Deaths	4	1	3
Asian or Pacific Islander			
Infant Deaths	3	3	0
Neonatal Deaths	3	3	0
Postneonatal Deaths	0	0	0
Black or African American			
Infant Deaths	5	5	0
Neonatal Deaths	2	2	0
Postneonatal Deaths	3	3	0
Hispanic			
Infant Deaths	107	61	46
Neonatal Deaths	75	46	29
Postneonatal Deaths	32	15	17
White			
Infant Deaths	55	38	17
Neonatal Deaths	34	22	12
Postneonatal Deaths	21	16	5

See Technical Appendix for information on infant mortality and race/ethnicity.

Mortality Section

**Table M-24(b) Infant Death Rates by Sex and Race/Ethnicity
New Mexico, 2012**

Race/Ethnicity	Both Sexes	2012	
		Male	Female
All Races			
Infant Deaths	6.9	8.1	5.6
Neonatal Deaths	4.7	5.6	3.7
Postneonatal Deaths	2.2	2.5	1.9
American Indian or Alaska Native			
Infant Deaths	4.9	3.0	6.8
Neonatal Deaths	3.7	2.4	4.9
Postneonatal Deaths	1.2	0.6	1.9
Asian or Pacific Islander			
Infant Deaths	5.4	10.5	0.0
Neonatal Deaths	5.4	10.5	0.0
Postneonatal Deaths	0.0	0.0	0.0
Black or African American			
Infant Deaths	10.9	20.4	0.0
Neonatal Deaths	4.4	8.2	0.0
Postneonatal Deaths	6.5	12.2	0.0
Hispanic			
Infant Deaths	7.3	8.2	6.4
Neonatal Deaths	5.1	6.2	4.1
Postneonatal Deaths	2.2	2.0	2.4
White			
Infant Deaths	7.0	9.3	4.4
Neonatal Deaths	4.3	5.4	3.1
Postneonatal Deaths	2.7	3.9	1.3

Infant death rates are numbers of infant deaths under 1 year of age per 1,000 live births. Neonatal deaths=under 28 days. Postneonatal deaths= 28 days to under 1 year. See Technical Appendix for information on infant mortality and race/ethnicity. Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution. See numbers in Table M-24(a).

Mortality Section

**Table M-25 Number of Infant Deaths by Cause
New Mexico, 2012, and United States, 2010**

Cause of Death	ICD-10 Code -- 10th Revision	United States	New Mexico
All Causes		24,586	186
Congenital malformations, deformations and chromosomal abnormalities	Q00-Q99	5,107	48
Disorders related to short gestation and low birthweight, not classified elsewhere	P07	4,148	36
Sudden infant death syndrome	R95	2,063	0
Newborn affected by maternal complications of pregnancy	P01	1,561	9
Accidents (unintentional injuries)	V01-X59	1,110	11
Newborn affected by complications of placenta, cord, and membranes	P02	1,030	10
Bacterial sepsis of newborn	P36	583	5
Respiratory distress of newborn	P22	514	1
Diseases of the circulatory system	I00-I99	507	8
Neonatal hemorrhage	P50-P52, P54	469	2
Assault	X85-Y09, Y871	311	1
All other causes	residual	7,183	55

See Technical Appendix for information on infant mortality.

**Table M-26 Method of Final Disposition for Deaths Occurring in New Mexico
2003-2012**

Year	Total	Burial		Cremation		Removal		Other and Unspecified	
	Number	Number	Percent	Number	Percent	Number	Percent	Number	Percent
2012	16,190	5,926	36.6	8,885	54.9	611	3.8	768	4.7
2011	15,982	6,024	37.7	8,699	54.4	633	4.0	626	3.9
2010	15,511	5,997	38.7	8,320	53.6	619	4.0	575	3.7
2009	15,197	6,132	40.4	8,062	53.0	634	4.2	369	2.4
2008	15,594	6,539	41.9	8,067	51.7	635	4.1	353	2.3
2007	15,160	6,815	45.0	7,454	49.2	599	4.0	292	1.9
2006	15,044	7,021	46.7	7,169	47.7	597	4.0	257	1.7
2005	14,755	6,786	46.0	6,787	46.0	973	6.6	209	1.4
2004	14,046	6,556	44.4	6,193	42.0	1,073	7.3	224	1.5
2003	14,548	6,955	49.5	6,305	44.9	1,039	7.4	249	1.8

Removal refers to removal of remains from New Mexico to another state or country. Other and Unspecified includes donation, entombment, and all other and unspecified means of disposition.


Fetal Mortality Section



FETAL MORTALITY SECTION

A fetal death, sometimes referred to as a “stillbirth,” is defined as a death in utero. By New Mexico statute, all fetal deaths in which the fetus weighs 500 or more grams must be reported. In 2012, there were 69 reported fetal deaths, and the fetal mortality rate was 2.5 fetal deaths per 1,000 live births plus fetal deaths (Table F-1).


**Figure F-1, Fetal Mortality Rates
New Mexico, 1998-2012, United States, 1998-2006**



New Mexico’s fetal mortality rate has remained well below the United States’ rate which has ranged from 6.7 in 1998 to 6.1 fetal deaths per 1,000 live births plus fetal deaths in 2006 (Figure F-1).

Fetal Mortality Section

**Figure F-2, Fetal Mortality Rates by Mother's Age
New Mexico, 2012, and United States, 2006**



In New Mexico, for mothers under 15 years of age, there was a higher rate of fetal death than for mothers in older age groups. Except for the under 15 year age group, fetal mortality rates by mother's age category in the state were consistently below national rates. The U.S. fetal death rate was also highest among mothers under age 15 years (Figure F-2). Due to the relatively small number of fetal deaths in New Mexico in recent years, numbers of fetal deaths and rates by age can fluctuate from year to year,

From 2010-2012, the three leading causes of fetal death were: Fetus affected by complications of placenta, cord and membranes; Congenital malformation, deformation and chromosomal abnormalities; and Fetus affected by maternal complications of pregnancy. The number of unspecified causes of fetal death remained high, with 35% of fetal death reports listing this as the cause of fetal death (Table F-2).

Fetal Mortality Section

**Table F-1 Number of Fetal Deaths and Fetal Death Rates by Mother's Race/Ethnicity
New Mexico, 2012, and United States, 2006**

Mother's Race/ Ethnicity	Mother's Age													
	All Ages		Under 15 Years		15 to 19 Years		20 to 24 Years		25 to 29 Years		30 to 39 Years		40 to 49 Years	
	#	Rate	#	Rate	#	Rate	#	Rate	#	Rate	#	Rate	#	Rate
New Mexico														
All Races	69	2.5	1	19.6	8	2.4	19	2.4	14	1.9	25	3.2	2	3.4
American Indian or Alaska Native	11	3.3	0	0.0	1	2.1	3	2.8	2	2.3	5	6.3	0	0.0
Asian or Pacific Islander	2	3.6	0	0.0	0	0.0	0	0.0	0	0.0	1	3.2	1	55.6
Black or African American	2	4.3	0	0.0	0	0.0	0	0.0	1	8.3	1	7.5	0	0.0
Hispanic	42	2.9	1	27.8	5	2.3	14	3.0	9	2.3	12	3.4	1	3.7
White	12	1.5	0	0.0	2	3.6	2	1.0	2	0.8	6	2.1	0	0.0
United States														
All Races	25,972	6.1	85	13.1	3,153	7.2	6,444	5.9	6,557	5.5	8,595	5.9	1,138	10.1

See Technical Appendix for information on race/ethnicity.

Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

U.S. Data Source: CDC, National Center for Health Statistics (NCHS). 2006 U.S. data are the latest available at time of publication.

Fetal Mortality Section

**Table F-2 Number of Fetal Deaths and Fetal Death Rates by Cause
New Mexico, 2010-2012**

	2010-2012	
	Number	Rate
All fetal death causes	202	2.5
Congenital malformation, deformation and chromosomal abnormalities (Q00-Q99)	26	0.3
Fetus affected by maternal conditions that may be unrelated to present pregnancy (P00)	4	0.0
- Maternal hypertensive disorders (P00.0)	4	0.0
- All other and unspecified maternal conditions unrelated to present pregnancy (P00.1 - P00.9)	0	0.0
Fetus affected by maternal complications of pregnancy (P01)	9	0.1
Fetus affected by complications of placenta, cord and membranes (P02)	51	0.6
Fetus affected by other complications of labor and delivery (P03)	4	0.0
Disorders related to short gestation and low birthweight (P07)	4	0.0
Intrauterine hypoxia and birth asphyxia (P20 - P21)	4	0.0
Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0 - P70.2)	5	0.1
Fetal death of unspecified cause (P95)	70	0.9
All other causes	25	0.3

Rates based on fewer than 20 events may be statistically unreliable and should be interpreted with caution.

Intentionally Left Blank


Abortion Section



ABORTION SECTION

In New Mexico, reporting of legal induced abortion became law in 1977, and 1978 was the first full year of reporting. The number of abortions in New Mexico in 2012 was 3,134. This is a 16.7% decrease from the 2011 number (Table A-1).


**Figure A-1, Percent Distribution of Induced Abortions by Weeks Gestation
New Mexico Occurrence, 2012
(New Mexico Residents Only)**



In 2012, 70.4% of abortions in New Mexico were to women who were less than nine weeks pregnant. Also, 86.7% of the abortions in the state occurred within the first twelve weeks of pregnancy (Figure A-1).

Abortion Section

**Figure A-2, Ratio of Induced Abortions
New Mexico Occurrence, 2004-2012
(New Mexico Residents Only)**



The abortion ratio is the number of abortions reported for every 1,000 live births. From 2004 to 2012, New Mexico’s abortion ratio ranged from the highest value of 200.8 to the lowest value of 122.4 (Figure A-2). The national ratio for 2010, the most recent year available, was 228 abortions per 1,000 live births. New Mexico’s 2012 ratio was 46.3% lower than the national ratio. Since 2004, New Mexico’s ratio has decreased by 39%.

For all racial/ethnic groups in 2012, 20-29 year-olds accounted for over half of abortions (54.2%) in New Mexico (Table A-1).

Abortion Section

**Table A-1 Number and Percent of Induced Abortions by Woman's Age and Race/Ethnicity
New Mexico Residents (New Mexico Occurrence), 2008-2012, and United States, 2008-2010**

Year	All Ages		Under 15 Years		15 to 19 Years		20 to 24 Years		25 to 29 Years		30 to 34 Years		35 to 39 Years		40 Plus Years		Unknown Age	
	#	%	#	%	#	%	#	%	#	%	#	%	#	%	#	%	#	%
United States - All Races																		
2010	677,227	99.5	3,155	0.5	98,698	14.6	222,656	32.9	166,040	24.5	103,601	15.3	60,082	8.9	22,995	3.4	-	-
2009	690,419	99.4	3,377	0.5	107,325	15.5	225,585	32.7	168,758	24.4	101,818	14.7	60,993	8.8	22,541	3.3	-	-
2008	728,604	99.3	3,721	0.5	117,725	16.2	238,645	32.8	177,517	24.4	104,447	14.3	64,085	8.8	22,464	3.1	-	-
New Mexico - All Races																		
2012	3,134	100.0	11	0.4	457	14.6	965	30.8	733	23.4	464	14.8	234	7.5	89	2.8	181	5.8
2011	3,761	100.0	24	0.6	533	14.2	1,182	31.4	809	21.5	513	13.6	259	6.9	133	3.5	308	8.2
2010	4,478	100.0	19	0.4	774	17.3	1,495	33.4	1,014	22.6	624	13.9	331	7.4	136	3.0	85	1.9
2009	4,771	100.0	37	0.8	863	18.1	1,595	33.4	1,111	23.3	597	12.5	340	7.1	137	2.9	91	1.9
2008	5,136	100.0	32	0.6	932	18.1	1,833	35.7	1,147	22.3	621	12.1	335	6.5	137	2.7	99	1.9
American Indian or Alaska Native																		
2012	387	100.0	2	0.5	45	11.6	149	38.5	87	22.5	51	13.2	30	7.8	7	1.8	16	4.1
2011	463	100.0	4	0.9	63	13.6	152	32.8	100	21.6	57	12.3	46	9.9	16	3.5	25	5.4
2010	527	100.0	3	0.6	104	19.7	171	32.4	106	20.1	83	15.7	47	8.9	12	2.3	1	0.2
2009	555	100.0	3	0.5	94	16.9	175	31.5	150	27.0	74	13.3	39	7.0	18	3.2	2	0.4
2008	624	100.0	8	1.3	99	15.9	203	32.5	166	26.6	93	14.9	42	6.7	12	1.9	1	0.2
Asian or Pacific Islander																		
2012	61	100.0	0	0.0	5	8.2	11	18.0	16	26.2	13	21.3	9	14.8	3	4.9	4	6.6
2011	61	100.0	0	0.0	3	4.9	22	36.1	8	13.1	14	23.0	9	14.8	2	3.3	3	4.9
2010	72	100.0	0	0.0	7	9.7	15	20.8	13	18.1	11	15.3	15	20.8	11	15.3	0	0.0
2009	69	100.0	0	0.0	6	8.7	25	36.2	6	8.7	14	20.3	13	18.8	5	7.2	0	0.0
2008	80	100.0	0	0.0	6	7.5	18	22.5	22	27.5	16	20.0	13	16.3	5	6.3	0	0.0
Black or African American																		
2012	83	100.0	1	1.2	22	26.5	19	22.9	16	19.3	13	15.7	3	3.6	4	4.8	5	6.0
2011	99	100.0	0	0.0	10	10.1	32	32.3	22	22.2	17	17.2	7	7.1	0	0.0	11	11.1
2010	125	100.0	2	1.6	24	19.2	27	21.6	40	32.0	16	12.8	12	9.6	3	2.4	1	0.8
2009	108	100.0	2	1.9	22	20.4	37	34.3	21	19.4	22	20.4	2	1.9	2	1.9	0	0.0
2008	142	100.0	1	0.7	27	19.0	50	35.2	35	24.6	17	12.0	9	6.3	0	0.0	3	2.1
Hispanic																		
2012	1,643	100.0	6	0.4	284	17.3	524	31.9	384	23.4	232	14.1	108	6.6	28	1.7	77	4.7
2011	1,862	100.0	14	0.8	308	16.5	618	33.2	415	22.3	240	12.9	98	5.3	48	2.6	121	6.5
2010	2,303	100.0	9	0.4	437	19.0	839	36.4	517	22.4	315	13.7	129	5.6	48	2.1	9	0.4
2009	2,463	100.0	21	0.9	483	19.6	867	35.2	565	22.9	306	12.4	159	6.5	50	2.0	12	0.5
2008	2,667	100.0	18	0.7	536	20.1	1,020	38.2	596	22.3	292	10.9	142	5.3	48	1.8	15	0.6
White																		
2012	771	100.0	1	0.1	80	10.4	220	28.5	190	24.6	123	16.0	68	8.8	42	5.4	47	6.1
2011	1,005	100.0	5	0.5	112	11.1	292	29.1	220	21.9	150	14.9	83	8.3	63	6.3	80	8.0
2010	1,077	100.0	2	0.2	156	14.5	351	32.6	257	23.9	158	14.7	99	9.2	50	4.6	4	0.4
2009	1,201	100.0	7	0.6	203	16.9	392	32.6	298	24.8	142	11.8	105	8.7	51	4.2	3	0.2
2008	1,318	100.0	3	0.2	226	17.1	454	34.4	270	20.5	178	13.5	118	9.0	66	5.0	3	0.2

See Technical Appendix for information on race/ethnicity and induced abortions. Unknown and Other race are included in All Races.

Due to rounding percents may not add to 100.

2008--2010 United States, All Races, Known Age: U. S. abortion distribution by age is based on women of known ages.

U.S. Data Source: CDC, Abortion Surveillance Reports. U.S. 2010 data are latest available at publication time.

Technical Appendix



TECHNICAL APPENDIX

DATA SOURCES

Birth Data

Nativity data are derived from items reported on the birth certificate and include demographic information on the mother and father; geographic information on place of birth and mother's residence; information on medical risk factors, labor, and delivery; and newborn health. Most birth certificate records are transmitted electronically from hospitals. Some paper birth certificates are completed by midwives and birthing centers.

To reflect the health status of New Mexicans, most of the data in this report are presented by New Mexico residence, rather than by occurrence. To facilitate reporting by residence, states, territories and other jurisdictions engage in an inter-jurisdictional exchange process. When births to New Mexico resident mothers occur out of state, abstracts or copies of the birth certificate are transmitted to New Mexico's Bureau of Vital Records and Health Statistics (NMVRHS) for statistical reporting.

Death Data

Mortality statistics are based on items reported on the death certificate and include demographic, geographic, injury, medical, and cause-of-death information on the decedent. In most cases, funeral directors work with an informant for the decedent (usually a relative or friend) to collect demographic and geographic information. The medical and cause-of-death section of the death certificate is completed by the attending physician or the Office of the Medical Investigator (OMI). Except for deaths occurring on tribal or military lands, the Office of the Medical Investigator has jurisdiction in determining cause-of-death for all unexpected and unattended deaths in New Mexico. When deaths to New Mexico residents occur out of state, information from the death certificate (or a shortened version of the death certificate) is transmitted to NMVRSH for statistical reporting.

Fetal Death Data

Fetal death information is obtained from the New Mexico Report of Fetal Death. A fetal death, sometimes referred to as a "stillbirth," is defined as a death in utero. Until 1980, New Mexico statute had required reporting of all fetal deaths of 20 weeks or more gestation. As of 1980, the statute requires that such deaths be reported if the fetus weighs 500 grams or more, regardless of the length of gestation. The fetal death report contains much of the same information as the birth certificate plus information on the cause of fetal death. States vary in their requirements for reporting fetal deaths. The majority of states require reporting of a fetal death if the fetus is delivered at 20 weeks or more gestation. Starting January 1, 2014 there will be new reporting requirements for fetal death certificate due to a change in the law. The new statute will change to require the reporting requirements of fetal deaths from fetus weights of 500 grams or more to 20 weeks gestation or 350 grams.

Abortion Data

Induced Terminations of Pregnancy (abortions) are reported to NMVRHS by medical providers. Only limited data are collected on the Report of Induced Termination of Pregnancy. The identities of the patient and provider are not collected. Because receipt of abortion data from other states is incomplete, abortion statistics presented in this report are for New Mexico residents who had an abortion in New Mexico.

Statistical File Timeline

To allow sufficient time to receive as many vital records as possible, including those that occur out of state, NMVRHS keeps the statistical files open for approximately 4 to 6 months after the end of the calendar data year. At that time, the statistical file is closed so that statistics for that year can be consistently generated in subsequent years.

THE NEW MEXICO REVISED BIRTH CERTIFICATE

Beginning January 1, 2008, New Mexico adopted a new birth certificate based on the 2003 U.S. standard certificate of live birth. Before 2008, the New Mexico birth certificate was based on the 1989 U.S. standard.

The 2003 revision of the U.S. standard certificate of live birth introduced a number of changes to data content and quality. A number of items (e.g. maternal age, birth order, marital status, attendant at birth, birthweight, gestational age) are common to both the old and new certificates and are considered compatible. However, several other items (educational attainment, tobacco use during pregnancy, month prenatal care began, and type of vaginal or cesarean delivery), although collected on both certificate revisions, were modified and are not considered comparable. For this reason, care should be taken when comparing numbers and rates related to those items between those stated for years 2008 and after and those stated for years 2007 and prior. For more information, see the Special Feature, *The Revised Birth Certificate: Changes to the New Mexico Birth File*, in the *New Mexico Selected Health Statistics, Annual Report, 2008* at http://www.vitalrecordsnm.org/reports/2008_AR.pdf.

In addition to the information collected for the U.S. standard certificate, New Mexico collects and reports on some state-specific birth information.

New Mexico adopted the 2003 revision of the United States Standard Death Certificate, with the addition of some state-specific items, in 2006.

National Vital Statistics Data

National vital statistics data are produced by the United States Centers for Disease Control's National Center for Health Statistics (NCHS). As part of the national vital statistics system, states, territories, and other jurisdictions provide birth, death, and fetal death data to NCHS.

Population Data

Population estimates used as the denominators in calculating birth and death rates in this report were produced by the University of New Mexico's Geospatial and Population Studies (GPS) Program for years 2000 to 2012, and by the United States Census Bureau, in collaboration with NCHS, for previous years. National vital statistics data use population estimates and census counts produced by the Census Bureau.

The population numbers and birth and death rates reflect the estimates in GPS' release, vintage September 2013, in which GPS recalculated its inter-censal county-level estimates for the years 2001-2010. Updated revisions of population estimates are reflected in that birth and death rate trends presented in this report differ from those in previous annual reports.

CALCULATIONS

For the figures and tables shown in this report, the numerators of the rates are events occurring to New Mexico residents, unless otherwise specified. The denominators are the resident population figures, including all races and both sexes, unless otherwise specified. Many of the calculations in this report were accessed through New Mexico's Indicator-Based Information System (NM-IBIS) through online dataset queries (see <https://ibis.health.state.nm.us/home/ContentUsage.html>).

Rates and Ratios for Natality

This report uses crude birth rates (also termed birth rates), age-specific birth rates, fertility rates, and ratios to measure natality. Birth rates, fertility rates, and ratios in this report are per 1,000 population.

Crude Birth Rate

Crude birth rates, called birth rates in this report, are the easiest to understand and are computed by dividing the number of births in a given year by the total population (including both males and females), and multiplying by 1,000. Since most of the rates in this report are calculated by residence, the New Mexico crude birth rate is the number of births to New Mexico residents divided by the population of New Mexico and multiplied by 1,000. Similar rates are calculated by county, region, or other specified area.

Age-specific Birth Rate

Age-specific birth rates limit the rate to females in a specified age group. These rates are calculated by dividing the number of births to females in a specific age group by the number of females in that age group and multiplying by 1,000. A teen birth rate, age 15-19, is an example of an age-specific birth rate.

Fertility Rate

While the crude birth rate measures the number of births to the total population, the fertility rate limits the rate to women of child-bearing age. NMVRHS uses the NCHS fertility rate definition. The fertility rate, which is also called the general fertility rate, is the number of births to mothers of all ages divided by the numbers of females in the 15-44 age group, multiplied by 1,000.

Ratio

A rate measures the risk of an event happening in a specified time period by comparing a subset of a group to the larger group. A ratio compares one group to another group. An example of a ratio is the sex ratio, which compares the number of males to the number of females. Like a rate, a ratio may be multiplied by a constant, such as 1,000.

Rates for Fetal Mortality

A fetal death or "stillbirth" is defined as a death in utero. Fetal mortality rates are calculated by dividing the number of fetal deaths by the sum of the number of live births and fetal deaths and then multiplying by 1,000 or 100,000.

Abortion Ratios

Abortion ratios, used to compare abortions to live births, are calculated by dividing the number of abortions by the number of live births for the same period and multiplying by 1,000.

Rates for Mortality

In addition to maternal mortality rates and infant mortality rates, this publication uses crude death rates, age-specific death rates, and age-adjusted death rates to measure mortality. Except for maternal and infant mortality rates, which use live births as the denominator, death rates in this report are per 100,000 population.

Crude Death Rate

Crude death rates are the easiest to understand and are computed by dividing the number of deaths by the population, and multiplying by 100,000. Most of the rates in this report are calculated by residence. The New Mexico crude death rate is the number of deaths to New Mexico residents divided by the population of New Mexico and multiplied by 100,000. Similar rates are calculated by county, region, or other specified area.

Age-specific Death Rate

Age-specific death rates limit the rate to a specific age category. For example, the age-specific death rate for 15-19 year-olds is calculated by dividing the number of deaths to people who were age 15-19 by the number of 15-19 year-olds in the population and multiplying by 100,000.

Age-adjusted Death Rate

Because crude death rates are influenced by the age composition of the population, comparisons over time or between groups may be misleading. To account for differences in population age distributions, the age-adjusted death rate is used to compare relative mortality risks between groups and over time. This rate should be viewed as an index for comparison, rather than as a direct or actual measure of mortality risk. It is calculated by weighting the age-specific death rates and summing the products. The weights represent the proportion of the population in each age group. Beginning with 1999 data, NMVRHS joined NCHS and other agencies in using the 2000 United States standard population. For more information on age adjustment, see <http://ibis.health.state.nm.us/docs/PHStatistics/statnt20.pdf>.

GENERAL NOTES

Race and Ethnicity

Following the United States standard birth and death certificates, race and Hispanic origin are collected as separate data items on New Mexico's birth and death certificates, reports of spontaneous fetal death, and reports of induced terminations of pregnancy (abortions). For birth certificates and fetal death reports, race and Hispanic origin of the infant's mother and father (when available) are provided by the mother. For death certificates, decedent's race, Hispanic origin, and tribal affiliation are provided by an informant (usually a relative or a friend) for the deceased.

Federal reports frequently present race and ethnicity (Hispanic origin) separately. Persons of Hispanic origin may be of any race. In the New Mexico Department of Health (DOH), race and ethnicity are presented as a single social and cultural construct. The categories used by DOH and used in the figures and tables in this report, are: American Indian or Alaska Native, Asian or Pacific Islander, Black or African American, Hispanic, and White.

Beginning in July 2013, DOH implemented modified guidelines for the presentation of race and ethnicity data as a single construct (Race/Ethnicity). The revised definition categorizes into the Hispanic category any individual reporting Hispanic ethnicity, whereas, previously, a person's race reported as American Indian or Alaska Native, Asian or Pacific Islander, or Black or African American, race was presented as such regardless of Hispanic origin. The revised New Mexico Department of Health Race/Ethnicity guidelines are followed in the *New Mexico Selected Health Statistics Annual Report 2012*, and historical data presented in this report have been recalculated using the new definition; therefore, differing from race and ethnicity data for those years presented in previous reports. **For more information on this new standard, please visit:** <http://ibis.health.state.nm.us/resources/RacEth2013.html>.

Geography

State Health Regions

In addition to county level data, this publication also reports data by New Mexico Health Regions. The State Health Regions include the following counties:

Northwest Region: Cibola, McKinley, and San Juan;

Technical Appendix

Northeast Region: Colfax, Guadalupe, Los Alamos, Mora, Rio Arriba, San Miguel, Santa Fe, Taos, Union, and Harding;

Metro Region: Bernalillo, Sandoval, Torrance, and Valencia;

Southeast Region: Chaves, Curry, De Baca, Eddy, Lea, Lincoln, Quay, and Roosevelt; and

Southwest Region: Catron, Doña Ana, Grant, Hidalgo, Luna, Otero, Sierra, and Socorro.

Residence Data

Residence data are presented by the place where the person normally resided, regardless of where the event occurred. Except where specified, all data presented in this report are residence data.

Occurrence Data

Occurrence data refers to the place where the event occurred, regardless of the usual residence of the person involved.

NOTES ON NATALITY DATA

Paternity

By statute, if the mother was married at the time of either conception or birth, her husband is recorded as the infant's father. If the mother is not married, the father's name and demographic information may be recorded if the mother and father sign an Acknowledgment of Paternity. If a determination of paternity has been made by a court, the name of the father as determined by the court is entered along with his demographic information. Birth records where the mother is unmarried and no acknowledgement of paternity has been signed will have missing demographic data for the father.

Birth Order

Birth order is the order in which this child (of all the children born to the mother) was born. As a fertility indicator, it is used to measure how many children a mother has.

Multiple Births

Twins and triplets are examples of multiple births. The multiple birth rate is defined as the number of twins, triplets, or higher-order multiple births per 1,000 live births. It is calculated by dividing the number of live births in multiple deliveries by the number of total live births and multiplying by 1,000.

Birthweight

In New Mexico, birthweight is reported in grams. Low birthweight infants weigh less than 2,500 grams (5 pounds, 9 ounces) at birth, while very low birthweight infants weigh less than 1,500 grams (3 pounds, 5 ounces). High birthweight infants weigh 4,000 grams (8 pounds, 14 ounces) or more at birth.

Gestational Age

The interval between the first day of mother's last normal menstrual period (LMP) and the date of birth is the primary method used to calculate gestational age. However, LMP is subject to error because of maternal recall or misidentification of the LMP due to other factors such as post-conception bleeding. Additionally, LMP may be unreported on the birth record. For records where LMP is missing or the computed gestation is inconsistent with the infant's birthweight (taking into account multiple births), the clinical estimate of gestation is used. The clinical estimate of gestation is recorded on the birth record based on what is reported in the mother's medical record by the mother's physician or other health professional. The final result is the calculated (also called combined) gestational age. To enable comparability with United States figures, NMVRHS calculates gestational age (presented in weeks) consistent with the imputation procedure of the National Center for Health Statistics

Technical Appendix

(NCHS). NCHS revised the methodology with the 2003 revision to the U.S. Standard Birth Certificate, implemented in New Mexico starting with 2008 birth data.

Measures of Prenatal Care

There are two primary ways that New Mexico reports on prenatal care: (1) the trimester prenatal care began and (2) the Kessner Index. Both rely heavily on when prenatal care was initiated.

Kessner Index

New Mexico's traditional measure of prenatal care is the modified Kessner Index. Level of prenatal care is defined using a combination of the month prenatal care began and the number of prenatal visits. Low level of care is defined as care that either begins in the third trimester, consists of less than five prenatal care visits, or no prenatal care. Mid level of care is defined as care that begins during the first trimester with five to eight total prenatal visits or care beginning in the fourth to sixth month of pregnancy with five or more visits. High (optimum) level of care is defined as care that begins during the first trimester with a total of nine or more prenatal care visits during that period.

Method of Delivery

Methods of delivery may be reported as vaginal birth, vaginal birth after previous cesarean-section, primary cesarean-section, and repeat cesarean-section.

Birth Attendant

Except in rare cases, newborns in New Mexico are delivered by the following birth attendants: Medical Doctor or Doctor of Osteopathy, Licensed Midwife, Certified Nurse Midwife, or Registered Nurse.

Definitions of midwives and nurses licensed in New Mexico are below:

Licensed Midwife (LM). A Licensed Midwife is a person who has successfully completed the prescribed course of studies in midwifery and has acquired the requisite qualifications to be registered and/or legally licensed to practice midwifery.

Certified Nurse Midwife (CNM). A Certified Nurse-Midwife (CNM) is an individual educated in the two disciplines of nursing and midwifery and who possesses evidence of certification according to the requirements of the American College of Nurse-Midwives. A certified nurse-midwife must be licensed in New Mexico as a registered nurse.

Registered Nurse (RN). A Registered Nurse (RN) is a nurse who has graduated from a formal program of nursing education (diploma school, associate degree or baccalaureate program) and is licensed by the appropriate New Mexico state authority.

NOTES ON MORTALITY DATA

Cause of Death

ICD Classification

Beginning with 1999 data, cause of death has been coded according to the tenth revision of the World Health Organization's International Classification of Diseases (ICD-10). The International Classification of Diseases (ICD) is a system of classification developed in partnership with the World Health Organization (WHO) and WHO Collaborating Centers. (The North American Collaborating Center is housed at NCHS in Hyattsville, Maryland.) WHO member nations are required to use this classification system for comparability in the collection and classification of health statistics (<http://www.who.int/classifications/icd/en/>).

Technical Appendix

New Mexico began systematic record keeping of causes of death and disease morbidity in 1929 and became part of the United States Vital Statistics System in the same year. Revisions to the ICD have occurred almost every ten years since the first version went into effect in 1900. The tenth revision is the exception, going into effect 20 years after the 1979 implementation of the ninth revision. Revisions to the ICD are necessary to keep up with advances in medical science. Changes in classification due to revision may lead to discontinuities in cause of death trends. To account for differences between revisions, comparability ratios have been applied to statistics of deaths occurring from 1978-1998 (http://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_02.pdf).

In addition to serving as a classification system, the ICD also includes coding rules that allow the nosologist (person who codes cause of death) to select the underlying cause of death - the single condition on the death certificate that is considered most informative from a public health point of view. Also included are definitions (such as "maternal death"), regulations on the compilation and publication of statistics, a prescribed format of the medical certification of death (part of the death certificate), and tabulation lists that indicate cause-of-death groupings that should be used to present comparable mortality data.

Leading Cause of Death

Causes of death are ranked following procedures that are consistent with the recommendations of the 1951 Public Health Conference on Records and Statistics, where causes are ranked by the number of deaths in each rankable cause category. With each ICD revision, the list of rankable causes has been revised. The current rankable cause list is based on the cause of death lists produced by NCHS (http://www.cdc.gov/nchs/data/dvs/im9_2002.pdf).

Alcohol-induced Deaths

Causes of death and corresponding ICD-10 codes used by NMVRHS for alcohol-induced deaths are: Alcohol-induced pseudo-Cushing's syndrome, E24.4; Mental and behavioral disorders due to alcohol use, F10; Degeneration of nervous system due to alcohol, G31.2; Alcoholic polyneuropathy, G62.1; Alcoholic myopathy, G72.1; Alcoholic cardiomyopathy, I42.6; Alcoholic gastritis, K29.2; Alcoholic liver disease, K70; Alcohol-induced chronic pancreatitis, K86.0; Finding of alcohol in blood, R78.0; Accidental poisoning by and exposure by alcohol, X45; Intentional self poisoning by and exposure to alcohol, X65; Poisoning by and exposure to alcohol, undetermined intent, Y15. Alcohol-induced causes exclude other unintentional injuries, including motor vehicle accidents; homicides; and other causes indirectly related to alcohol use. Infant deaths associated with maternal alcohol use are also excluded.

Drug-induced Deaths

This category includes Drug-induced folate deficiency anemia, D52.1; Drug-induced hemolytic anemia, D59.0; Drug-induced nonautoimmune hemolytic anemia, D59.2; Drug-induced aplastic anemia, D61.1; Secondary sideroblastic anemia due to drugs and toxins, D64.2; Drug-induced thyroiditis, E06.4; Drug-induced hypoglycemia without coma, E16.0; Drug-induced hypopituitarism, E23.1; Drug-induced Cushing's syndrome, E24.2; Drug-induced adrenocortical insufficiency, E27.3; Drug-induced obesity, E66.1; Mental and behavioral disorders due to psychoactive substance use (F11.0-F11.5, F11.7-F11.9, F12.0-F12.5, F12.7-F12.9, F13.0-F13.5, F13.7-F13.9, F14.0-F14.5, F14.7-F14.9, F15.0-F15.5, F15.7-F15.9, F16.0-F16.5, F16.7-F16.9, F17.0, F17.3-F17.5, F17.7-F17.9, F18.0-F18.5, F18.7-F18.9, F19.0-F19.5, and F19.7-F19.9); Other drug-induced secondary parkinsonism, G21.1; Drug-induced dystonia, G24.0; Drug-induced tremor, G25.1; Drug-induced chorea, G25.4; Drug-induced tics and other tics of organic origin, G25.6; Drug-induced headache, not elsewhere classified, G44.4; Drug-induced polyneuropathy, G62.0; Drug-induced myopathy, G72.0; Hypotension due to drugs, I95.2; Acute drug-induced interstitial lung disorders, J70.2; Chronic drug-induced interstitial lung disorders, J70.3; Drug-induced interstitial lung disorder, unspecified, J70.4; Drug-induced pemphigus, L10.5; Generalized skin eruption due to drugs and medicaments, L27.0; Localized skin eruption due to drugs and medicaments, L27.1; Drug-induced gout, M10.2; Drug-induced systemic lupus erythematosus, M32.0; Drug-induced osteoporosis with pathological fracture, M80.4; Drug-induced osteoporosis, M81.4; Other drug-induced osteomalacia in adults, M83.5; Osteonecrosis due to drugs, M87.1; Finding of opiate drug in blood, R78.1; Finding of cocaine in blood, R78.2; Finding of hallucinogen in blood, R78.3; Finding of other drugs of addictive potential in blood, R78.4; and

Technical Appendix

Finding of psychotropic drug in blood, R78.5; Accidental poisoning by and exposure to drugs, medicaments and biological substances, X40-X44; Intentional self-poisoning (suicide) by and exposure to drugs, medicaments and biological substances, X60-X64; Assault (homicide) by drugs, medicaments and biological substances, X85; and Poisoning by and exposure to drugs, medicaments and biological substances, undetermined intent, Y10-Y14. Other unintentional injuries, homicides, and other causes indirectly related to drug use are excluded from this category. Infant deaths due to mother's drug use are also excluded.

Firearm Deaths

Mortality due to firearm use includes suicide, homicide, unintentional injury, legal intervention and undetermined deaths. This category excludes firearm injury deaths due to explosives and other causes indirectly related to firearms. The specific causes of death and ICD-10 codes included in this category are: Accidental discharge of firearms, W32-W34; Intentional self-harm (suicide) by discharge of firearms, X72-X74; Assault (homicide) by discharge of firearms, X93-X95; Discharge of firearms, undetermined intent, Y22-Y24; and Legal intervention involving firearm discharge, Y35.0

Unintentional Injury Deaths

Unintentional injury deaths are those with ICD-10 codes V01-X59 and Y85-Y86. Categories of unintentional injury deaths are: Motor vehicle crash injuries, V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2; Fall injuries, W00-W19; Poisonings, X40-X49; and all other unintentional injuries, V01, V05-V06, V09.1, V09.3-V09.9, V10-V11, V15-V18, V19.3, V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9, V90-V99, W20-W99, X00-X39, X50-X59, Y85-Y86. The "other" category includes such injuries as accidental drowning and submersion; accidental exposure to smoke, fire, and flames; accidental firearm discharge; water, air, and space and other land transport accidents; and other/unspecified non-transport accidents.

Injury at Work

Information on deaths due to injuries sustained at work is from the injury at work check box on the death certificate. Except for deaths occurring on tribal lands or military facilities, the Office of the Medical Investigator investigates injury at work deaths that occur in New Mexico.

Maternal Mortality

Maternal deaths are defined by WHO as "the death of a woman while pregnant or within 42 days of termination of pregnancy, irrespective of the duration and the site of the pregnancy, from any cause related to or aggravated by the pregnancy or its management, but not from accidental or incidental causes." ICD-10 codes used to capture maternal deaths are: A34, O00-O95, and O98-O99.

Maternal mortality rates are calculated by dividing the number of deaths due to maternal causes by the number of live births for the same period and multiplying by 100,000. Maternal mortality rates are presented as the number of maternal deaths per 100,000 live births. Because only pregnant women are at risk of maternal mortality, live births are used as the denominator to approximate the population of pregnant women.

Infant Mortality

Infant mortality is defined as the death of an infant under one year of age and is often separated into two age groups: neonatal and postneonatal. The neonatal period represents infants less than 28 days old. Postneonatal infants are at least 28 days of age but less than one year of age. The infant mortality rate is one of the most widely used health indicators and is computed by dividing the number of infant deaths by the number of live births in a period and multiplying by a constant (1,000 or 100,000). Neonatal and postneonatal mortality rates also use the total number of live births as the denominator. Another measure of infant mortality is the infant death rate, which uses the population of infants as the denominator rather than the number of live births. In this report, only infant mortality rates are used.

Technical Appendix

Infant mortality rates by race/ethnicity are calculated by using the decedent's (the infant's) race/ethnicity reported on the death certificate and the mother's race/ethnicity reported on the birth certificate

Intentionally Left Blank

The Bureau of Vital Records and Health Statistics is located in the Public Health Nurse Memorial Bldg.
1105 St. Francis Drive
Santa Fe, NM 87505

For information on obtaining New Mexico birth and death certificates, please visit us on the web at www.vitalrecordsnm.org or call 866-534-0051.



ADMINISTRATION

Mark Kassouf, Bureau Chief and Director, State Center for Health Statistics
Liz Padilla, Financial Specialist
Lisa Sandoval, Administrative Secretary

STATISTICS AND EPIDEMIOLOGY

Camille Clifford, Epidemiologist
Irma Montoya, Management Analyst
Tony Ortiz, Epidemiologist
Rosemary Perez-Mendoza, Statistician
Anne Worthington, Epidemiologist

BIRTH AND DEATH REGISTRATION

Renee Valencia, New Mexico Acting State Registrar and Registration Manager,
Irene Chavez, Office Clerk
Martha Lucero, Office Clerk
Melinda Roybal, Statistician
Peter Varela, Statistician

VITAL RECORDS

Vacant, Records Unit Manager
Hector Cruz, Records Unit Supervisor
Rita Encinias, Customer Service Representative Unit Supervisor
Percy Armijo, Office Clerk
Peggy Armijo, Office Clerk
Amber Beauchamp, Office Clerk
George Dominguez, Office Clerk
Nancy Blake, Training and Development
Genoveva Garcia, Office Clerk
Natasha Krohn, Office Clerk
Debbie Laemmle, Customer Service Representative Unit
Matthew Medina, Clerk Specialist
Michelle Montoya-Griego, Office Clerk
Monika Romero, Deposit/Audit & Security Paper Management
Rebecca Ulibarri, Order Processing & Permits
Valerie Voight, Sanchez, Office Clerk

