VALENCIA COUNTY HEALTH PROFILE

2003 NEW MEXICO COUNTY HEALTH PROFILES NEW MEXICO DEPARTMENT OF HEALTH PUBLIC HEALTH DIVISION OFFICE OF NEW MEXICO VITAL RECORDS AND HEALTH STATISTICS

Welcome to the fifth update of the New Mexico County Health Profiles. This publication is dedicated to users of county-level data as well as to those who contributed data, suggestions, interpretations, and comments. Thanks to so many of you, there are many improvements in this year's version. Some of you work in the Department of Health, but a great many of you work in other agencies and it is gratifying to be able to have collaborated with you.

As in prior versions, the latest data available from each source is used. Questions about the data, their collection, limitations, or interpretation should be directed to the appropriate data source. To aid in this, a source is listed at the bottom of each table throughout the profile.

The compilation of the Profiles is possible because of assistance and contributions from many agencies and data providers: NM Department of Health, District Health Offices, Diabetes Prevention and Control Program, Family Food and Nutrition Services, Family Health Bureau, Infectious Disease Bureau, Injury Prevention and EMS Bureau, Long Term Services Division, Family Infant Toddler Program, Behavioral Health, and Office of Epidemiology; NM Children, Youth and Families Department, Child Protective Services; NM Commission on Higher Education; NM Department of Education; NM Department of Labor; NM Health Policy Commission; NM Department of Transportation, Traffic Safety Bureau; NM Human Services Department, Income Support Division, Medical Assistance Division; University of New Mexico's Bureau of Business and Economic Research (BBER) and New Mexico Tumor Registry; Center for Disease Control's National Center for Health Statistics; US Department of Commerce, Bureau of the Census and Bureau of Economic Analysis; and US Bureau of Labor Statistics.

Without outstanding assistance from the entire staff at the Office of New Mexico Vital Records and Health Statistics, these profiles would not have been produced and we especially wish to thank: Mary M. Gonzales, John Hedrick, Brad Whorton, Brian Woods, Radha S. Biswas, Tony Ortiz, Renee Valencia, Michael Patterson, Veronica Tapia, Pete Varela, Rebecca Rodriguez, and Melinda Roybal.

Your comments are welcome. You may call 827-0167 to provide comments or suggestions for future reports.

Celine Sanchez
State Registrar and Acting Chief
Office of New Mexico Vital Records and Health Statistics

Kimberley Peters

Director, Designated State Center for Health Statistics

Mary Houdek
Manager, Information Systems and GIS

Mary Anne Escudero

Manager, Fraud Program & Deputy State Registrar

Copyright 2003 Office of New Mexico Vital Records and Health Statistics. Permission to quote from or reproduce materials from this report is granted with attributions.



Office of New Mexico Vital Records & Health Statistics County Health Profile Team

Mary M. Gonzales Management Analyst & Co-editor

Jonathan Hedrick Statistician & Co-editor

Radha S. Biswas Epidemiologist
Tony Ortiz Epidemiologist
Michael Patterson Statistician
Rebecca Rodriguez Statistician
Melinda Roybal Statistician
Veronica Tapia Clerk Specialist


Renee Valencia
Pete Varela
Brad Whorton
Brian Woods
Statistician
Epidemiologist
Epidemiologist

TABLE OF CONTENTS


| Population5 |
|---|
| Growth and Change |
| Distribution by Age and Gender |
| Distribution by Race/Ethnicity |
| Distribution by Hispanic Ethnicity |
| Special Populations |
| Special i opulations |
| Socio-Economic |
| Per Capita Income / Average Wages / Total Personal Income |
| |
| Unemployment / TANF & Food Stamps / Medicaid / Poverty |
| Actornal and Infant Health |
| Maternal and Infant Health 10 |
| Live Births / Crude Birth Rates / Fertility Rates |
| Number of Births and Birth Rates by Age of Mother |
| Percent of Total Births to Teen Mothers / Single Mothers |
| Percent of Births to Mothers with < 12 Years Education / First Trimester Prenatal Care |
| Prenatal Care – Modified Kessner Index |
| Adequacy of Prenatal Care Utilization - Kotelchuck Index |
| Live Births by Type of Attendant |
| Low Birth Weight Births / Very Low Birth Weight Births / High Birth Weight |
| Supplemental Programs / Births Paid by Medicaid / Female Population Age 15-44 |
| Number Infant, Neonatal and Postneonatal Deaths / Infant Mortality - SIDS Deaths/Birth Defects |
| Trainer main, Treonaur and Testines main Penalty (1928) Penalty (1928) |
| |
| Thild Health (1-14) Vears of Age |
| Child Health (1-14) Years of Age. 20 Children's Chronic Conditions Registry Data / Child Abuse |
| Children's Chronic Conditions Registry Data / Child Abuse |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age. 24 |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age. 24 |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |
| Children's Chronic Conditions Registry Data / Child Abuse Family Infant Toddler (FIT) Data School Enrollment Mortality – Deaths / Death Rates / Childhood Leading Causes Adolescent / Youth (15-19) Years of Age |

TABLE OF CONTENTS cont'd

..........




| | 2002 POPULATION ESTIMATES BY AGE AND GENDER | | | | | | | | | | | |
|----------------|--|-------------|-------------|------------|---------|-----------|--|--|--|--|--|--|
| Age Groups | Val | lencia Cou | nty | New Mexico | | | | | | | | |
| rige Groups | Male | Female | Both | Male | Female | Both | | | | | | |
| Less than 1 yr | 436 | 481 | 917 | 13,746 | 13,361 | 27,107 | | | | | | |
| 1 to 4 years | 1,964 | 1,943 | 3,907 | 53,236 | 51,780 | 105,016 | | | | | | |
| 5 to 9 years | 2,648 | 2,563 | 5,211 | 67,768 | 65,647 | 133,415 | | | | | | |
| 10 to 14 years | 3,049 | 2,896 | 5,945 | 74,247 | 71,746 | 145,993 | | | | | | |
| 15 to 19 years | 2,850 | 2,736 | 5,586 | 76,356 | 72,288 | 148,644 | | | | | | |
| 20 to 24 years | 2,177 | 2,005 | 4,182 | 67,182 | 64,847 | 132,029 | | | | | | |
| 25 to 29 years | 1,884 | 1,806 | 3,690 | 56,796 | 56,273 | 113,069 | | | | | | |
| 30 to 34 years | 2,270 | 2,241 | 4,511 | 59,596 | 59,961 | 119,557 | | | | | | |
| 35 to 39 years | 2,640 | 2,612 | 5,252 | 64,222 | 65,623 | 129,845 | | | | | | |
| 40 to 44 years | 2,808 | 2,783 | 5,591 | 70,951 | 73,496 | 144,447 | | | | | | |
| 45 to 49 years | 2,480 | 2,464 | 4,944 | 66,325 | 70,011 | 136,336 | | | | | | |
| 50 to 54 years | 2,161 | 2,200 | 4,361 | 60,317 | 63,383 | 123,700 | | | | | | |
| 55 to 59 years | 1,746 | 1,763 | 3,509 | 47,075 | 50,411 | 97,486 | | | | | | |
| 60 to 64 years | 1,342 | 1,431 | 2,773 | 36,530 | 40,431 | 76,961 | | | | | | |
| 65 to 69 years | 1,117 | 1,171 | 2,288 | 30,979 | 34,231 | 65,210 | | | | | | |
| 70 to 74 years | 819 | 987 | 1,806 | 26,175 | 29,427 | 55,602 | | | | | | |
| 75 to 79 years | 615 | 803 | 1,418 | 19,298 | 25,786 | 45,084 | | | | | | |
| 80 to 84 years | 430 | 485 | 915 | 12,434 | 17,304 | 29,738 | | | | | | |
| 85 and over | 272 | 500 | 772 | 8,926 | 16,894 | 25,820 | | | | | | |
| All Ages | 33,708 | 33,870 | 67,578 | 912,159 | 942,900 | 1,855,059 | | | | | | |
| | Sour | ce: U.S. Bu | reau of the | Census | | | | | | | | |




| | 2002 POPULATION ESTIMATES BY AGE AND RACE/ETHNICITY | | | | | | | | | | | |
|----------------|---|-----------|--|--------------------------------|-------------------------|---------------------|--|--|--|--|--|--|
| Age Groups | White | ONE Black | RACE American Indian / Alaska Native | Asian / Pacific Islander | Two or More Races | Total Population | Hispanic or Latino Ethnicity Any Race ^{/1} | | | | | |
| Under 5 years | 4,461 | 64 | 166 | 35 | 98 | 4,824 | 3,078 | | | | | |
| 5 to 9 years | 4,774 | 76 | 212 | 43 | 106 | 5,211 | 3,466 | | | | | |
| 10 to 14 years | 5,392 | 93 | 294 | 41 | 125 | 5,945 | 3,795 | | | | | |
| 15 to 19 years | 5,119 | 93 | 228 | 48 | 98 | 5,586 | 3,545 | | | | | |
| 20 to 24 years | 3,834 | 66 | 175 | 47 | 60 | 4,182 | 2,657 | | | | | |
| 25 to 29 years | 3,426 | 63 | 128 | 36 | 37 | 3,690 | 2,308 | | | | | |
| 30 to 34 years | 4,148 | 85 | 201 | 31 | 46 | 4,511 | 2,612 | | | | | |
| 35 to 39 years | 4,873 | 84 | 203 | 39 | 53 | 5,252 | 3,097 | | | | | |
| 40 to 44 years | 5,174 | 85 | 235 | 42 | 55 | 5,591 | 2,900 | | | | | |
| 45 to 49 years | 4,621 | 76 | 176 | 29 | 42 | 4,944 | 2,359 | | | | | |
| 50 to 54 years | 4,091 | 65 | 133 | 37 | 35 | 4,361 | 1,983 | | | | | |
| 55 to 59 years | 3,303 | 50 | 106 | 22 | 28 | 3,509 | 1,569 | | | | | |
| 60 to 64 years | 2,638 | 23 | 77 | 11 | 24 | 2,773 | 1,215 | | | | | |
| 65 to 69 years | 2,169 | 27 | 60 | 13 | 19 | 2,288 | 982 | | | | | |
| 70 to 74 years | 1,730 | 14 | 31 | 19 | 12 | 1,806 | 711 | | | | | |
| 75 to 79 years | 1,362 | 23 | 17 | 7 | 9 | 1,418 | 526 | | | | | |
| 80 to 84 years | 871 | 18 | 17 | 4 | 5 | 915 | 309 | | | | | |
| 85 and over | 746 | 5 | 12 | 4 | 5 | 772 | 267 | | | | | |
| All Ages | 62,732 | 1,010 | 2,471 | 508 | 857 | 67,578 | 37,379 | | | | | |

Hispanic or Latino origin may be of any race. Adding the Hispanic or Latino population to the racial categories will result in double counting.


SOURCE: U.S. Bureau of the Census




| UNDER 20 POPULATION 2000, 2001 and 2002 | | | | | | | | | | |
|--|--------------|--------------|--------|--|--|--|--|--|--|--|
| Age Groups 2000 2001 2002 | | | | | | | | | | |
| Under 5 years | 5,050 | 4,914 | 4,824 | | | | | | | |
| 5 to 9 years | 5,608 | 5,407 | 5,211 | | | | | | | |
| 10 to 14 years | 5,936 | 5,972 | 5,945 | | | | | | | |
| 15 to 19 years | 5,151 | 5,410 | 5,586 | | | | | | | |
| Under 20 years | 21,745 | 21,703 | 21,566 | | | | | | | |
| County Population 66,152 66,851 67,578 | | | | | | | | | | |
| Source: U | .S. Bureau o | of the Censu | S | | | | | | | |





| 20 TO 64 POPULATION 2000, 2001 and 2002 | | | | | | | | | | |
|--|--------|--------|--------|--|--|--|--|--|--|--|
| Age Groups | 2000 | 2001 | 2002 | | | | | | | |
| 20 to 24 years | 3,719 | 3,920 | 4,182 | | | | | | | |
| 25 to 29 years | 3,993 | 3,764 | 3,690 | | | | | | | |
| 30 to 34 years | 4,781 | 4,648 | 4,511 | | | | | | | |
| 35 to 39 years | 5,472 | 5,308 | 5,252 | | | | | | | |
| 40 to 44 years | 5,352 | 5,473 | 5,591 | | | | | | | |
| 45 to 49 years | 4,599 | 4,841 | 4,944 | | | | | | | |
| 50 to 54 years | 4,100 | 4,268 | 4,361 | | | | | | | |
| 55 to 59 years | 3,110 | 3,261 | 3,509 | | | | | | | |
| 60 to 64 years | 2,558 | 2,675 | 2,773 | | | | | | | |
| 20 to 64 years | 37,684 | 38,158 | 38,813 | | | | | | | |
| County Population | 66,152 | 66,851 | 67,578 | | | | | | | |



| | 65 AND OLDER POPULATION 2000, 2001 and 2002 | | | | | | | | | | |
|--------------------|--|--------------|--------|--|--|--|--|--|--|--|--|
| Age Groups | 2000 | 2001 | 2002 | | | | | | | | |
| 65 to 69 years | 2,195 | 2,258 | 2,288 | | | | | | | | |
| 70 to 74 years | 1,731 | 1,778 | 1,806 | | | | | | | | |
| 75 to 79 years | 1,322 | 1,369 | 1,418 | | | | | | | | |
| 80 to 84 years | 838 | 873 | 915 | | | | | | | | |
| 85 and over | 637 | 712 | 772 | | | | | | | | |
| 65 years and older | 6,723 | 6,990 | 7,199 | | | | | | | | |
| County Population | 66,152 | 66,851 | 67,578 | | | | | | | | |
| Source: U | J.S. Bureau o | of the Censu | I S | | | | | | | | |








r/ Revised p/ Preliminary

SOURCE: U.S. Dept of Commerce, Bureau of Economic Analysis, released December 2002.



r/ Revised p/ Preliminary

See glossary for definition of Total Personal Income

SOURCE: U.S. Dept of Commerce, Bureau of Economic Analysis, released May 2003.

^{1/} The employment estimates used to compute the average wage are a job, not person, count.

SOCIO-ECONOMIC - UNEMPLOYMENT / TANF / FOOD STAMPS / MEDICAID / POVERTY




AVERAGE MONTHYLY TANF AND FOOD STAMPS CASES FISCAL YEAR 2003 (7/02 - 6/03)

| Supplement | Valend | cia Co. | New Mexico | | | |
|-------------|--------|----------|------------|----------|--|--|
| Supprement | Number | Percent* | Number | Percent* | | |
| TANF | 838 | 1.2 | 16,382 | 0.9 | | |
| Food Stamps | 3,471 | 5.1 | 72,411 | 3.9 | | |

^{*}Percent of county's estimated 2002 population.

NOTE: Starting July 1, 1999, changes were made in the methodology for counting TANF (Temporary Assistance for Needy Families) caseloads. Caution should be used when comparing data to previous fiscal years.

SOURCE: NM Human Services Department, Income Support Division.




Income Support Division

ESTIMATED NUMBER AND PERCENT OF PEOPLE IN POVERTY

See glossary for definition of poverty. *Percent of Specified Age Group.

SOURCE: U.S. Census Bureau, "Estimates Model 1999."

AVERAGE MONTHLY PERCENT TANF AND FOOD STAMP CASES FISCAL YEAR 2003





TANF = Temporary Assistance for Needy Families.

SOURCE: NM Human Services Department Income Support Division.

¹⁹⁹⁹ Valencia Co. New Mexico Age Number Percent* Number Percent* 11,220 326,223 18.2 All Ages in Poverty 17.3 -Children age < 18 years 5,028 25.5 133,194 26.4 -Related children ages 5-17 years 3,341 23.1 92,333 25.0

| | RESIDENT LIVE BIRTHS BY COUNTY OF RESIDENCE AND STATE 1992-2002 | | | | | | | | | | | | |
|--------------|---|-----------|-----------|-----------|-----------|------------|----------|-----------|----------|-----------|--------|--|--|
| | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* | | |
| Valencia Co. | 788 | 769 | 818 | 857 | 965 | 969 | 964 | 1,000 | 992 | 1,000 | 952 | | |
| New Mexico | 27,910 | 27,831 | 27,585 | 26,914 | 27,216 | 26,844 | 27,294 | 27,133 | 27,206 | 27,101 | 27,708 | | |
| | *2002 data are provisional | | | | | | | | | | | | |
| SOUR | RCE: NM I | De partme | nt of Hea | lth, Offi | ce of Nev | v Me xi co | Vital Re | cords and | Health S | tatistics | | | |





| VALENCIA COUNTY RESIDENT LIVE BIRTHS BY AGE OF MOTHER 1992-2002 | | | | | | | | | | | |
|---|--------------|-------|------|------|------|------|------|------|------|------|-------|
| AGE | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* |
| 15-19 | 149 | 138 | 164 | 170 | 207 | 176 | 174 | 207 | 184 | 181 | 174 |
| 20-24 | 227 | 229 | 233 | 254 | 242 | 291 | 295 | 291 | 296 | 321 | 281 |
| 25-29 | 219 | 186 | 206 | 215 | 250 | 252 | 250 | 239 | 246 | 255 | 251 |
| 30-34 | 126 | 140 | 145 | 144 | 179 | 165 | 147 | 172 | 173 | 153 | 143 |
| 35-39 | 59 | 54 | 55 | 64 | 70 | 66 | 76 | 66 | 63 | 71 | 77 |
| 40-44 | 5 | 14 | 10 | 4 | 14 | 10 | 18 | 14 | 26 | 13 | 21 |
| *2002 data | a are provis | ional | | | | | | | | | |

SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics

| | VALENCIA COUNTY RESIDENT BIRTH RATES** BY AGE OF MOTHER 1992-2002 | | | | | | | | | | | |
|------------|---|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--|
| AGE | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* | |
| 15-19 | 84.2 | 73.3 | 81.6 | 77.0 | 87.6 | 70.3 | 66.9 | 78.3 | 73.6 | 68.5 | 63.6 | |
| 20-24 | 172.9 | 166.7 | 160.8 | 162.2 | 147.1 | 167.7 | 164.8 | 159.1 | 163.2 | 167.6 | 140.1 | |
| 25-29 | 136.4 | 118.7 | 132.7 | 134.7 | 151.9 | 148.9 | 143.3 | 134.1 | 125.1 | 139.9 | 139.0 | |
| 30-34 | 59.4 | 64.3 | 65.1 | 62.7 | 78.4 | 73.4 | 67.6 | 81.5 | 73.0 | 66.5 | 63.8 | |
| 35-39 | 29.6 | 25.7 | 24.9 | 26.8 | 27.9 | 25.4 | 28.8 | 24.9 | 23.4 | 27.5 | 29.5 | |
| 40-44 | 2.8 | 7.3 | 4.9 | 1.8 | 5.9 | 4.0 | 6.9 | 5.2 | 9.8 | 4.7 | 7.5 | |
| *2002 data | **Rate per 1,000 women in specified group | | | | | | | | | | | |

SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics

| | NEW MEXICO RESIDENT BIRTH RATES** BY AGE OF MOTHER 1992-2002 | | | | | | | | | | |
|------------|--|-------|-----------|-------------|-------------|--------------|-------|-------|-------|-------|-------|
| AGE | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* |
| 15-19 | 79.8 | 80.7 | 77.2 | 74.3 | 70.8 | 68.0 | 69.1 | 67.2 | 65.5 | 63.1 | 62.3 |
| 20-24 | 153.9 | 150.4 | 142.2 | 134.0 | 135.0 | 133.6 | 135.8 | 137.6 | 136.1 | 134.2 | 132.1 |
| 25-29 | 129.0 | 131.7 | 129.7 | 125.5 | 131.8 | 129.8 | 132.1 | 128.0 | 117.0 | 119.3 | 123.3 |
| 30-34 | 75.8 | 74.7 | 76.5 | 77.2 | 76.6 | 79.1 | 79.6 | 82.7 | 77.2 | 77.2 | 79.1 |
| 35-39 | 31.3 | 30.8 | 31.3 | 31.3 | 32.5 | 31.5 | 33.8 | 34.1 | 33.3 | 33.5 | 35.0 |
| 40-44 | 6.9 | 5.9 | 7.0 | 6.6 | 7.4 | 7.0 | 6.8 | 7.1 | 7.6 | 7.5 | 7.1 |
| *2002 data | are provis | ional | **Rate ne | r 1.000 wor | nen in spec | cified group |) | | | | |


SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics

| | UNITED STATES BIRTH RATES* BY AGE OF MOTHER 1992-2002 | | | | | | | | | | | | | |
|-------|---|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--|--|--|
| AGE | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | | | |
| 15-19 | 60.7 | 59.6 | 58.9 | 56.8 | 54.4 | 52.3 | 51.1 | 49.6 | 47.7 | 45.3 | 43.0 | | | |
| 20-24 | 114.6 | 112.6 | 111.1 | 109.8 | 110.4 | 110.4 | 111.2 | 111.0 | 109.7 | 106.2 | 103.6 | | | |
| 25-29 | 117.4 | 115.5 | 113.9 | 112.2 | 113.1 | 113.8 | 115.9 | 117.8 | 113.5 | 113.4 | 113.6 | | | |
| 30-34 | 80.2 | 80.8 | 81.5 | 82.5 | 83.9 | 85.3 | 87.4 | 89.6 | 91.2 | 91.9 | 91.5 | | | |
| 35-39 | 32.5 | 32.9 | 33.7 | 34.3 | 35.3 | 36.1 | 37.4 | 38.3 | 39.7 | 40.6 | 41.4 | | | |
| 40-44 | 5.9 | 6.1 | 6.4 | 6.6 | 6.8 | 7.1 | 7.3 | 7.4 | 8.0 | 8.1 | 8.3 | | | |


* Rate per 1,000 women in specified group.

SOURCE: CDC's, National Center for Health Statistics: NVSR Volume 52, Number 10.


| | RESIDENT LIVE BIRTHS TO MOTHERS UNDER 20 YEARS OF AGE 1992-2002 | | | | | | | | | | | | | |
|----------------|--|-----------|-----------|------------|------------|----------|-----------|------------|-----------|----------|-------|--|--|--|
| | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* | | | |
| Valencia Co. | 152 | 145 | 168 | 176 | 209 | 181 | 177 | 214 | 186 | 185 | 177 | | | |
| New Mexico | New Mexico 4,749 4,965 4,964 4,950 4,850 4,789 4,964 4,841 4,728 4,603 4,592 | | | | | | | | | | | | | |
| *2002 data are | *2002 data are provisional | | | | | | | | | | | | | |
| SOI | URCE: NM | I Departm | ent of He | ealth, Off | ice of Nev | w Mexico | Vital Rec | ords and l | Health St | atistics | | | | |



| | RESIDENT LIVE BIRTHS TO SINGLE MOTHERS 1992-2002 | | | | | | | | | | | | | |
|----------------|---|------|------|------|------|------|------|------|------|------|-------|--|--|--|
| | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* | | | |
| Valencia Co. | 288 | 298 | 328 | 371 | 391 | 395 | 382 | 431 | 408 | 438 | 419 | | | |
| New Mexico | New Mexico 11,021 11,522 11,408 11,388 11,347 11,590 12,021 12,236 12,400 12,549 12,984 | | | | | | | | | | | | | |
| *2002 data are | 2002 data are provisional | | | | | | | | | | | | | |
| SO | SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics | | | | | | | | | | | | | |




| | RESIDENT LIVE BIRTHS TO MOTHERS WITH LESS THAN 12 YEARS OF EDUCATION 1992-2002 | | | | | | | | | | | | | |
|----------------|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--|--|--|
| | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* | | | |
| Valencia Co. | 207 | 193 | 229 | 250 | 255 | 236 | 250 | 280 | 280 | 293 | 275 | | | |
| New Mexico | 7,898 | 7,806 | 7,545 | 7,269 | 7,415 | 6,961 | 7,309 | 7,342 | 7,408 | 7,459 | 7,679 | | | |
| *2002 data are | 2002 data are provisional NOTE: Births to mothers of all ages. Some mothers are still in school. | | | | | | | | | | | | | |
| S | SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics | | | | | | | | | | | | | |



NEW MEXICO RESIDENT LIVE BIRTHS WITH FIRST TRIMESTER PRENATAL CARE 1992-2002 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002* Valencia Co. 440 490 563 583 622 628 544 503 598 650 643 17,932 18,293 17,344 17,893 18,011 18,264 17,411 16,790 17,757 17,741 New Mexico 16,656 *2002 data are provisional

SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics





KESSNER INDEX VERSUS KOTELCHUCK INDEX


Both the Kessner Index and the Kotelchuck Index utilize the month of pregnancy in which prenatal care began and the number of prenatal visits. The Kessner Index relies very heavily on the timing of the initiation of prenatal care. For example, inadequate prenatal care is defined as no prenatal care or care beginning in the third trimester. The Kessner Index has been modified by most states to allow for insufficient visits even when care was initiated early. The exact combination varies from state to state so that there is no standard modified Kessner.


The Kotelchuck Index is more sophisticated and uses two dimensions, the timing of the initiation of care and the adequacy of services received (ratio of observed visits to expected visits), which are combined into a single summary index.

RESIDENT LIVE BIRTHS ADEQUACY OF PRENATAL CARE UTILIZATION (Kotelchuck Index) 1996-2002 1996 1997 1998 1999 2000 2001 2002* **Inadequate Care** 287 199 Valencia Co. 218 190 263 269 209 New Mexico 5,811 5,541 6.284 6.499 6.333 5.854 6.321 Intermediate Care 151 Valencia Co. 197 176 174 147 135 133 New Mexico 4.943 4,845 4,660 4,331 4,122 4.187 4,663 Adequate Care Valencia Co. 267 295 230 244 269 258 New Mexico 8.826 8.417 8.239 8.052 7.915 8.165 8.498 **Intensive Care** Valencia Co. 199 216 172 175 264 265 315 7,097 6,182 6,341 6,109 5,885 New Mexico 7,090 6,526 * 2002 data are provisional SOURCE: NM Department of Health; ONMVRHS












| RESIDENT LIVE BIRTHS LEVEL OF PRENATAL CARE (Modified Kessner Index) 1996-2002 | | | | | | | | | | | | | |
|--|--------|-------------------------|---------|---------|----------|--------|--------|--|--|--|--|--|--|
| | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* | | | | | | |
| | | L | ow or N | o Pren | atal Ca | re | | | | | | | |
| Valencia Co. | 119 | 106 | 156 | 175 | 149 | 87 | 111 | | | | | | |
| New Mexico | 3,203 | 2,973 | 3,413 | 3,665 | 3,523 | 3,065 | 3,356 | | | | | | |
| | | Mid-Level Prenatal Care | | | | | | | | | | | |
| Valencia Co. | 311 | 303 | 306 | 301 | 283 | 310 | 271 | | | | | | |
| New Mexico | 8,337 | 8,166 | 8,551 | 8,258 | 8,045 | 8,190 | 8,485 | | | | | | |
| | | | High I | Prenata | l Care | | | | | | | | |
| Valencia Co. | 452 | 475 | 383 | 382 | 486 | 533 | 501 | | | | | | |
| New Mexico | 14,331 | 14,139 | 13,476 | 13,025 | 14,035 | 14,164 | 14,284 | | | | | | |
| | | U | nknow | n Prena | ıtal Car | ·e | | | | | | | |
| Valencia Co. | 83 | 85 | 119 | 142 | 74 | 70 | 69 | | | | | | |
| New Mexico | 1,345 | 1,566 | 1,854 | 2,185 | 1,603 | 1,682 | 1,581 | | | | | | |
| * 2002 data are provisional | | | | | | | | | | | | | |
| | CE: NM | | tmento | f Healt | h. ONM | VRHS | | | | | | | |









M

UNKNOWN LEVEL ?

MID-LEVEL





| | | | | | | E BIRTHS FTENDAN 002 | | | | | | | | |
|---|-----------|-------|-------|-------|-------|----------------------------|-------|-------|-------|-------|-------|--|--|--|
| | Physician | | | | | | | | | | | | | |
| | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* | | | |
| Valencia Co. | 586 | 569 | 638 | 669 | 735 | 693 | 643 | 579 | 560 | 544 | 512 | | | |
| New Mexico 23,454 22,492 22,309 21,965 22,101 21,387 20,932 19,696 19,407 19,063 19,231 | | | | | | | | | | | | | | |
| Certified Nurse Midwife | | | | | | | | | | | | | | |
| Valencia Co. | 189 | 184 | 171 | 173 | 217 | 264 | 305 | 409 | 420 | 444 | 431 | | | |
| New Mexico | 3,977 | 4,756 | 4,738 | 4,474 | 4,622 | 5,040 | 5,912 | 7,011 | 7,338 | 7,619 | 7,937 | | | |
| | | | | | L | ay Midwife | • | | | | | | | |
| Valencia Co. | 6 | 2 | 2 | 5 | 1 | 4 | 6 | 6 | 8 | 7 | 2 | | | |
| New Mexico | 286 | 293 | 292 | 272 | 254 | 240 | 248 | 269 | 290 | 268 | 336 | | | |
| | Other | | | | | | | | | | | | | |
| Valencia Co. | 7 | 14 | 7 | 10 | 12 | 8 | 10 | 6 | 4 | 5 | 7 | | | |
| New Mexico | 193 | 290 | 246 | 203 | 239 | 177 | 202 | 157 | 171 | 151 | 204 | | | |


*2002 data are provisional

SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics


PERCENT OF RESIDENT LIVE BIRTHS BY TYPE OF ATTENDANT 1992-2002

| | Physician | | | | | | | | | | | | | |
|--------------|-------------------------|------|------|------|------|------------|------|------|------|------|-------|--|--|--|
| | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* | | | |
| Valencia Co. | 74.4 | 74.0 | 78.0 | 78.1 | 76.2 | 71.5 | 66.7 | 57.9 | 56.5 | 54.4 | 53.8 | | | |
| New Mexico | 84.0 | 80.8 | 80.9 | 81.6 | 81.2 | 79.7 | 76.7 | 72.6 | 71.3 | 70.3 | 69.4 | | | |
| | Certified Nurse Midwife | | | | | | | | | | | | | |
| Valencia Co. | 24.0 | 23.9 | 20.9 | 20.2 | 22.5 | 27.2 | 31.6 | 40.9 | 42.3 | 44.4 | 45.3 | | | |
| New Mexico | 14.2 | 17.1 | 17.2 | 16.6 | 17.0 | 18.8 | 21.7 | 25.8 | 27.0 | 28.1 | 28.6 | | | |
| | | | | · | L | ay Midwife | e | · | · | | | | | |
| Valencia Co. | 0.8 | 0.3 | 0.2 | 0.6 | 0.1 | 0.4 | 0.6 | 0.6 | 0.8 | 0.7 | 0.2 | | | |
| New Mexico | 1.0 | 1.1 | 1.1 | 1.0 | 0.9 | 0.9 | 0.9 | 1.0 | 1.1 | 1.0 | 1.2 | | | |
| | | _ | | | | Other | | | | _ | | | | |
| Valencia Co. | 0.9 | 1.8 | 0.9 | 1.2 | 1.2 | 0.8 | 1.0 | 0.6 | 0.4 | 0.5 | 0.7 | | | |
| New Mexico | 0.7 | 1.0 | 0.9 | 0.8 | 0.9 | 0.7 | 0.7 | 0.6 | 0.6 | 0.6 | 0.7 | | | |
| *2002 1 / | | 1.0 | 0.7 | 0.0 | 0.7 | 0.7 | 0.7 | 0.0 | 0.0 | 0.0 | | | | |

*2002 data are provisional



| | RESIDENT LIVE BIRTHS LOW BIRTH WEIGHT BIRTHS (< 2500 grams) 1992-2002 | | | | | | | | | | | | | |
|----------------|---|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--|--|--|
| | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | | | |
| Valencia Co. | 74 | 49 | 67 | 58 | 85 | 89 | 72 | 87 | 83 | 85 | 77 | | | |
| New Mexico | 2,014 | 2,032 | 2,023 | 2,017 | 2,041 | 2,086 | 2,042 | 2,078 | 2,180 | 2,147 | 2,224 | | | |
| *2002 data are | provision | al | | • | | | | | | , | | | | |
| sou | SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics | | | | | | | | | | | | | |



Low Birth Weight

Low birth weight (LBW) is defined as an infant weight of less than 2,500 grams (5 1/2 lbs) at the time of delivery. Birth weight is one of the most important factors in determining the survival and health of an infant. Low birth weight is a public health problem since infants born LBW have increased infant mortality, morbidity, incidence of learning disabilities, and medical costs. Nationally, the risk factors for LBW include maternal age of less than seventeen and greater than 34 years, Black race/ethnicity, low socioeconomic status, single marital status, lower levels of maternal education, smoking, inadequate weight gain, low pre-pregnancy weight, and a variety of medical risk factors (Source: Data to Action).

Smoking, which retards fetal growth, is the single largest modifiable risk factor for LBW and infant mortality. It is estimated to account for 20 to 30% of all LBW births.

RESIDENT LIVE BIRTHS HIGH BIRTH WEIGHT BIRTHS (4090+ grams, 9 pounds) 1994-2002

| | 1994- | 1996 | 1997- | 1999 | 2000-2002* | | |
|--------------|--------|---------|--------|---------|------------|---------|--|
| | Number | Percent | Number | Percent | Number | Percent | |
| Valencia Co. | 108 | 4.1 | 135 | 4.6 | 147 | 5.0 | |
| New Mexico | 4,010 | 5.2 | 3,810 | 4.9 | 3,674 | 4.7 | |

*2002 Data are provisional

SOURCE: NM Department of Health, ONMVRHS

RESIDENT LIVE BIRTHS

VERY LOW BIRTH WEIGHT BIRTHS (<1500 grams) 1994-2002

| | 1994- | -1996 | 1997- | -1999 | 2000-2002* | | |
|--------------|--------|---------|--------|---------|------------|---------|--|
| | Number | Percent | Number | Percent | Number | Percent | |
| Valencia Co. | 29 | 1.1 | 37 | 1.3 | 40 | 1.4 | |
| New Mexico | 894 | 1.1 | 880 | 1.1 | 1,005 | 1.2 | |

*2002 data are provisional

SOURCE: NM Department of Health, ONMVRHS

High Birth Weight

A woman with a history of gestational diabetes or delivery of a baby weighing >9 pounds is at risk of developing type 2 diabetes. In addition, if the baby has macrosomia (generally defined as a weight of 4000 g or more), he or she is at risk for obesity and also for developing type 2 diabetes later on in life. It should be noted that there is no widely agreed-upon weight definition. Any evaluation of fetal weight must be considered in the context of gestational age.


Sources: "Diabetes in the Life Cycle and Research," Vol. 4. 2003. Editor M. Franz. Published by the American Association of Diabetes Educators, Chicago Illinois. Diabetes Care Journal, Supplement 1, January 2003 Vol. (26).

MATERNAL AND INFANT HEALTH - MEDICAID BIRTHS / SUPPLEMENTAL PROGRAMS / FEMALE POPULATION

Medicaid is a Federal-State matching entitlement program that pays for medical assistance. Originating in 1965 as Title XIX of the Social Security Act, it was created as a medical extension of the federally funded cash income assistance program emphasizing care for dependent children and their mothers, the disabled, and the elderly.

Federal and state initiatives intending to expand access to maternity and child health care have resulted in dramatic changes in policy and manner of health care delivery. Federal Omnibus Budget Reconciliation Acts (OBRA) of 1986, 1987, and 1989 required states to grant Medicaid eligibility/coverage for pregnant woman and infants, ultimately, at or below the 133% of federal poverty levels, and allow for eligibility up to 185%. The State of New Mexico followed this progression and complimented the initiatives by extending eligibility up to 225% of federal poverty levels.

Statewide, Medicaid paid for 49% of births.



NEW MEXICO RESIDENT BIRTHS BY MEDICAID STATUS 1999-2000

| Residence | Non- Medicaid Births | Medicaid Paid Births | Total Births |
|------------|----------------------------|-------------------------|-----------------|
| Valencia | 953 | 1,037 | 1,990 |
| New Mexico | 27,710 | 26,600 | 54,310 |

SOURCE: NM Department of Human Services and Department of Health, O.N.M.V.R.H.S.

Compared to mothers of children whose births were not paid by Medicaid, Medicaid mothers in New Mexico tended to:

- Be younger;
- Be White Hispanic;
- Have fewer years of education;
- Have lower levels of prenatal care;
- Be outside of the major metropolitan areas;
- Be more at risk for preterm and low weight births.

WOMEN, INFANT & CHILDREN (WIC) PROGRAM & COMMODITY SUPPLEMENTAL FOOD PROGRAM (CSFP) POTENTIAL ELIGIBLES FISCAL YEAR 2001 Estimated CSFP WIC **Potential** Percent **Eligible** Recipients Recipients Served Valencia Co. 3,937 2,170 22 55.7

> SOURCE: NM Department of Health Family Food & Nutrition Services, WIC Program

96,571

New Mexico

55,934

4,889

PERCENT OF ALL NEW MEXICO
WIC AND CSFP
ESTIMATED ELIGIBLES
FISCAL YEAR 2001

4.1

95.9

Valencia Co. All Other Counties

SOURC E: NM Department of Health
Family Food & Nutrition Sevices, WIC Program

| VALENCIA FEMALE POPULATION AGES 15-44 2000, 2001 and 2002 | | | | | | | | | | | | | | |
|---|-----------------------------------|-------|-------|----------------|--------|--------|--------|--|--|--|--|--|--|--|
| Age Groups | 2000 | 2001 | 2002 | Age Groups | 2000 | 2001 | 2002 | | | | | | | |
| 15 to 19 years | 2,500 | 2,644 | 2,736 | 35 to 39 years | 2,693 | 2,583 | 2,612 | | | | | | | |
| 20 to 24 years | 1,814 | 1,915 | 2,005 | 40 to 44 years | 2,659 | 2,778 | 2,783 | | | | | | | |
| 25 to 29 years | 1,967 | 1,823 | 1,806 | 15 to 44 years | 14,002 | 14,045 | 14,183 | | | | | | | |
| 30 to 34 years | 30 to 34 years 2,369 2,302 2,241 | | | | | | | | | | | | | |
| | Source: U.S. Bureau of the Census | | | | | | | | | | | | | |

63.0


RESIDENT INFANT DEATHS NEONATAL AND POSTNEONATAL MORTALITY RATES 1993-1997 & 1998-2002 AVERAGE


| | All Infant Deaths | | Neo | natal | Postneonatal | | |
|--------------|-------------------|------------|-----------|------------|--------------|------------|--|
| | 1993-1997 | 1998-2002* | 1993-1997 | 1998-2002* | 1993-1997 | 1998-2002* | |
| Valencia Co. | 8.2 | 6.9 | 6.1 | 3.8 | 2.1 | 3.0 | |
| New Mexico | 7.1 | 6.6 | 4.2 | 4.0 | 2.9 | 2.6 | |

^{* 2002} data are provisional.

Rate per 1,000 live births in mid year.

SOURCE: NM Department of Health, ONMVRHS





INFANT MORTALITY RATES BY COUNTY OF RESIDENCE FOR SELECTED CAUSES 1993-1997 & 1998-2002 AVERAGE

| | All Infant Deaths | | Birth Defects | | Short Gestation | | SIDS | |
|--------------|-------------------|------------|---------------|------------|-----------------|------------|-----------|------------|
| | 1993-1997 | 1998-2002* | 1993-1997 | 1998-2002* | 1993-1997 | 1998-2002* | 1993-1997 | 1998-2002* |
| Valencia Co. | 8.2 | 6.9 | 2.5 | 2.2 | 1.5 | 1.1 | 1.0 | 0.4 |
| New Mexico | 7.1 | 6.6 | 1.7 | 1.8 | 1.3 | 1.4 | 1.1 | 0.6 |

^{*2002} data are provisional

NOTE: Rate per 1,000 live births. Subcategory rates may not add up to all infant death rate due to rounding.

Due to the implementation of ICD-10 in 1999, comparability ratios have been applied to statistics for years prior to 1999. Rates and numbers may not equal those previously reported due to the application of comparability ratios.

NEW MEXICO RESIDENTS AGES 20 AND YOUNGER SELECTED DIAGNOSES REPORTED TO CHILDRENS CHRONIC CONDITIONS REGISTRY 2000

| Diagnasas | Valenc | ia Co. | New Mexico | | |
|----------------------------|--------|--------|------------|--------|--|
| Diagnoses | Number | Rate | Number | Rate | |
| Children in Registry | 2,891 | 1282.0 | 90,763 | 1534.5 | |
| Congenital Anomalies - All | 775 | 343.7 | 22,241 | 376.0 | |
| Neural Tube Defects | 33 | 14.6 | 870 | 14.7 | |
| Cleft Lip/Cleft Palate | 78 | 34.6 | 2,946 | 49.8 | |
| Down Syndrome | 33 | 14.6 | 997 | 16.9 | |
| All Other Anomalies | 631 | 279.8 | 17,428 | 294.6 | |
| Asthma | 628 | 278.5 | 20,904 | 353.4 | |

* = 1-3 **= 0

Rates per 10,000 population ages 20 years and younger

NOTE: Children in registry totals include all other conditions not listed above.

SOURCE: NM Department of Health Children's Medical Services

CHILD ABUSE CASES NUMBER AND TYPE OF INVESTIGATIONS 2002


| | Substantiated | Unsubstantiated | Total |
|--------------|---------------|-----------------|--------|
| Valencia Co. | 134 | 370 | 504 |
| New Mexico | 4,089 | 10,067 | 14,156 |

Substantiated means that through the course of the investigation, the social worker determined that the child(ren) who is the subject of the report has been determined to have been the victim of abuse and/or neglect. In other words the allegations are true.

An *unsubstantiated* investigation indicates that the investigator was unable to determine that the allegation(s) was, true.

SOURCE: NM Children, Youth and Families Department, Child Protective Services







Sometimes children have special developmental needs or health concerns that may put them at risk for or cause a developmental delay.

Through the Department of Health's Family Infant Toddler (FIT) Program early intervention services are available to help families who have concerns about the development of their young child (birth to three).

INDIVIDUALIZED FAMILY SERVICE PLAN (IFSP) ELIGIBILITY STATUS FOR FY03

| ELIGIBILITY | Valencia | New Mexico |
|-----------------------------|----------|------------|
| ELIGIDILI I Y | NUMBER | NUMBER |
| At Risk, Environmental | 27 | 665 |
| At Risk, Medical/Biological | 40 | 1,150 |
| Developmental Delay | 98 | 2,937 |
| Established Condition | 13 | 470 |

The eligibility category of children with an active IFSP during FY 2003 is reported. Children are eligible for FIT Program service through one of four eligibility categories: a developmental delay of 25% or more in one of five developmental areas, an established condition (such as Down's Syndrome) that usually leads to a development delay, or a medical / biological risk or an environmental risk for a developmental delay.

SOURCE: DOH Long Term Services Division Family Infant Toddler (FIT) Program

INDIVIDUALIZED FAMILY SERVICE PLAN (IFSP) BY RACE / ETHNICITY FOR FY03

| Valencia | New Mexico | | | | | | | |
|----------|------------|--|--|--|--|--|--|--|
| NUMBER | NUMBER | | | | | | | |
| 11 | 779 | | | | | | | |
| 0 | 43 | | | | | | | |
| ^ | 130 | | | | | | | |
| 121 | 2,696 | | | | | | | |
| 45 | 1,574 | | | | | | | |
| | | | | | | | | |

The race / ethnicity of children with active IFSP during FY 2003 is reported ^Statistics not displayed due to less than 5 cases.

SOURCE: DOH Long Term Services Division Family Infant Toddler (FIT) Program

INDIVIDUALIZED FAMILY SERVICE PLAN (IFSP) BY AGE OF CHILD on 12/1/2002

| AGE OF CHILD | Vale | encia | New Mexico | | |
|-------------------|--------|---------|------------|---------|--|
| AGE OF CHILD | NUMBER | PERCENT | NUMBER | PERCENT | |
| Less Than 1 year | 16 | 18.4 | 458 | 17.9 | |
| 1 to 2 years | 26 | 29.9 | 707 | 27.6 | |
| 2 to 3 years | 36 | 41.4 | 1048 | 41.0 | |
| 3 years and older | 9 | 10.3 | 345 | 13.5 | |
| Total | 87 | 100.0 | 2558 | 100.0 | |

The age of child with an IFSP on December 1st as reported

SOURCE: DOH Long Term Services Division Family Infant Toddler (FIT) Program

Fiscal Year 2003 = July 01, 2002 to June 30, 2003

PRIVATE AND PUBLIC SCHOOL ENROLLMENT KINDERGARTEN THROUGH 8TH GRADE 2002-2003

| GRADE | Valend | cia Co. | New Mexico | | |
|--------------|---------|---------|------------|---------|--|
| GRADE | PRIVATE | PUBLIC | PRIVATE | PUBLIC | |
| Kindergarten | 38 | 991 | 3,525 | 22,638 | |
| 1st Grade | 36 | 954 | 2,667 | 23,706 | |
| 2nd Grade | 38 | 957 | 2,738 | 23,550 | |
| 3rd Grade | 33 | 959 | 2,601 | 24,012 | |
| 4th Grade | 40 | 1,028 | 2,553 | 24,995 | |
| 5th Grade | 29 | 1,079 | 2,554 | 25,310 | |
| 6th Grade | 40 | 1,117 | 2,666 | 25,702 | |
| 7th Grade | 34 | 1,116 | 2,628 | 25,425 | |
| 8th Grade | 42 | 1,150 | 2,555 | 25,630 | |
| Total | 330 | 9,351 | 24,487 | 220,968 | |


SOURCE: NM Department of Education


| HOMESCHOOL STUDENTS ALL GRADES 1998-1999 & 1999-2000 | | | | | | |
|--|-------|--------|-------|--------|--|--|
| | 1998 | -1999 | 1999- | -2000 | | |
| | FALL | SPRING | FALL | SPRING | | |
| Valencia Co. | 454 | 539 | 437 | 515 | | |
| New Mexico | 5,796 | 5,728 | 6,080 | 6,732 | | |

NOTE: These are the number of home school students reporting to the New Mexico Department of Education.

SOURCE: NM Department of Education








NEW MEXICO RESIDENT DEATHS FOR AGES 1-14 YEARS 1992-2002

| Year | Valencia Co. | New Mexico |
|-------|--------------|------------|
| 1992 | 6 | 120 |
| 1993 | 9 | 133 |
| 1994 | 3 | 125 |
| 1995 | 6 | 123 |
| 1996 | 2 | 128 |
| 1997 | 5 | 106 |
| 1998 | 3 | 115 |
| 1999 | 2 | 100 |
| 2000 | 4 | 79 |
| 2001 | 0 | 94 |
| 2002* | 3 | 92 |
| | | |

*2002 data are provisional

SOURCE: NM Department of Health ONMVRHS



SOURCE: NM Department of Health
Office of New Mexico Vital Records and Health Statistics

NEW MEXICO RESIDENT DEATHS LEADING CAUSES OF DEATH - AGES 1-14 YEARS 1998-2002* AVERAGE


| VALENCIA COUNT | NEW MEXICO | | | | |
|--------------------------------------|------------|-------|------------------------------------|--------|-------|
| Causes | Number | Rate | Causes | Number | Rate |
| All Causes | 12 | 15.4 | All Causes | 480 | 24.4 |
| Accidents (Unintentional Injuries) | 6 | 7.8 | Accidents (Unintentional Injuries) | 190 | 9.7 |
| (Motor Vehicle Accidents) | (2) | (2.4) | (Motor Vehicle Accidents) | (128) | (6.5) |
| Congenital Malformations | 2 | 2.6 | Congenital Malformations | 45 | 2.3 |
| Conditions Orig. in Perinatal Period | 1 | 1.3 | Malignant Neoplasms | 39 | 2.0 |

^{* 2002} data are provisional

Rate per 100,000 population ages 1-14 years. Rate numerator was deaths for years 1998-2002, divided by 5; denominator was the 2000 population.

Due to the implementation of ICD-10 in 1999, comparability ratios have been applied to statistics for years prior to 1999. Rates and numbers may not equal those previously reported due to the application of comparability ratios.

SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics




*2002 data are provisional. **Accidents are considered Unintentional Injuries. *** Motor Vehicle Accidents (MVA) are included in Accidents. Rate numerator was deaths for years 1998-2002, divided by 5; denominator was the 2000 population.


Due to the implementation of ICD-10 in 1999, comparability ratios have been applied to statistics for years prior to 1999. Rates and numbers may not be equal to those previously reported due to application of comparability ratios.

PUBLIC HIGH SCHOOL ENROLLMENT **GRADES 9-12** SCHOOL YEARS 2000-2003 SCHOOL YEAR Valencia Co. New Mexico 2000-2001 3,650 95,427 2001-2002 3,728 95,224 2002-2003 3,778 95,767 SOURCE: NM Department of Education

PRIVATE AND PUBLIC HIGH SCHOOL ENROLLMENT **GRADES 9 - 12 SCHOOL YEAR 2000-2001** Valencia Co. New Mexico **GRADE** PRIVATE PUBLIC PRIVATE PUBLIC 24 1.092 2,454 28,944 9th 10th 24 1,033 2,092 25,476 11th 29 867 1,901 21,905 12th 14 658 19,102 1,690 Total 91 3,650 8,137 95,427

SOURCE: NM Department of Education





Rate calculated by dividing the number of dropouts by the number of students enrolled in grades 9-12 on the 40th day of enrollment. A dropout is a student who was enrolled in school the previous year, but did not enroll the following year and did not graduate or meet certain approved exclusions (e.g., transfer, illness).


SOURCE: NM Department of Education


| RESIDENT FALL ENROLLMENT TO NEW MEXICO | | | | | | | | |
|---|---------------|---------------|------------|--------|--|--|--|--|
| HIGHER EDUCATION INSTITUTIONS 2001-2002 | | | | | | | | |
| INSTITUTE | Vale | | New Mexico | | | | | |
| | 2001 | 2002 | 2001 | 2002 | | | | |
| UNM | 821 | 863 | 19,751 | 20,497 | | | | |
| UNM Gallup | 1 | 2 | 2,125 | 2,485 | | | | |
| UNM Los Alamos | 2 | 1 | 824 | 918 | | | | |
| UNM Medical School | 11 | 8 | 263 | 268 | | | | |
| UNM Taos | 1 | 1 | 1,190 | 1,349 | | | | |
| UNM Valencia | 1,268 | 1,429 | 1,563 | 1,717 | | | | |
| NMSU | 179 | 187 | 10,897 | 11,330 | | | | |
| NMSU Alamogordo | 1 | 0 | 1,300 | 1,380 | | | | |
| NMSU Carlsbad | 3 | 3 | 1,028 | 1,108 | | | | |
| NMSU Dona Ana | 23 | 25 | 4,342 | 4,744 | | | | |
| NMSU Grants | 34 | 16 | 619 | 640 | | | | |
| ENMU | 26 | 24 | 3,144 | 3,120 | | | | |
| ENMU Roswell | 13 | 62 | 3,046 | 3,536 | | | | |
| ENMU Ruidoso | 0 | 0 | 752 | 751 | | | | |
| NMHU | 79 | 83 | 2,987 | 2,841 | | | | |
| WNMU | 10 | 11 | 2,109 | 2,277 | | | | |
| Other NM Institutions* | 653 | 812 | 38,739 | 41,773 | | | | |
| SOURCE | E: NM Commiss | ion on Higher | Education | | | | | |

NOTE: Not all resident students report their original county of origin. In addition, out of state students are not included. Resident students attending school out of state are also not included. Counts reflect all students at a specific site, including extended services enrollments.

*Includes all other institutions of higher education not listed above, including community and technical colleges.


| PERSONS 19 AND UNDER KILLED OR INJURED AS VEHICLE OCCUPANTS IN CRASHES BY SEATBELT STATUS 2002 | | | | | | |
|--|--|---------|------|-------|---------|-------|
| | | KILLED | | | INJURED | |
| | TOTAL | NO BELT | BELT | TOTAL | NO BELT | BELT |
| Valencia Co. | 4 | 3 | 1 | 142 | 26 | 116 |
| New Mexico | 81 | 59 | 22 | 6,407 | 974 | 5,433 |
| NOTE: Number excludes pedestrians, bicyclists, motorcyclists, and unknown. | | | | | | |
| SOURCE: | SOURCE: New Mexico Department of Transportation, Traffic Safety Bureau | | | | | |





| NUMBER OF PERSONS ARRESTED FOR DRIVING WHILE IMPAIRED (DWI) 2001 & 2002 | | | | | |
|---|--------------|----------------------------------|------|------|--|
| | 19 Years and | 19 Years and Younger All Persons | | | |
| | 2001 | 2002 | 2001 | 2002 | |
| Valencia Co. | 58 | 39 | 562 | 528 | |
| New Mexico 3,617 3,593 21,576 21,603 | | | | | |
| SOURCE: NM Department of Transportation, Traffic Safety Bureau | | | | | |





NEW MEXICO RESIDENT LIVE BIRTHS TO MOTHERS AGE 15-17 YEARS 1998-2002* AVERAGE

| | Number | Rate |
|--------------|--------|------|
| Valencia Co. | 357 | 43.5 |
| New Mexico | 8,687 | 40.0 |

*2002 data are provisional


Rate is per 1,000 female population ages 15-17 years. Rate numerator was births to mothers age 15-17 years for 1998-2002, divided by 5; denominator was 2000 female population age 15-17.

SOURCE: NM Department of Health **ONMVRHS**


NEW MEXICO RESIDENT DEATHS FOR AGES 15-19 YEARS 1992-2002

| Year | Valencia Co. | New Mexico | |
|----------------------------|--------------|------------|--|
| 1992 | 7 | 117 | |
| 1993 | 5 | 132 | |
| 1994 | 9 | 128 | |
| 1995 | 7 | 150 | |
| 1996 | 3 | 158 | |
| 1997 | 6 | 118 | |
| 1998 | 6 | 144 | |
| 1999 | 4 | 150 | |
| 2000 | 5 | 143 | |
| 2001 | 6 | 107 | |
| 2002* | 7 | 136 | |
| *2002 data are provisional | | | |

SOURCE: NM Department of Health **ONMVRHS**



Office of New Mexico Vital Records and Health Statistics



SOURCE: NM Department of Health Office of New Mexico Vital Records and Health Statistics

NEW MEXICO RESIDENT DEATHS LEADING CAUSES OF DEATH - AGES 15-19 YEARS 1998-2002* AVERAGE


| VALENCIA COUNTY | | | NEW MEXICO | | |
|------------------------------------|--------|--------|------------------------------------|--------|--------|
| Cause | Number | Rate | Cause | Number | Rate |
| All Causes | 28 | 108.7 | All Causes | 680 | 93.3 |
| Accidents (Unintentional Injuries) | 11 | 42.8 | Accidents (Unintentional Injuries) | 312 | 42.8 |
| (Motor Vehicle Accidents) | (9) | (34.9) | (Motor Vehicle Accidents) | (235) | (32.2) |
| Intentional Self-Harm (Suicide) | 6 | 23.3 | Assault (Homicide) | 117 | 16.0 |
| Assault (Homicide) | 6 | 23.3 | Intentional Self-Harm (Suicide) | 117 | 16.0 |
| Malignant Neoplasms | 1 | 3.9 | Malignant Neoplasms | 35 | 4.8 |

^{* 2002} data are provisional


Rate per 100,000 population ages 15-19 years. Rate numerator was deaths for years 1998-2002, divided by 5; denominator was the 2000 population.

Due to the implementation of ICD-10 in 1999, comparability ratios have been applied to statistics for years prior to 1999. Rates and numbers may not equal those previously reported due to the application of comparability ratios.


GASTROINTESTINAL DISEASES



| HEPATITIS A NUMBER OF CASES | | | |
|---|--------------|------------|--|
| Year | Valencia Co. | New Mexico | |
| 1994-1996 | 43 | 2,254 | |
| 1997-1999 | 9 | 557 | |
| 2000-2002 | 12 | 141 | |
| Data from NETSS* | | | |
| SOURCE: NM Department of Health Office of Epidemiology | | | |



| CAMPYLOBACTER INFECTION NUMBER OF CASES | | | |
|--|--------------|------------|--|
| Year | Valencia Co. | New Mexico | |
| 1994-1996 | 53 | 1,117 | |
| 1997-1999 | 50 | 1,093 | |
| 2000-2002 | 50 | 1,059 | |
| Data from NETSS* | | | |
| SOURCE: NM Department of Health | | | |
| Office of Epidemiology | | | |
| | | | |




| SALMONELLOSIS NUMBER OF CASES | | | | |
|----------------------------------|------------|--|--|--|
| | New Mexico | | | |
| 1 | 1,016 | | | |
| 7 | 984 | | | |
| 3 | 856 | | | |
| Data from NETSS* | | | | |
| SOURCE: NM Department of Health | | | | |
| Office of Epidemiology | | | | |
| | | | | |


| Cases/100,000 population/per 30 | | 15.1 19.9 1997-1999 | Valencia Co. New Mexico 10.2 2000-2002 |
|---|---|---------------------------|--|
| Data from NETSS | • | | |
| SOURCE: NM Department of Health, Office of Epidemiology | | | |

| SHIGELLOSIS NUMBER OF CASES | | | |
|---|--------------|------------|--|
| Year | Valencia Co. | New Mexico | |
| 1994-1996 | 16 | 1,900 | |
| 1997-1999 | 38 | 786 | |
| 2000-2002 | 12 | 560 | |
| Data from NETSS* | | | |
| SOURCE: NM Department of Health Office of Epidemiology | | | |

*National Electronic Telecommunication System for Surveillance (NETSS)

BLOODBORNE DISEASES







| HEPATITIS B NUMBER OF CASES | | | | |
|---|--------------|------------|--|--|
| Year | Valencia Co. | New Mexico | | |
| 1994-1996 | 24 | 905 | | |
| 1997-1999 | 24 | 736 | | |
| 2000-2002 | 18 | 419 | | |
| Data from NETSS* | | | | |
| SOURCE: NM Department of Health Office of Epidemiology | | | | |

*National Electronic Telecommunication System for Surveillance (NETSS)

| AIDS NUMBER OF CASES | | | |
|-----------------------------------|--------------|------------|--|
| Year | Valencia Co. | New Mexico | |
| 1994-1996 | 17 | 559 | |
| 1997-1999 | 7 | 357 | |
| 2000-2002* | 7 | 270 | |
| NOTE: *2002 counts may be low due | | | |
| to delays in reporting. | | | |
| SOURCE: New Mexico HIV/AIDS | | | |
| Reporting System | | | |

RESPIRATORY DISEASES






| PERTUSSIS NUMBER OF CASES | | | | | |
|---|--------------|------------|--|--|--|
| Year | Valencia Co. | New Mexico | | | |
| 1994-1996 | 6 | 247 | | | |
| 1997-1999 | 13 | 453 | | | |
| 2000-2002 | 9 | 426 | | | |
| Data from NETSS* | | | | | |
| SOURCE: NM Department of Health Office of Epidemiology | | | | | |


| TUBERCULOSIS NUMBER OF CASES | | | | |
|---------------------------------|---------------------------|------------|--|--|
| Year | Valencia Co. | New Mexico | | |
| 1997-1998 | 5 | 139 | | |
| 1999-2000 | 2 | 110 | | |
| 2001-2002 | 0 | 111 | | |
| Data from TIMS** | | | | |
| SOURCE: NM Department of Health | | | | |
| Info | Infectious Disease Bureau | | | |

*Tuberculosis Information Management System (TIMS)


SEXUALLY TRANSMITTED DISEASES



| CHLAMYDIA NUMBER OF CASES | | | | |
|--|--------------|------------|--|--|
| Year | Valencia Co. | New Mexico | | |
| 2001 | 156 | 6,669 | | |
| 2002 | 163 | 7,419 | | |
| Data from S | TD MIS* | | | |
| SOURCE: NM Department of Health Infectious Disease Bureau | | | | |



| GONORRHEA NUMBER OF CASES | | | | |
|---------------------------------|--------------|------------|--|--|
| Year | Valencia Co. | New Mexico | | |
| 2001 | 20 | 1,090 | | |
| 2002 | 55 | 1,472 | | |
| Data from S | TD MIS* | | | |
| SOURCE: NM Department of Health | | | | |
| Infectious Disease Bureau | | | | |



| S YPHILIS NUMBER OF CASES | | | | |
|---------------------------------|--------------|------------|--|--|
| Year | Valencia Co. | New Mexico | | |
| 2001 | 0 | 19 | | |
| 2002 | 1 | 39 | | |
| Data from S | TD MIS* | | | |
| SOURCE: NM Department of Health | | | | |
| Infectious Disease Bureau | | | | |

NOTE: The scales on the above graphs are not comparable.

*Sexually Transmitted Diseases Management Information Systems (STD MIS)




NUMBER OF FATAL AND INJURY CRASHES VEHICLE MILES TRAVELED, LICENSED DRIVERS AND ECONOMIC LOSS $2001\,$

| | NUMBER OF CRASHES | | R OF CRASHES | | Economic | 100 MVM | |
|--------------|-------------------|-------|--------------|-----------|-------------|----------|--|
| | Total | Fatal | Injury | Drivers | Loss *1000 | Traveled | |
| Valencia Co. | 859 | 20 | 361 | 44,630 | \$86,548 | 4.6 | |
| New Mexico | 50,236 | 415 | 17,879 | 1,225,793 | \$3,537,921 | 227.1 | |


NOTE: MVM = Million Vehicle Miles


SOURCE: NM Department of Transportation, Traffic Safety Bureau "New Mexico Traffic Crash Information", 2001 Report.



ALL PERSONS KILLED OR INJURED AS VEHICLE OCCUPANTS IN CRASHES BY SEATBELT STATUS 2002 KILLED **INJURED** TOTAL NO BELT BELT TOTAL NO BELT BELT Valencia Co. 12 10 458 50 408 349 236 New Mexico 113 24,771 2,818 21,953 NOTE: Number excludes pedestrians, bicyclists, motorcyclists, and unknown

SOURCE: New Mexico Department of Transportation, Traffic Safety Bureau.







ALCOHOL RELATED CRASHES NEW MEXICO - 2001


- 43 percent of all fatal crashes involved alcohol.
- A person died in an alcohol-involved crash every 44 hours.
- A person was injured in an alcohol-involved crash every 174 minutes.
- An alcohol-involved crash occurred every 147 minutes.

ALCOHOL INVOLVED CRASHES PERSONS INVOLVED AND DWI CONVICTIONS 2002 CRASHES PERSONS INVOLVED

| | CRAS | SHES | | PERSONS INVOLVED | | |
|--------------|-------|--------|--------|------------------|-----------------|----------|
| | Fatal | Injury | Killed | Injured | DWI Conviction* | DWI Rate |
| Valencia Co. | 11 | 39 | 11 | 82 | 414 | 9.3 |
| New Mexico | 198 | 1,774 | 221 | 2,921 | 12,544 | 10.1 |

*DWI convictions are those handed down during year listed. The citations which resulted in these convictions could have been issued prior to these years. Convictions are counted by county of driver's residence, not the county in which the citation was issued.

SOURCE: NM Department of Transportation, Traffic Safety Bureau "Driving While Impared", 2002 Report.



*DWI convictions are those handed down during year listed. The citations which resulted in these convictions could have been issued prior to these years. Convictions are counted by county of driver's residence, not the county in which the citation was issued

SOURCE: NM Department of Transportation, Traffic Safety Bureau "Driving While Impared", various years.


| VALENCIA COUNTY INJURY DEATHS BY MECHANISM 1999-2001 | | | | | | |
|--|-----------------------------------|---|---|--|--|--|
| Mechanism of Injury | Number of Deaths in 3 Years | Age Adjusted Rate/100,000* Per Year | New Mexico Age Adjusted Rate/100,000* Per Year | | | |
| Cut/pierce | 2 | 1.0 | 1.8 | | | |
| Drowning | 2 | 1.0 | 1.4 | | | |
| Fall | 15 | 9.6 | 10.2 | | | |
| Fire/burn | 2 | 1.2 | 1.0 | | | |
| Firearm | 37 | 19.4 | 15.7 | | | |
| Motor Vehicle Traffic | 51 | 27.0 | 22.1 | | | |
| Poisoning | 43 | 22.0 | 17.0 | | | |
| Other** | 24 | 13.1 | 14.0 | | | |
| All Injury | 176 | 94.3 | 83.2 | | | |

| NON-FATAL INJURY HOSPITALIZATIONS 1998-2000 | | | | | |
|--|--------------------|---------------|--|--|--|
| Number/Rate | Valencia County | New Mexico | | | |
| Total Number of | | | | | |
| Hospital Discharges in | | | | | |
| 3 Years | 1,286 | 30,596 | | | |
| Age Adjusted Rate per | | | | | |
| 100,000* Population per | | | | | |
| Year | 694.2 | 585.2 | | | |
| NOTE III III III III | | | | | |

NOTE: Injury as the principal diagnosis of discharge.

* Rates are adjusted to the 2000 U. S. Standard
Population.

SOURCE: NM Health Policy Commission



Note: Injury categories are from the "Preliminary External Cause of Injury Mortality Matrix for ICD-10," June 12, 2001 by Centers for Disease Control and Prevention, National Center for Injury Prevention and Control.

New Mexico Department of Health, Office of Epidemiology, Injury Epidemiology Unit Source: Office of New Mexico Vital Records and Health Statistics

^{*}Age-adjusted death rate adjusted to the 2000 U.S standard population. Age adjusted death rates weight the age-specific rates for a given population by the age distribution of a standard population, thereby enabling the direct comparison of rates between populations with differing age structures.

^{**} Other includes machinery, cyclist, pedestrian, natural /environmental, struck by/against, suffocation, other specified and unspecified categories.

CANCER INCIDENCE AMONG MALES NUMBER AND INCIDENCE RATE 1994-2000 AGGREGATE

| SITE | Valencia | | New Mexico | |
|----------------------------------|----------|-------|------------|-------|
| SHE | NUMBER | RATE | NUMBER | RATE |
| All Cancers | 873 | 520.4 | 23,565 | 468.7 |
| Stomach | 13 | 8.5 | 515 | 10.6 |
| Colon and Rectum | 80 | 49.3 | 2,482 | 50.4 |
| Liver and Intrahepatic Bile Duct | 22 | 14.1 | 412 | 8.2 |
| Gallbladder | ^ | ^ | 52 | 1.1 |
| Pancreas | 20 | 13.4 | 570 | 11.7 |
| Lung and Bronchus | 120 | 73.5 | 3,053 | 62.2 |
| Melanomas of the Skin | 31 | 17.6 | 974 | 18.7 |
| Breast | ^ | ^ | 51 | 1.0 |
| Prostate | 242 | 145.4 | 7,320 | 145.5 |
| Urinary Bladder | 63 | 39.2 | 1,324 | 27.4 |
| Kidney and Renal Pelvis | 27 | 15.0 | 741 | 14.3 |
| Non-Hodgkins Lymphomas | 32 | 19.0 | 912 | 17.8 |
| Leukemias | 29 | 16.8 | 727 | 14.2 |
| All Other Cancers | 190 | 106.3 | 4,432 | 85.3 |

Rates are per 100,000 and age-adjusted to the 2000 U.S. (19 age groups) standard.

- ~ Statistic could not be calculated.
- ^ Statistic not displayed due to less than 5 cases.

SOURCE: New Mexico Tumor Registry, University of New Mexico

Colorectal Cancer Screening and Early Detection

Screening tests can reduce deaths from colorectal cancer by identifying polyps, which are tiny growths inside the colon, before they become cancerous. Screening also detects early cancers when treatment is most effective and survival rates are high.

SOURCE: www.cdc.gov


Prostate Cancer Fact

The American Cancer Society estimates that in 2003, 220,900 new cases of prostate cancer will be diagnosed and that 28,900 men will die of the disease in the United States.

Two methods for detecting prostate cancer are commonly used by clinicians:

- 1) Digital rectal examination (DRE)
- 2) Prostate-specific antigen (PSA)

SOURCE: www.cdc.gov



| CANCER INCIDENCE AMONG FEMALES NUMBER AND INCIDENCE RATE 1994-2000 AGGREGATE | | | | | | |
|--|--------|-------|--------|-------|--|--|
| SITE Valencia New Mexico | | | | | | |
| SHE | NUMBER | RATE | NUMBER | RATE | | |
| All Cancers | 703 | 363.9 | 21,575 | 353.8 | | |
| Stomach | 14 | 7.4 | 333 | 5.5 | | |
| Colon and Rectum | 67 | 37.1 | 2,173 | 35.6 | | |
| Liver and Intrahepatic Bile Duct | 7 | 4.0 | 211 | 3.4 | | |
| Gallbladder | ^ | ^ | 113 | 1.8 | | |
| Pancreas | 14 | 7.7 | 543 | 8.9 | | |
| Lung and Bronchus | 69 | 36.5 | 2,208 | 36.0 | | |
| Melanomas of the Skin | 29 | 15.1 | 745 | 12.2 | | |
| | | | | | | |

215

21

32

18

14

21

14

110.4

10.4

16.7

9.5

7.0

10.3

6.8

82.9

7,067

1,116

566

484

472

759

532

4,253

116.5

9.3

18.3

7.9

7.7

12.5

8.6

69.5

Rates are per 100,000 and age-adjusted to the 2000 U.S. (19 age groups) standard.

~ Statistic could not be calculated.

Corpus and Uterus, NOS

Kidney and Renal Pelvis

Non-Hodgkins Lymphomas

Urinary Bladder

Leukemias

All Other Cancers

^ Statistic not displayed due to less than 5 cases.

SOURCE: New Mexico Tumor Registry, University of New Mexico

Breast Cancer Screening

Breast

Cervix


According to the American Cancer Society, an estimated 211,300 new invasive cases of breast cancer are expected to occur among women in the United States during 2003. Mammography is the best way to detect breast cancer in its earliest, most treatable stage, an average of 1.7 years before a woman can feel the lump. Mammography also locates cancers too small to be felt during a clinical breast examination.

SOURCE: www.cdc.gov

Cervical Cancer Screening

Cervical cancer is nearly 100 percent preventable, yet according to the American Cancer Society, in 2003, an estimated 12,200 new cases of invasive cervical cancer will be diagnosed and about 4,100 women will die of the disease. The occurrence of deaths from cervical cancer has declined significantly over the past 40 years. This decline is largely attributed to the effectiveness of Pap test screening in detecting precancerous lesions and early stage cervical cancer.

SOURCE: www.cdc.gov



MODIFIED MAJOR DIAGNOSTIC CATEGORIES (MMDC)

➤ Hospitals use a classification scheme, called Major Diagnostic Categories (MDC), to group the diagnoses of patients upon their discharge from the hospital. Data shown in the following pages are based on a modification of the Major Diagnostic Categories. The Modified Major Diagnostic Categories (MMDCs) separates injuries and neoplasms into their own unique groupings while conventional MDCs distribute these diagnoses across other categories, which obscures their impact.


METHODOLOGY NOTES:

- The size of the county's population and the population demographics, such as average age of residents, should be taken into account in interpreting reported data.
- Indian Health Service facilities do not report discharge data to the New Mexico Health Policy Commission, the state agency that maintains the hospital discharge database. As such, counties with large Native American populations may have artificially lower rates.
- Only hospitalization to New Mexico hospitals are included. Counties bordering other states may have a significant number of hospitalization in other states.


| HOSPITAL INPATIENT DISCHARGE DATA - AI | LL AGES |
|--|---------|
| MODIFIED MAJOR DIAGNOSTIC CATEGORIES | (MMDCs) |
| 1999-2001 | |

| MMDC | 19: | 99 | 200 | 00 | 2001 | | |
|----------------------------------|--------------|----------------|-----------------|------------|--------------|------------|--|
| MINIDC | Valencia Co. | New Mexico | Valencia Co. | New Mexico | Valencia Co. | New Mexico | |
| 01: Nervous System | 203 | 6,319 | 244 | 6,416 | 258 | 6,799 | |
| 02: Eye Diseases | 15 | 639 | 13 | 386 | 4 | 205 | |
| 03: Ear, Nose, Mouth, Throat | 67 | 1,879 | 68 | 1,612 | 54 | 1,702 | |
| 04: Respiratory System | 582 | 17,875 | 462 | 15,151 | 450 | 15,329 | |
| 05: Circulatory System | 665 | 20,349 | 799 | 20,923 | 702 | 20,629 | |
| 06: Digestive System | 407 | 12,425 | 491 | 13,099 | 419 | 12,921 | |
| 07: Hepatobiliary System | 191 | 5,084 | 204 | 5,306 | 194 | 5,343 | |
| 08: Musculoskeletal System | 279 | 7,400 | 286 | 7,206 | 312 | 7,293 | |
| 09: Skin, Subcutaneous, Breast | 91 | 2,272 | 107 | 2,392 | 100 | 2,583 | |
| 10: Endocrine, Metabolic | 150 | 4,658 | 107 | 4,726 | 157 | 5,103 | |
| 11: Kidney & Urinary Tract | 148 | 4,789 | 139 | 4,787 | 172 | 5,186 | |
| 12: Male Reproductive System | 22 | 772 | 18 | 662 | 22 | 767 | |
| 13: Female Reproductive System | 144 | 3,330 | 107 | 3,496 | 127 | 3,542 | |
| 14: Pregnancy, Childbirth | 1,072 | 26,121 | 1,114 | 26,521 | 1,132 | 26,606 | |
| 15: Newborns, Neonates | 994 | 23,859 | 1,031 | 24,401 | 1,031 | 24,567 | |
| 16: Blood & Blood Forming-Organs | 32 | 1,052 | 34 | 1,170 | 44 | 1,253 | |
| 17: Myeloproliferative Diseases | 330 | 8,616 | 323 | 8,345 | 341 | 8,118 | |
| 18: Infectious Diseases | 60 | 2,128 | 54 | 1,795 | 50 | 1,915 | |
| 19: Mental Diseases | 326 | 8,878 | 287 | 9,064 | 334 | 9,485 | |
| 20: Alcohol / Drug Abuse | 151 | 3,295 | 131 | 2,957 | 96 | 2,909 | |
| 21: Injuries & Poisonings | 523 | 14,682 | 594 | 14,193 | 550 | 14,113 | |
| 22: Burns | 14 | 270 | 19 | 233 | 12 | 229 | |
| 23: Factors Influencing Health | 93 | 4,148 | 182 | 4,274 | 180 | 4,387 | |
| 24: Multiple Trauma | 0 | 0 | 0 | 0 | 0 | 0 | |
| 25: HIV | 3 | 148 | 0 | 96 | 0 | 86 | |
| | SOURCI | E: NM Health P | olicy Commissio | on | | | |

SOURCE: NM Health Policy Commission



NEW MEXICO ASTHMA DISCHARGE DATA BY COUNTY 1998-2000



SOURCE: NM Health Policy Commission

| HOSPITAL DISCHARGES BY PRIMARY PAYER 1999 & 2000 | | | | | | | | | |
|--|-------------------------------------|---------------|---------------|----------------|--|--|--|--|--|
| Payee | Vale 1999 | encia 2000 | New N 1999 | lexico 2000 | | | | | |
| Medicare | | | | 1.1.1 | | | | | |
| Medicare | 984 | 1,154 | 47,144 | 47,959 | | | | | |
| Medicaid | 1,633 | 1,546 | 39,147 | 39,468 | | | | | |
| Private | | | | | | | | | |
| Insurance | 3,421 | 3,556 | 71,401 | 74,271 | | | | | |
| Uninsured | 378 | 479 | 12,354 | 13,859 | | | | | |
| Others | 194 | 119 | 5,431 | 4,738 | | | | | |
| Total | | | | | | | | | |
| Discharges | 6,610 | 6,854 | 175,477 | 180,295 | | | | | |
| so | SOURCE: NM Health Policy Commission | | | | | | | | |

NEW MEXICO ASTHMA HOSPITAL
DISCHARGES* BY COUNTY
1998-2000
Number Rate

| | Number | Rate |
|--------------|--------|------|
| Valencia Co. | 179 | 9.2 |
| New Mexico | 4,936 | 9.4 |

*Primary discharges from non-Federal Hospitals Average rates per 10,000 population per year.

SOURCE: NM Health Policy Commission

ESTIMATED DIABETES PREVALENCE AMONG NEW MEXICO ADULTS 2002*

| County | 2002 Adult Population | Number of Adults with Diabetes | Percentage of Adults with Diabetes |
|--------------|-----------------------|-----------------------------------|-------------------------------------|
| Bernalillo | 430,698 | 38,586 | 9.0% |
| Catron | 2,844 | 228 | 8.0% |
| Chaves | 43,413 | 3,806 | 8.8% |
| Cibola | 18,460 | 1,960 | 10.6% |
| Colfax | 10,765 | 951 | 8.8% |
| Curry | 31,536 | 2,672 | 8.5% |
| De Baca | 1,666 | 142 | 8.5% |
| Dona Ana | 127,663 | 12,049 | 9.4% |
| Eddy | 37,146 | 3,195 | 8.6% |
| Grant | 22,677 | 2,022 | 8.9% |
| Guadalupe | 3,513 | 352 | 10.0% |
| Harding | 616 | 55 | 8.9% |
| Hidalgo | 3,801 | 344 | 9.1% |
| Lea | 39,662 | 3,451 | 8.7% |
| Lincoln | 15,533 | 1,269 | 8.2% |
| Los Alamos | 13,763 | 1,075 | 7.8% |
| Luna | 18,007 | 1,639 | 9.1% |
| McKinley | 47,057 | 5,630 | 12.0% |
| Mora | 3,960 | 398 | 10.0% |
| Otero | 43,945 | 3,827 | 8.7% |
| Quay | 7,524 | 646 | 8.6% |
| Rio Arriba | 29,687 | 3,093 | 10.4% |
| Roosevelt | 13,160 | 1,109 | 8.4% |
| Sandoval | 68,949 | 6,313 | 9.2% |
| San Juan | 82,679 | 8,169 | 9.9% |
| San Miguel | 22,029 | 2,197 | 10.0% |
| Santa Fe | 103,491 | 9,359 | 9.0% |
| Sierra | 10,497 | 859 | 8.2% |
| Socorro | 13,177 | 1,246 | 9.5% |
| Taos | 23,651 | 2,246 | 9.5% |
| Torrance | 11,901 | 1,030 | 8.7% |
| Union | 2,962 | 251 | 8.5% |
| Valencia | 48,121 | 4,450 | 9.2% |
| New Mexico** | 1,354,553 | 120,555 | 8.9% |

*County estimates were derived in the following manner: The Behavioral Risk Factor Surveillance System (BRFSS, 2002), a telephone survey of adults' (age 18 and over) health in New Mexico, was utilized to generate the racial/ethnic-specific diagnosed diabetes prevalence rates for the entire state. County populations were then obtained from the US Census Bureau (2002) by racial/ethnic group. The BRFSS rates were applied to these groups, generating an estimate of the number of American Indians, non-Hispanic Whites, Hispanics and Others (Asian/Pacific Islanders, non-Hispanic Blacks, and other) with diagnosed diabetes. The numbers were then combined to yield a total estimate of adults in each county with diagnosed diabetes. For those with undiagnosed diabetes, estimates by racial/ethnic group from NHANES III (National Health and Nutrition Examination Survey) were obtained from Harris, MI et al. (1998). The estimated numbers of adults with diagnosed and undiagnosed diabetes were combined for the final estimate.

**State estimates were derived in the following manner: The Behavioral Risk Factor Surveillance System (BRFSS, 2002) was utilized to generate an overall estimate of diagnosed diabetes among New Mexican adults. That is, in 2002, an estimated 6.2% of New Mexican adults indicated they had been diagnosed with diabetes. An estimate for undiagnosed diabetes (from NHANES III data) was obtained from Harris, MI et al. (1998). The estimated numbers of adults with diagnosed and undiagnosed diabetes were combined for the final estimate. Note: adding up the county estimates will not result in the state estimate. For the state estimate, there was no initial stratification by race/ethnicity, it was simply a calculation of those with diabetes (all races) divided by those responding to the question.

Source: Department of Health, Diabetes Prevention & Control Program

HOSPITAL INPATIENT DISCHARGE DATA MODIFIED MAJOR DIAGNOSTIC CATEGORIES (MMDC) VALENCIA COUNTY 1999-2001


| | | 1999 | | | 2000 | | | 2001 | |
|----------------------------------|---------|---------|------------|------------|----------|-------|-----|-------|-------|
| MMDC | | All | % All | | All | % All | | All | % All |
| | 65+ | Ages | Ages | 65+ | Ages | Ages | 65+ | Ages | Ages |
| 01: Nervous System | 106 | 203 | 52.2 | 133 | 244 | 54.5 | 129 | 258 | 50.0 |
| 02: Eye Diseases | 7 | 15 | 46.7 | 6 | 13 | 46.2 | 1 | 4 | 25.0 |
| 03: Ear, Nose, Mouth, Throat | 10 | 67 | 14.9 | 12 | 68 | 17.6 | 11 | 54 | 20.4 |
| 04: Respiratory System | 268 | 582 | 46.0 | 193 | 462 | 41.8 | 200 | 450 | 44.4 |
| 05: Circulatory System | 357 | 665 | 53.7 | 429 | 799 | 53.7 | 388 | 702 | 55.3 |
| 06: Digestive System | 131 | 407 | 32.2 | 179 | 491 | 36.5 | 138 | 419 | 32.9 |
| 07: Hepatobiliary System | 46 | 191 | 24.1 | 57 | 204 | 27.9 | 51 | 194 | 26.3 |
| 08: Musculoskeletal System | 82 | 279 | 29.4 | 108 | 286 | 37.8 | 143 | 312 | 45.8 |
| 09: Skin, Subcutaneous, Breast | 24 | 91 | 26.4 | 41 | 107 | 38.3 | 21 | 100 | 21.0 |
| 10: Endocrine, Metabolic | 49 | 150 | 32.7 | 36 | 107 | 33.6 | 54 | 157 | 34.4 |
| 11: Kidney & Urinary Tract | 54 | 148 | 36.5 | 61 | 139 | 43.9 | 74 | 172 | 43.0 |
| 12: Male Reproductive System | 15 | 22 | 68.2 | 12 | 18 | 66.7 | 13 | 22 | 59.1 |
| 13: Female Reproductive System | 18 | 144 | 12.5 | 16 | 107 | 15.0 | 17 | 127 | 13.4 |
| 14: Pregnancy, Childbirth | 0 | 1,072 | 0.0 | 0 | 1,114 | 0.0 | 0 | 1,132 | 0.0 |
| 15: Newborns, Neonates | 0 | 994 | 0.0 | 1 | 1,031 | 0.1 | 0 | 1,031 | 0.0 |
| 16: Blood & Blood-Forming Organs | 9 | 32 | 28.1 | 18 | 34 | 52.9 | 16 | 44 | 36.4 |
| 17: Myeloproliferative Diseases | 124 | 330 | 37.6 | 112 | 323 | 34.7 | 123 | 341 | 36.1 |
| 18: Infectious Diseases | 22 | 60 | 36.7 | 14 | 54 | 25.9 | 15 | 50 | 30.0 |
| 19: Mental Diseases | 28 | 326 | 8.6 | 38 | 287 | 13.2 | 39 | 334 | 11.7 |
| 20: Alcohol / Drug Abuse | 5 | 151 | 3.3 | 7 | 131 | 5.3 | 8 | 96 | 8.3 |
| 21: Injuries & Poisonings | 125 | 523 | 23.9 | 176 | 594 | 29.6 | 167 | 550 | 30.4 |
| 22: Burns | 0 | 14 | 0.0 | 1 | 19 | 5.3 | 1 | 12 | 8.3 |
| 23: Factors Influencing Health | 49 | 93 | 52.7 | 106 | 182 | 58.2 | 107 | 180 | 59.4 |
| 24: Multiple Trauma | 0 | 0 | | 0 | 0 | | 0 | 0 | |
| 25: HIV | 0 | 3 | 0.0 | 0 | 0 | | 0 | 0 | |
| | SOURCE: | New Mex | ico Healtl | h Policy C | ommissio | n | | | |

VALENCIA COUNTY HOSPITAL INPATIENT DISCHARGE STATUS FOR AGE 65 YEARS AND OLDER 1999-2001

| | 1999 | 2000 | 2001 | | | | |
|-------------------------------------|-------|-------|-------|--|--|--|--|
| Routine Discharge | 1,004 | 1,126 | 1,120 | | | | |
| Transfer-Hospital | 21 | 31 | 32 | | | | |
| Transfer-Skilled Nursing Facility | 129 | 159 | 133 | | | | |
| Transfer-Intermediate Care Facility | 23 | 70 | 64 | | | | |
| Transfer-Other | 63 | 112 | 125 | | | | |
| Transfer-Home Care | 174 | 174 | 161 | | | | |
| Left Against Medical Advice | 3 | 1 | 6 | | | | |
| Died | 110 | 78 | 72 | | | | |
| Other | 3 | 10 | 4 | | | | |
| TOTAL DISCHARGED | 1,530 | 1,761 | 1,717 | | | | |

SOURCE: New Mexico Health Policy Commission





ADULT ABUSE CASES NUMBER AND TYPE OF INVESTIGATION Annual 2002

| Residence | | ntiated gations | Unsubstantiated Investigations | | Total Investigations |
|--------------|--------|--------------------|-----------------------------------|---------|-------------------------|
| | Number | Percent | Number | Percent | Number |
| Valencia Co. | 57 | 30.8 | 128 | 69.2 | 185 |
| New Mexico | 1475 | 29.2 | 3569 | 70.8 | 5,044 |

Source: NM Children, Youth and Families Department Protective Services Division Fact Book, Fourth Quarter & Annual 2002




NEW MEXICO RESIDENT DEATHS AGES 65 YEARS AND OLDER 1992-2002

| Year | Valencia Co. | New Mexico |
|-------|--------------|------------|
| 1992 | 185 | 7,600 |
| 1993 | 219 | 7,971 |
| 1994 | 223 | 8,368 |
| 1995 | 234 | 8,628 |
| 1996 | 274 | 8,663 |
| 1997 | 269 | 8,772 |
| 1998 | 283 | 9,100 |
| 1999 | 288 | 9,515 |
| 2000 | 293 | 9,441 |
| 2001 | 337 | 9,864 |
| 2002* | 311 | 9,840 |

*2002 data are provisional

SOURCE: NM Department of Health ONMVRHS



NEW MEXICO RESIDENT DEATHS LEADING CAUSES OF DEATH - AGES 65 YEARS AND OLDER 1998-2002* AVERAGE

| VALENCIA COUNTY | NEW MEXICO | | | | |
|------------------------------------|------------|--------|------------------------------------|--------|--------|
| Cause | Number | Rate | Cause | Number | Rate |
| All Causes | 1,512 | 4498.0 | All Causes | 47,760 | 4500.9 |
| Diseases of Heart | 400 | 1189.9 | Diseases of Heart | 13,459 | 1268.4 |
| Malignant Neoplasms | 357 | 1060.6 | Malignant Neoplasms | 10,313 | 971.9 |
| Chronic Lower Respiratory Diseases | 126 | 374.7 | Chronic Lower Respiratory Diseases | 3,630 | 342.1 |
| Cerebrovascular Diseases | 102 | 303.8 | Cerebrovascular Diseases | 3,407 | 321.1 |
| Diabetes Mellitus | 60 | 178.7 | Diabetes Mellitus | 1,969 | 185.5 |


^{* 2002} data are provisional

Rate per 100,000 population ages 65 years and older. Rate numerator was deaths for years 1998-2002, divided by 5; denominator was the 2000 population.

Due to the implementation of ICD-10 in 1999, comparability ratios have been applied to statistics for years prior to 1999. Rates and numbers may not equal those previously reported due to the application of comparability ratios.

| NEW MEXICO RESIDENT DEATHS COUNTY AND STATE TOTALS 1992-2002 | | | | | | | | | | | |
|--|-----------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002* |
| Valencia Co. | 306 | 351 | 351 | 389 | 432 | 413 | 404 | 442 | 451 | 499 | 478 |
| New Mexico | 11,130 | 11,689 | 12,106 | 12,500 | 12,456 | 12,613 | 12,858 | 13,433 | 13,384 | 14,016 | 14,114 |
| * 2002 data ar | * 2002 data are provisional | | | | | | | | | | |

SOURCE: NM Department of Health, Office of New Mexico Vital Records and Health Statistics



NEW MEXICO RESIDENT DEATHS LEADING CAUSES OF DEATH 1998-2002* AVERAGE

| VALENCIA COUNT | Y | NEW MEXICO | | | |
|------------------------------------|--------|------------|------------------------------------|---------|--------|
| All Races/All Ages | Number | Rate | All Races/All Ages | Number | Rate |
| All Causes | 2,273 | 687.2 | All Causes | 67,805 | 745.5 |
| Malignant Neoplasms | 522 | 157.7 | Diseases of Heart | 16,186 | 178.0 |
| Diseases of Heart | 500 | 151.1 | Malignant Neoplasms | 14,440 | 158.8 |
| Accidents (Unintentional Injuries) | 190 | 57.4 | Accidents (Unintentional Injuries) | 4,759 | 52.3 |
| (Motor Vehicle Accidents) | (77) | (23.3) | (Motor Vehicle Accidents) | (1,904) | (20.9) |
| Chronic Lower Respiratory Diseases | 151 | 45.7 | Chronic Lower Respiratory Diseases | 4,125 | 45.3 |
| Cerebrovascular Diseases | 116 | 35.1 | Cerebrovascular Diseases | 3,890 | 42.8 |

^{* 2002} data are provisional

Rate per 100,000 population. Rate numerator was deaths for years 1998-2002, divided by 5; denominator was the 2000 population.


Due to the implementation of ICD-10 in 1999, comparability ratios have been applied to statistics for years prior to 1999. Rates and numbers may not equal those previously reported due to the application of comparability ratios.


| CHILDREN UNDER 6 WITH ELEVATED BLOOD LEAD LEVELS | | | | | | | | |
|---|-------|-------|--|--|--|--|--|--|
| OF THOSE CHILDREN SCREENED, 2001-2002 | | | | | | | | |
| | 2001 | 2002 | | | | | | |
| Valencia Co. | | | | | | | | |
| Number of Children with Elevated Blood Lead Levels | 0 | 1 | | | | | | |
| New Mexico | | | | | | | | |
| Number of Children Screened | 2,691 | 2,876 | | | | | | |
| Number of Children with Elevated Blood Lead Levels | 37 | 35 | | | | | | |
| Percent of Children with Elevated Blood Lead Levels 1.4 | | | | | | | | |
| NOTE: Elevated Blood Lead Levels is defined as ≥ 10 µg/dl of blood. | | | | | | | | |

In Valencia County, 11.3% of the houses were built prior to 1950. More than 27% pre-1950 housing in an area is considered a greater than average risk for lead poisoning in children.

- · Lead poisoning is the number one preventable environmental health problem for children under 6 years.
- · Risk factors for lead poisoning include older housing, poverty, and adult occupational exposure to lead.
- The only way to determine a child's lead level is with a blood lead test.
- The national estimate (1997) of children with an elevated blood lead level is 4.4%.²

It is estimated that 60% to 80% of children with elevated blood lead levels are in the Medicaid population.² Note: Screening rates have declined significantly in New Mexico (NM) since the mid 1990's.





 $^{^{1} \}ge 10 \,\mu\text{g/dl}$ of blood (greater than or equal to 10 micrograms per deciliter of blood), the Centers for Disease Control and Prevention (CDC) identified level of concern.

SOURCE: NM Department of Health, Office of Epidemiology

² General Accounting Office (GAO) Report, 1999.



| HEALTH CARE CLINICS | | | | | | |
|------------------------------------|---------------|-------------------------|-------------|-------|-------|--|
| FACILITY | PHONE | STREET ADDRESS | CITY | STATE | ZIP | |
| First Choice Community Health Care | (505)861-1013 | 120 S 9th St | Belen | NM | 87002 | |
| First Choice Community Health Care | (505)865-4618 | 1259 Hwy 314 | Los Lunas | NM | 87031 | |
| La Vida Felicidad | (505)865-4651 | 555 Don Pasqual NW | Los Lunas | NM | 87031 | |
| Leach, Donald B., OD | (505)865-4812 | 943 Main St SW | Los Lunas | NM | 87031 | |
| Parents Reaching Out | (505)247-0192 | 1920 B. Columbia Dr. SE | Albuquerque | NM | 87106 | |
| Presbyterian Family Healthcare | (505)864-5454 | 609 S Christopher Rd | Belen | NM | 87002 | |
| University of New Mexico-VC | (505)925-8500 | 280 La Entrada Rd | Los Lunas | NM | 87031 | |
| Valencia Counseling Services, Inc. | (505)865-3350 | 735 Don Pasqual NW | Los Lunas | NM | 87031 | |

SOURCE: NM Department of Health, District I Health Office

| VALENCIA COUNTY SELECTED COMMUNITY PARTNERS FOR PUBLIC HEALTH | | | | |
|--|----------|--|--|--|
| COMMUNITY PARTNER | PHONE# | | | |
| Belen Schools | 966-1105 | | | |
| Child Youth Families Department | 865-4634 | | | |
| DWI Councils | 866-8041 | | | |
| First Choice Community Healthcare Belen | 861-1013 | | | |
| First Choice Community Healthcare Los Lunas | 865-4618 | | | |
| La Vida Felicidad | 865-4651 | | | |
| Los Lunas Schools | 866-2440 | | | |
| Maternal Child Health Council | 865-0345 | | | |
| New Mexico Breast Feeding Task Force | 841-5335 | | | |
| Valencia Community Partnership | 865-0345 | | | |
| Valencia County News Bulletin | 864-4472 | | | |
| VC Local Emergency Planning Committee | 861-2178 | | | |
| Youth Development, Inc. | 865-0345 | | | |
| SOURCE: NM Department of Health, District I Health Office | | | | |

| LICENSED HEALTH PROFESSIONALS BY COUNTY OF LICENSE 2002 | | | |
|---|--------------|------------|--|
| | Valencia Co. | New Mexico | |
| Medical Doctor | 33 | 5,809 | |
| Medical Doctor Resident | 1 | 1,144 | |
| Physician Assistant | 6 | 287 | |
| Nurse (RN & LPN) | 475 | 19,575 | |
| Certified Nurse Midwives | 1 | 148 | |
| Licensed Midwives | 0 | 79 | |
| SOURCE: New Mexico Health Policy Commission | | | |

ACTIVE EMERGENCY MEDICAL SERVICES PERSONNEL BY LEVEL AS OF AUGUST 20, 2003

| Level | Valencia Co. | New Mexico |
|-----------------------|--------------|------------|
| First Responder | 8 | 760 |
| Dispatcher | 9 | 398 |
| Dispatcher-Instructor | 0 | 16 |
| EMT-Basic | 99 | 3,321 |
| EMT-Intermediate | 42 | 1,190 |
| EMT-Paramedic | 35 | 955 |
| Total | 193 | 6,641 |

NOTE: There are personnel with current active licenses at more than one level in this snapshot. The actual state personnel total is 6,546.

SOURCE: NM Department of Health



ACTIVE EMERGENCY MEDICAL SERVICES PERSONNEL BY GENDER AS OF AUGUST 20, 2003

| Gender | Valencia Co. | New Mexico |
|---------|--------------|------------|
| Male | 118 | 4,574 |
| Female | 74 | 1,995 |
| Unknown | 1 | 72 |
| Total | 193 | 6,641 |

NOTE: There are personnel with current active licenses at more than on level in this snapshot. The actual state personnel total is 6,546.

SOURCE: NM Department of Health



AGE-ADJUSTED—Weights the age-specific rates for a given population by the age distribution of a standard population, thereby enabling the direct comparison of rates between populations with differing age structures.

CHILDREN'S CHRONIC CONDITIONS REGISTRY—A registry of data on children with special needs; data are from programs and physicians who provide services to these children. Maintained by the New Mexico Department of Health.

COMPARABILITY RATIOS—With each revision of the *International Classification of Diseases (ICD)*, discontinuities are created in cause-of-death trend data. The classification and rule changes have a substantial impact as deaths that may have been coded to one category in the previous revision are coded to another category in the new revision. To compensate for these shifts, comparability (also called bridge-coding) studies are undertaken to determine the extent of comparability between the two revisions. As a result of these studies, comparability ratios are developed that reflect the degree of comparability between the two revisions.

CRUDE BIRTH RATE—Number of live births per 1, 000 population.

CRUDE DEATH RATE—Number of deaths per 1,000 or 100,000 population.

D.W.I.—Driving While Impaired; the current legal limit for blood alcohol concentration is .08 for 21 and older, and .02 for younger than 21.

FERTILITY RATE—Number of live births per 1,000 women 15-44 years of age.

FOOD STAMP PROGRAM (FSP)—The FSP is an entitlement program based on Federal rules and mandates. The purpose of the FSP is to increase food purchasing power and to supplement the nutrition level of eligible households. In addition, the FSP provides a nutrition education program to eligible households. Eligibility for participation in the FSP is based on a gross income test at 130% of the Federal Poverty Guidelines for the size of the household. Eligibility is also based on a resource test; the household's liquid and nonliquid resources cannot exceed \$2000.00. Food stamp benefits are issued on a monthly basis through an electronic benefit transfer (EBT) system. The FSP has a change reporting and work requirement for adults in the household. Failure to report changes or comply with work requirements results in a reduction or closure of Food Stamp benefits. Under contract with another agency, the Department provides a free nutrition education program for Food Stamp applicants and recipients.

HOSPITAL DISCHARGE DATA—Data, such as diagnosis, gender, age, residence, on hospital inpatient discharges; collected by the NM Health Policy Commission from licensed, non-federal general and specialty hospitals.

INCIDENCE—Measure of the frequency at which new events/cases of a disease or condition occur in a specified population over a given time period.

INCIDENCE RATE—Number of new cases of disease/condition over a given time period (often, a year) that occur in a population at risk for that disease/condition (ergo number of new lung cancer cases per 100,000 persons in a year). An estimate of the risk of developing a disease/condition during a specified time period.

INFANT DEATH—Death of an infant under one year of age (neonatal + postneonatal).

INFANT MORTALITY RATE—Number of infant deaths per 1,000 (or 100,000) live births.

LIVE BIRTH—A birth that shows any sign of life after delivery.

LOW BIRTH WEIGHT—A live birth weighing less than 2500 grams (5-1/2 pounds).

MEDICAID—A federally and state-funded program to pay for medical and other services to eligible New Mexico citizens.

NEONATAL DEATH—Death of newborn under 28 days of age.

NEONATAL MORTALITY RATE—Number of neonatal deaths per 1,000 (or 100,000) live births.

NEURAL TUBE DEFECTS—Malformations of the spine and brain that arise in the first 29 days of pregnancy.

OCCURRENCE DATA—Data allocated by place where event occurred, regardless of the person's place of residence.

GLOSSARY CONTINUED

POPULATION ESTIMATE—Estimate of the number of persons for either a past or current time period.

POSTNEONATAL DEATH—Death of an infant of at least 28 days, but less than one year of age.

POSTNEONATAL MORTALITY RATE—Number of postneonatal deaths per 1,000 or (100,000) live births.

POVERTY—Poverty status is based on the 1964 Social Security Administration definition (modified in 1969 and 1980). At the core of this definition was the 1961 Department of Agriculture economy food plan, which arose from findings of a 1955 Department of Agriculture survey of food consumption. The 1955 findings showed that families of three or more spent about 1/3 of their income on food. Poverty status/level is tied to the cost of the economy food plan that is revised annually for cost of living as reflected in the Consumer Price Index.

PRENATAL CARE—1) Level of Prenatal Care – categorization of prenatal care services into no/low, middle and high levels of care based on the number of visits and month care began (modified Kessner index). 2) Adequacy of Prenatal Care Utilization Index – an alternative measure of prenatal care utilization that adjusts for some of the weaknesses of the trimester care began and the Kessner index (Kotelchuck index).

RESIDENCE DATA—Data allocated by place of residence of the decedent (deaths) or by place of residence of the child's mother (births).

SCHOOL ENROLLMENT—The count of high school students on the current roll on the 40th official school day, by school district.

SUBSTANTIATED—Through the course of the investigation, the social worker determined that the child(ren) who is the subject of the report have been determined to have been the victim of abuse and/or neglect. In other words the allegations are true.

TEMPORARY ASSISTANCE FOR NEEDY FAMILIES (TANF)—TANF is the State-administered, county-operated, monthly cash assistance program funded by a Federal block grant under Title IV-A of the Social Security Act. In July 1997, TANF replaced the previous entitlement program known as Aid to Families with Dependent Children (AFDC). The TANF program is not an entitlement program and has a 60-month term limit in New Mexico, which means that adults receiving assistance from the TANF block grant program can receive assistance for a 60 month maximum limit. TANF is available to both single and two-parent families. To receive TANF, there must be at least one dependent child (under age 18) in the home. TANF eligibility is based on a gross income test at 85% of the Federal Poverty Guideline for the size of the family. Eligibility is also based on a resource test; the family's liquid resources must be less than \$1500.00 and non liquid resources must be less than \$2000.00. Cash assistance is issued through an electronic benefit transfer system, similar to an ATM debit card. The TANF program has a mandatory change reporting and work requirement for adults and a mandatory school attendance requirement for dependent children. The adults in the family must cooperate with child support enforcement efforts on behalf of the dependent children. There are cash payment reductions for failure to comply with change reporting, work requirements, child support enforcement and school attendance.

TOTAL PERSONAL INCOME (TPI)—Includes the earnings (wages and salaries, other labor income, proprietors' income), dividends, interest, and rent, and transfer payments received by the residents of county. Data shown in thousands.

UNEMPLOYED—Based on monthly surveys, persons who did not work during a specified week but made specific efforts to find a job in the previous 4 weeks and were available for work. Also, persons who did not work, were available for work but were not actively seeking work because they (a) had been laid off and were waiting to be called back; or (b) were waiting to report to a job within 30 days. Does not include those no longer seeking work other than in situations (a) and (b).

UNEMPLOYMENT RATE—Number of persons unemployed as a percent of the civilian labor force.

UNSUBSTANTIATED—Investigation indicates that the investigator was unable to determine that the allegation(s) were true.

VERY LOW BIRTH WEIGHT—A live birth weighing less than 1500 grams (about 2 lbs. 5 oz.) at birth.

W.I.C.—Women, Infants and Children; a federally-funded, special supplemental nutrition program administered by the New Mexico Department of Health and tribal agencies. Its purpose is to provide high-nutrient density food to promote optimal health outcome during the perinatal period. Services include food assistance, nutrition education and referral to health services.