New Mexico Stroke Assessment Tool (NMSAT)

The patient may still be having a stroke even if all NMSAT criteria are not met. If all criteria are not met, and stroke is still suspected, treat the patient under stroke treatment and transport protocols. Notify the receiving facility that the patient is a RULE OUT STROKE.

DO NOT DELAY RAPID TRANSPORT of a suspected stroke patient. Obtain as much history and assessment as possible WHILE ENROUTE. If necessary, transport the patient with witness or a family member.

PATIENT NAME:	PATIENT AGE	·	M / F (circle one)
PROVIDERS NAME:	TOĐAYS DATI	E:	540
WITNESS NAME:		IE:	
EVENT HISTORY/ COMMENTS:			
		<u> </u>	700
****LAST TIME THE PATIENT WAS SEEN WI	THOUT SYMPTOMS	**** DATE: _	TIME:
SCREENING CRITERIA		1	TRUE FALSE
. Sudden onset of new onset of neurologic symptoms in last 12 hou	ırs,	1000	
. Blood Glucose between 60 and 400.		747	
. Symptoms were not immediately preceded by a seizure.	- 18 A	00	
. Focused assessment (IF 2 OR MORE SYMPTOMS PRESENT,	CHECK TRUE)	PRESENT	
Facial droop present on one side only? (ask the patient to smile)	j .		
Arm drift present on one side only? (ask the patient to hold their arms out in front of them and kee	p them level)		86
Leg Weakness present on one side only? (NEW ONSET difficu	dt or unable to walk)	·	
Speech deficit present? (i.e. slurred or inappropriate words or m	ute)	9	
operation presents (i.e. statted of mappiopriate words of in			

References: Los Angeles Prehaspital Stroke Screen, Cincinnati Prohospital Stroke Scale, Miami Emergency Neurologic Deficit (MEND) Prehospital Checklist