

Mental Health


June 2015 | Page 1 of 2

Why is mental health a public health issue?

- Mental health affects a person's ability to lead a **healthy, productive life**
- Globally and in the United States, mental disorders make up the **highest burden of all diseases** in middle- and high-income countries
- Mental disorders increase risk of **substance abuse and suicide**
- Mental disorders are associated with **diabetes, heart disease, cancer,** and other chronic physical conditions

How does New Mexico compare to other states?

- New Mexico and the United States have similar rates of mental illness for youth and adults
- However, the New Mexico suicide rate is 59% higher than the United States
 - ◇ *Suicide is the **2nd leading cause of death** for New Mexico residents 15 to 44 years old*
 - ◇ *Suicide accounts for **9.8% of all New Mexico Years of Potential Life Lost** under 65 years*
 - ◇ *New Mexico has the **5th highest suicide rate** in the United States*
 - ◇ *Suicide rates have been increasing in New Mexico and the United States since 2000*



Jessica Reno | Mental Health Epidemiologist | Jessica.Reno@state.nm.us

Mental Health

What are the risk factors for mental disorders?

- **Genes** 17-28% of risk can be accounted for by variations in common genes
- **Environment** People who experience traumatic childhood events are
 - ◇ 17x more likely to have behavioral problems
 - ◇ 5x more likely to have alcohol problems
 - ◇ 3x more likely to have job problems
 - ◇ 2.6x more likely to have depression
- **Substance Abuse** People with mental disorders are
 - ◇ 30x more likely to develop illicit drug dependence
 - ◇ 3x more likely to develop alcohol dependence
 - ◇ 2.3x as likely to develop nicotine dependence
- **Biology** Head injuries and brain abnormalities

How can we prevent mental illness and suicides at a state level?

- The evidence-based practices to reduce mental disorders in the general population are:
 - ◇ Mental health benefits with financial coverage and access to services, and
 - ◇ Multicomponent, healthcare system-level intervention that screens for depression and links primary care providers to mental health specialists
- Approximately **half** of suicide victims make a healthcare visit within 4 weeks of death
 - ◇ 25% visit specialty care
 - ◇ 21% visit primary care
 - ◇ 13% visit the emergency department
 - ◇ ...but only 24% are diagnosed with a mental disorder

**If you or someone you know is experiencing an emotional crisis, please call the
New Mexico Crisis and Access Line 24/7 at 1 (855) NMCRISIS (662-7474)**

Sources:

World Health Organization. The global burden of disease: 2004 update. Geneva, Switzerland 2008.

New Mexico Death Certificate Database, Bureau of Vital Records and Health Statistics, NMDOH.

National Center for Health Statistics, Centers for Disease Control and Prevention.

Guide to Community Preventive Services, Centers for Disease Control and Prevention.

Substance Abuse and Mental Health Services Administration (SAMHSA), Center for Behavioral Health Statistics and Quality. (September 4, 2014).

The NSDUH Report: Substance Use and Mental Health Estimates from the 2013 National Survey on Drug Use and Health: Overview of Findings. Rockville, MD.

Ronald C. Kessler et al., Lifetime Prevalence and Age-of-Onset Distributions of DSM-IV Disorders in the National Comorbidity Survey Replication, 62 Arch. Gen. Psychiatry 593, 595 (2005).

Ahedani, B K et al. Health Care Contacts in the Year Before Suicide Death. J Gen Intern Med 29(6):870-7.

Jessica Reno | Mental Health Epidemiologist | Jessica.Reno@state.nm.us

