Code:	Code:
NEW MEXICO DEPARTMENT OF I	NEW MEXICO////
Date of Issue:	Date of Issue:
Program:	Program:
Program Contact Information:	Program Contact Information:
Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A)	Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A)
This individual is trained and certified through the approved New Mexico Department of Health overdose prevention and naloxone administration course.	This individual is trained and certified through the approved New Mexico Department of Health overdose prevention and naloxone administration course.
For contact information and a list of programs and service times, please visit: www.nmhivguide.org	For contact information and a list of programs and service times, please visit: www.nmhivguide.org
Code:	Code:
NEW MEXICO DEPARTMENT OF	NEW MEXICO DEPARTMENT OF
Date of Issue:	Date of Issue:
	Program:
•	
Program Contact Information:	Program Contact Information:
Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A)	Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A)
This individual is trained and certified through the approved	This individual is trained and certified through the approved
New Mexico Department of Health overdose prevention and naloxone administration course.	New Mexico Department of Health overdose prevention and naloxone administration course.
For contact information and a list of programs and service times, please visit: www.nmhivquide.org	For contact information and a list of programs and service times, please visit: www.nmhivquide.org
Code:	Code:
NEW MEXICO DEPARTMENT OF	NEW MEXICO DEPARTMENTO:
Date of leaves	Date of Issue:
Date of Issue:	
Program:	Program:
Program Contact Information:	Program Contact Information:
Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A)	Nelevane description appropriation to legally sormy (NIMCA 24-22-4-A)
This individual is trained and certified through the approved New Mexico Department of Health overdose prevention and naloxone administration course.	Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A) This individual is trained and certified through the approved New Mexico Department of Health overdose prevention and naloxone administration course.
For contact information and a list of programs and service times, please visit: www.nmhivquide.org	For contact information and a list of programs and service times, please visit: www.nmhivguide.org
Code: NEW MEXICO DEPARTMENT OF	Code: NEW MEXICO DEPARTMENT OF
Date of Issue:	Date of Issue:
Program:	Program:
Program Contact Information:	Program Contact Information:
Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A)	Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A)
This individual is trained and certified through the approved	This individual is trained and certified through the approved
New Mexico Department of Health overdose prevention and naloxone administration course.	New Mexico Department of Health overdose prevention and naloxone administration course.
For contact information and a list of programs and service times, please visit: www.nmhivguide.org	For contact information and a list of programs and service times, please visit: www.nmhivguide.org
Code:	Code:
NEW MEXICO DEPARTMENT OF I J J J J J J J J J J J J J J J J J J	NEW MEXICO DEPARTMENTOR HEALTH
Date of Issue:	Date of Issue:
Program:	Program:
Program Contact Information:	Program Contact Information:
Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A) This individual is trained and certified through the approved New Mexico Department of Health overdose prevention and naloxone administration course.	Naloxone does not require a prescription to legally carry (NMSA 24-23-1-A) This individual is trained and certified through the approved New Mexico Department of Health overdose prevention and naloxone administration course.

For contact information and a list of programs and service times, please visit: $\underline{www.nmhivguide.org}$

For contact information and a list of programs and service times, please visit: www.nmhivguide.org

Naloxone...also called Narcan

- It blocks the effects of opiates.
- It takes effect in 3-5 minutes and lasts for 30-90 minutes.
- It may cause some withdrawal symptoms.
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Stay with the person, naloxone loses effect 30-90 minutes after use.


Using "all-in-one" device

- blister pack
- Place nozzle end into
- nostril 3. Press firmly on base of



- <u>Using device with separate atomizer*</u>

 1. Remove colored caps on medicine vial &
- syringe barrel Insert vial into barrel & gently turn until it stops
- Twist nasal atomizer onto barrel tip. It is ready to use



Press firmly on base of the vial, spraying half into nostril 6. Repeat in the other nostril

*If an atomizer is not available, slowly drip the naloxone under the tongue

Naloxone...also called Narcan

- It blocks the effects of opiates.
- It takes effect in 3-5 minutes and lasts for 30-90 minutes.
- It may cause some withdrawal symptoms
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Stay with the person, naloxone loses effect 30-90 minutes after use.

<u>Using "all-in-one" device</u> 1. Remove device from

- blister pack
- Place nozzle end into nostril
- 3. Press firmly on base of device,



spraying medication into

<u>Using device with separate atomizer*</u>

1. Remove colored caps on medicine vial &

- syringe barrel Insert vial into barrel & gently turn until it
- Twist nasal atomizer onto barrel tip. It is
- ready to use
 Place assembled naloxone atomizer into one nostril
- Press firmly on base of the vial, spraying half into nostril
- Repeat in the other nostril 6.

*If an atomizer is not available, slowly drip the naloxone under the tongue

Naloxone...also called Narcan

- It blocks the effects of opiates.
- It takes effect in 3-5 minutes and lasts for 30-90 minutes
- It may cause some withdrawal symptoms.
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Stay with the person, naloxone loses effect 30-90 minutes after use.

Using "all-in-one" device

- Remove device from blister pack
- Place nozzle end into nostril
- Press firmly on base of device. spraying medication into



<u>Using device with separate atomizer*</u> 1. Remove colored caps on medicine vial &

- syringe barrel Insert vial into barrel & gently turn until it
- Twist nasal atomizer onto barrel tip. It is
- 3. ready to use
- Place assembled naloxone atomizer into one nostril Press firmly on base of the vial, spraying half into nostril
- 6. Repeat in the other nostril

*If an atomizer is not available, slowly drip the naloxone under the tongue

Naloxone...also called Narcan

- It blocks the effects of opiates.
- It takes effect in 3-5 minutes and lasts for 30-90 minutes.
- It may cause some withdrawal symptoms.
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Stay with the person, naloxone loses effect 30-90 minutes after use.

Using "all-in-one" device 1. Remove device from

- blister pack
- Place nozzle end into nostril
- 3. device



spraying medication into





 Remove colored caps on medicine vial & syringe barrel 2. Insert vial into barrel & gently turn until it

Using device with separate atomizer

3 Twist nasal atomizer onto barrel tip. It is

ready to use

Place assembled naloxone atomizer into one nostril Press firmly on base of the vial, spraying half into nostril

6. Repeat in the other nostril

*If an atomizer is not available, slowly drip the naloxone under the tongue

Naloxone...also called Narcan

- It blocks the effects of opiates.
- It takes effect in 3-5 minutes and lasts for 30-90 minutes.
- It may cause some withdrawal symptoms.
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Stay with the person, naloxone loses effect 30-90 minutes after use.

Using "all-in-one" device 1. Remove device from

- blister pack Place nozzle end into
- nostril
 3. Press firmly on base of device.
 - spraying nostril
 - medication into

Using device with separate atomizer Remove colored caps on medicine vial &

- svringe barrel
- 2. Insert vial into barrel & gently turn until it stops 3.
- Twist nasal atomizer onto barrel tip. It is ready to use
- Place assembled naloxone atomizer into one nostril Press firmly on base of the vial, spraying half into nostril 6. Repeat in the other nostril
- *If an atomizer is not available, slowly drip the naloxone under the tongue

Naloxone...also called Narcan

- It blocks the effects of opiates.
- It takes effect in 3-5 minutes and lasts for 30-90 minutes.
- It may cause some withdrawal symptoms.
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Stay with the person, naloxone loses effect 30-90 minutes after use.

Using "all-in-one" device

- blister pack
- Place nozzle end into nostril
- 3. Press firmly on base of device,



spraying medication into nostril

- <u>Using device with separate atomizer*</u>

 1. Remove colored caps on medicine vial & syringe barrel
- 2. Insert vial into barrel & gently turn until it stops
- Twist nasal atomizer onto barrel tip. It is
- ready to use
 Place assembled naloxone atomizer into one nostril
- Press firmly on base of the vial, spraying half into nostril
- Repeat in the other nostril

*If an atomizer is not available, slowly drip the naloxone under the tongue

Naloxone. .also called Narcan

- It takes effect in 3-5 minutes and lasts for 30-90 minutes.

- <u>Using "all-in-one" device</u>

 1. Remove device from blister pack
- Place nozzle end into nostril
- 3. Press firmly on base of device, spraying medication into nostril

3.

Press firmly on base of the vial, spraying half into nostril 6. Repeat in the other nostril *If an atomizer is not available, slowly drip the naloxone under the tongue

Naloxone...also called Narcan

- It may cause some withdrawal symptoms.

Using "all-in-one" device

- Remove device from blister pack
- Place nozzle end into nostril



device. spraying medication into nostril

stops

- ready to use
- Press firmly on base of the vial, spraying half into nostril 5.

6. Repeat in the other nostril

- It blocks the effects of opiates.
- It may cause some withdrawal symptoms.

Stay with the person, naloxone loses effect 30-90 minutes after use.

- Using "all-in-one" device 1. Remove device from
- blister pack Place nozzle end into



3.

spraying medication into

- Using device with separate atomizer
- 3 Twist nasal atomizer onto barrel tip. It is
- ready to use Place assembled naloxone atomizer into one nostril
- 6. Repeat in the other nostril *If an atomizer is not available, slowly drip the naloxone under the tongue


- Naloxone...also called Narcan
- It takes effect in 3-5 minutes and lasts for 30-90 minutes.
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Using "all-in-one" device 1. Remove device from Using device with separate atomizer Remove colored caps on medicine vial & blister pack
 - stops 3. Twist nasal atomizer onto barrel tip. It is

 - 6. Repeat in the other nostril



- It blocks the effects of opiates.


- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Stay with the person, naloxone loses effect 30-90 minutes after use.



- Twist nasal atomizer onto barrel tip. It is

- It blocks the effects of opiates.
- It takes effect in 3-5 minutes and lasts for 30-90 minutes.
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.
- Stay with the person, naloxone loses effect 30-90 minutes after use.
 - syringe barrel
 - 3. Twist nasal atomizer onto barrel tip. It is
 - Place assembled paloxone atomizer into one postril

- Naloxone...also called Narcan
- It takes effect in 3-5 minutes and lasts for 30-90 minutes.
- If not breathing use Rescue Breathing (mouth to mouth) 3-4 cycles of 12 breaths a minute before giving a second dose of naloxone.



Press firmly on base of the vial, spraying half into nostril

It blocks the effects of opiates.

spraying

nostril

medication into

- It may cause some withdrawal symptoms.
 - Stay with the person, naloxone loses effect 30-90 minutes after use.
 - Place nozzle end into nostril Press firmly on base of device.
 - Place assembled naloxone atomizer into one nostril Press firmly on base of the vial, spraying half into nostril
 - *If an atomizer is not available, slowly drip the naloxone under the tongue

- svringe barrel 2. Insert vial into barrel & gently turn until it
- ready to use

- It may cause some withdrawal symptoms



- ready to use
 Place assembled naloxone atomizer into one nostril

- - <u>Using device with separate atomizer*</u>

 1. Remove colored caps on medicine vial &
 - 2. Insert vial into barrel & gently turn until it