

**Supplement**  
**Racial and Ethnic Health Disparities Report Card**  
(Explanation of indicators and sources of data)

**1. Prenatal Care – Late or No Care (Care Beginning After the 3<sup>rd</sup> month of Pregnancy or no Prenatal Care):**

Late or no prenatal care is the percent of total births to women who did not receive prenatal care at all or began prenatal care after the first trimester of pregnancy divided by the total number of live births (excluding the births with unknown prenatal care).

Race	2009-2011 per 100	Confidence Interval <sup>1</sup>
African-American/Black	33.0	28.2-37.8
American Indian/Alaska Native	45.3	43.1-47.5
Asian/Pacific Islander	25.1	21.1-29.2
Hispanic	32.5	31.5-33.4
White	25.8	24.7-27.0

**Source of data:**

*New Mexico data* - New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2003 -2011 birth files.

*United States data* – National Center for Health Statistics, CDC as reported in national vital statistics reports for final birth data for calendar years 2003-2009 (<http://www.cdc.gov/nchs/births.htm>).

**2. Infant Mortality:**

Infant mortality consists of deaths to infants before the first birthday and is the ratio of deaths divided by live births.

Race	2009-2011	Confidence Interval <sup>1</sup>
African-American/Black	14.0	9.1-20.7
American Indian/Alaska Native	6.3	5.0-7.9
Asian/Pacific Islander	6.4*	3.2-11.5
Hispanic	5.4	4.8-6.1
White	3.9	3.1-4.8

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

**Source of data:**

*New Mexico data* - New Mexico Department of Health, Bureau of Vital Records and Health Statistics, **Deaths**: 2003- 2011 death files; **Births**: 2003- 2011 birth file;

*United States data* - National Center for Health Statistics, CDC as reported in national vital statistics reports for final death data for calendar years 2003-preliminary 2010 (<http://www.cdc.gov/nchs/deaths.htm>.)

### 3. Teen Births Ages 15-17:

Teen births are the number of births to females ages 15-17 divided by the number of females ages 15-17.

Race	2009-2011 Rate (per 1,000)	Confidence Interval <sup>1</sup>
African-American/Black	22.8	18.3-27.4
American Indian/Alaska Native	30.1	27.7-32.5
Asian/Pacific Islander	8.6	5.1-12.2
Hispanic	42.2	40.6-43.8
White	12.1	11.1-13.2

**Source of data:**

**New Mexico -Births:** New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2003 – 2011 birth files;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011

**United States data:** National Center for Health Statistics, CDC as reported in national vital statistics reports for calendar year 2003-2010 (<http://www.cdc.gov/nchs/births.htm>).

### 4. Adults with Diabetes Not Receiving All Recommended Diabetes Preventive Services:

Adults with Diabetes not receiving all recommended diabetes preventive services are the percent of individuals with diabetes who indicated that they had not received all three recommended services during the past year. The recommended services are lab work (two A1C tests), a foot exam and a dilated eye exam. Rates are estimated percents of population based on sample response. Estimates based on less than 50 records cannot be presented.

Race	2011 Percent	Confidence Interval <sup>1</sup>
African-American/Black	*	*
American Indian/Alaska Native	49.4	38.2-60.6
Asian/Pacific Islander	*	*
Hispanic	59.3	52.7-65.5
White	55.7	49.4-61.7

\*Survey results based on less than 50 interviews may fluctuate greatly and so are not presented.

**Source of data:**

**New Mexico data:** New Mexico Behavioral Risk Factor Surveillance System 2011 and the 2008-2010, 2007-2009, 2006-2008, 2005-2007, 2004-2006, and 2003-2005 Combined data sets; New Mexico Department of Health, Epidemiology and Response Division, Survey Unit.

Please note that changes to the weighting system and addition of cell phone interview preclude combining 2011 data with previous years

**United States** - CDC Behavioral Risk Factor Surveillance System online database.  
<http://apps.nccd.cdc.gov/brfss/index.asp>

### 5. Diabetes Deaths

Diabetes deaths are deaths for which diabetes is listed as the underlying cause of death on the death certificate. The number of deaths is divided by the total population and age-adjusted to the 2000 United States standard population<sup>2</sup>.

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	38.2	26.2-50.2
American Indian/Alaska Native	64.0	56.0-72.1
Asian/Pacific Islander	22.4*	10.7-34.2
Hispanic	38.4	35.7-41.1
White	18.7	17.3-20.1

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

#### Source of data:

*New Mexico Data:* Deaths: New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2003-2011 death files;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011.

*United States Data:* Centers for Disease Control and Prevention, National Center for Health Statistics, compressed Mortality File for calendar years 2002-2009 CDC WONDER on-line database, (<http://wonder.cdc.gov/>).

#### 6. Obesity Among Adults:

Obesity among adults is the percent of adults whose self-reported weight and height indicate a body mass index (BMI) of 30 or greater. BMI is a measure of body fat based on height and weight and people with a BMI of 30 or greater are considered obese. The challenge of survey data is to take a sample of the population and to derive generalizations about the entire population. Rates used are estimated percents of population based on sample response.

Race	2011 Percent	Confidence Interval <sup>1</sup>
African-American/Black	23.9	15.9-34.1
American Indian/Alaska Native	35.3	30.8-40.2
Asian/Pacific Islander	5.3	2.1-12.3
Hispanic	29.9	27.7-32.1
White	22.6	21.0-24.3

#### Source of data:

*New Mexico Data:* Mexico Behavioral Risk Factor Surveillance System 2011 and 2008-2010, 2007-2009, 2006-2008, 2005-2007, 2004-2006, and 2003-2005 Combined data sets; - New Mexico Department of Health, Epidemiology and Response Division, Survey Unit;

Please note that changes to the weighting system and addition of cell phone interview preclude combining 2011 data with previous years

*United States data-* CDC Behavioral Risk Factor Surveillance System online database.  
<http://apps.nccd.cdc.gov/brfss/index.asp>

#### 7. Obesity Among Youth:

Obesity among youth is the percent of youth whose self-reported height and weight indicate a body mass index (BMI) greater than the 95<sup>th</sup> percentile of youth, based on national multi-year reference data. BMI is a measure of body fat based on height and weight.

Race	2011 Percent	Confidence Interval <sup>1</sup>
African-American/Black	12.0	8.5-16.5
American Indian/Alaska Native	21.3	16.6-26.8
Asian/Pacific Islander	14.6	10.1-20.5
Hispanic	11.3	9.7-13.2
White	9.5	8.1-11.2

**Source of data:**

*New Mexico data:* 2011 New Mexico Youth and Resiliency Survey, school district-level sample. (New Mexico Department of Health, Epidemiology & Response Division and New Mexico Public Education Department). Note, in order to have adequate numbers of African-Americans and Asian/Pacific Islanders for reporting purposes, data reported here are from a separate but simultaneously administered survey. For this reason, these estimates may differ slightly from other reported statewide estimates. (Please note that the YRRS is conducted every other year so this indicator will be updated on a biennial basis);

*United States data* - the CDC Youth Risk Behavior Survey Online database: Comprehensive result.

#### 8. Not Had Pneumonia Vaccination (Adults 65 +):

Not had pneumonia vaccination is the percent of adults ages 65 or older who indicate they have never received a Pneumococcal vaccination. Rates are estimated percents of population based on sample response.

Race	2011 Percent	Confidence Interval <sup>1</sup>
African-American/Black	*	*
American Indian/Alaska Native	34.3	22.9-47.9
Asian/Pacific Islander	*	*
Hispanic	35.5	30.9-40.5
White	26.7	24.2-29.4

\*Survey results based on less than 50 interviews may fluctuate greatly and so are not presented.

**Source of data**

*New Mexico data:* New Mexico Behavioral Risk Factor Surveillance System 2011 and 2008-2010, 2007-2009, 2006-2008, 2005-2007, 2004-2006, and 2003-2005 Combined data sets - New Mexico Department of Health, Epidemiology and Response Division., Survey Unit;

Please note that changes to the weighting system and addition of cell phone interview preclude combining 2011 data with previous years

*United States data:* CDC Behavioral Risk Factor Surveillance System online database  
<http://apps.nccd.cdc.gov/brfss/index.asp>

#### 9. Pneumonia and Influenza Deaths:

Pneumonia and Influenza deaths are deaths for which pneumonia or influenza is indicated as the underlying cause of death on the death certificate. The number of deaths is divided by the total population and age-adjusted to the 2000 United States standard population.<sup>2</sup>

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	7.3*	2.0-12.6
American Indian/Alaska Native	34.1	28.1-40.1
Asian/Pacific Islander	18.4*	7.0-29.8
Hispanic	15.4	13.6-17.1
White	13.6	12.4-14.8

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

**Source of data:**

*New Mexico Data: Deaths:* New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2002-2011 death files;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011.

*United States data* - Centers for Disease Control and Prevention, National Center for Health Statistics, Compressed Mortality File for calendar years 2002-2009, CDC WONDER on-line database, (<http://wonder.cdc.gov/deaths>).

## **10. Chlamydia (females aged 15-24):**

Chlamydia cases are females ages 15-24 diagnosed with Chlamydia divided by number of females age 15-24. Data by race excludes cases between 2009-2011 for which race was other or unknown.

Race	2009-2011 Rate
African-American/Black	2203.8
American Indian/Alaska Native	3296.1
Asian/Pacific Islander	193.6*
Hispanic	3582.8
White	1403.1

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

### **Source of data:**

**New Mexico data:** **Cases:** New Mexico Department of Health, Sexually Transmitted Disease Program.  
33.8% of cases have race/ethnicity reported as other or unknown.

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011

**United States data:** Centers for Disease Control and Prevention – 2010.

## **11. Hepatitis B:**

Hepatitis B cases Individuals diagnosed with acute/chronic Hepatitis B divided by the total population.

Race	2009-2011 Rate
African-American/Black	11.4
American Indian/Alaska Native	1.5*
Asian/Pacific Islander	122.6
Hispanic	2.5
White	3.3

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

### **Source of data:**

**New Mexico Cases:** New Mexico Department of Health, HIV & Hepatitis Epidemiology Program.

*Approximately 67 cases did not have race/ethnicity recorded.*

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011.

## **12. HIV/AIDS:**

HIV/AIDS cases are New Mexico residents who were diagnosed with HIV infection divided by the population.

Race	2009-2011 Rate
African-American/Black	17.7
American Indian/Alaska Native	11.0
Asian/Pacific Islander	1.7*
Hispanic	8.2
White	4.4

\*Rates denoted by an asterisk are based on less than 20 events during the three year period and may fluctuate greatly from year to year.

**Source of data:**

*New Mexico data:* **Cases:** New Mexico Department of Health, HIV & Hepatitis Epidemiology Program, data are for cases diagnosed among residents of New Mexico;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011.

*United States data:* CDC/HIV/AIDS – US data by gender is only available for adults and adolescents .

### 13. Pertussis (Whooping Cough)

Pertussis cases are the number of New Mexicans of any age diagnosed with Pertussis during the divided by the midyear population.

Race	2009-2011 Rate (per 100,000)
African-American/Black	3.0
American Indian/Alaska Native	7.4
Asian/Pacific Islander	5.6
Hispanic	9.8
White	5.5

\*Rates based on less than 20 events and may fluctuate greatly from year to year. 66 cases unknown/missing/multirace designation.

**Source of data:**

*New Mexico data:* Pertussis cases-New Mexico Department of Health, Infectious Disease Epidemiology, Epidemiology and Response Division;

**Population:** Bureau of Business and Economic Research, UNM 2003-2011;

*United States data:* Centers for Disease Control and Prevention.

### 14. Motor Vehicle Deaths:

Motor Vehicle Deaths are deaths for which the underlying cause of death is listed as motor vehicle accident on the death certificate, divided by the total population and age-adjusted to the 2000 United States standard population.

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	8.6*	3.7-13.4
American Indian/Alaska Native	26.2	22.1-30.4
Asian/Pacific Islander	8.9*	3.6-14.2
Hispanic	16.2	14.6-17.8
White	12.2	10.8-13.6

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

**Source of data:**

*New Mexico:* **Deaths:** New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2002- 2011 death files;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011.

*United States data -* Centers for Disease Control and Prevention, National Center for Health Statistics, Compressed Mortality File for calendar years 2002-2009, CDC WONDER on-line database, (<http://wonder.cdc.gov/deaths>).

### 15. Fall-Related Deaths (Adults 65+)

Fall-related deaths are deaths for which the underlying cause of death is listed as a fall on the death certificate and divided by the population age 65 and older.

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	23.8*	0.0-57.0
American Indian/Alaska Native	70.9	47.4-94.4
Asian/Pacific Islander	22.6*	0.0-53.9
Hispanic	74.4	63.8-85.0
White	103.6	94.7-112.4

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

#### Source of data:

**New Mexico data: Deaths:** New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2003-2011 death files;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011.

**United States data -** Centers for Disease Control and Prevention, National Center for Health Statistics, Compressed Mortality File for calendar years 2002-2009, CDC WONDER on-line database, (<http://wonder.cdc.gov/deaths>).

#### 16. Suicide:

Suicides are deaths for which the underlying cause of death is listed as suicide on the death certificate and divided by the total population and age-adjusted to the 2000 United States standard population.<sup>2</sup>

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	8.6	3.6-13.6
American Indian/Alaska Native	18.2	14.8-21.5
Asian/Pacific Islander	9.7*	4.2-15.2
Hispanic	14.3	12.8-15.8
White	24.6	22.7-26.6

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

#### Source of data:

**New Mexico data: Deaths:** New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2002-2011 death files;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011

**United States data -** Centers for Disease Control and Prevention, National Center for Health Statistics, Compressed Mortality File for calendar years 2002-2009, CDC WONDER on-line database, (<http://wonder.cdc.gov/deaths>).

#### 17. Youth Suicide:

Youth Suicide are deaths to youth ages 15-24 for which the underlying cause of death is listed as suicide on the death certificate and divided by the 15-24 population.

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	10.1*	0.0-21.6
American Indian/Alaska Native	33.0	22.7-43.4
Asian/Pacific Islander	21.4*	0.4-42.4
Hispanic	19.0	14.8-23.2
White	22.0	16.7-27.4

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

**Source of data:**

**New Mexico data: Deaths:** New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2002-2011 death files;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2009

**United States data - Centers for Disease Control and Prevention, National Center for Health Statistics, Compressed Mortality File for calendar years 2002-2009, CDC WONDER on-line database, (<http://wonder.cdc.gov/deaths>).**

**18. Homicide:**

Homicides are deaths for which the underlying cause of death is listed as homicide and divided by the total population. Death rates are age-adjusted to the 2000 United States standard population.<sup>2</sup>

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	14.4	8.3-20.4
American Indian/Alaska Native	15.4	12.2-18.6
Asian/Pacific Islander	2.2*	0.0-4.8
Hispanic	9.0	7.9-10.2
White	4.1	3.3-5.0

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

**Source of data:**

**New Mexico data: Deaths:** New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2002-2011 death files;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2009.

**United States data - Centers for Disease Control and Prevention, National Center for Health Statistics, Compressed Mortality File for calendar years 2002-2009, CDC WONDER on-line database, (<http://wonder.cdc.gov/deaths>).**

**19. Smoking Among Adults:**

Smoking among adults is the percent of adults who indicate they currently smoke. Rates are estimated percents of population based on sample response.

Race	2011 Percent	Confidence Interval <sup>1</sup>
African-American/Black	29.3	20.6-39.8
American Indian/Alaska Native	21.5	17.6-26.0
Asian/Pacific Islander	15.8	8.5-27.6
Hispanic	22.8	20.8-24.9
White	20.5	18.8-22.2

**Source of data:**

**New Mexico data:** New Mexico Behavioral Risk Factor Surveillance System 2011 and the 2008-2010, 2005-2007, 2004-2006, and 2003-2005 Combined data sets - New Mexico Department of Health, Epidemiology and Response Division.

Please note that changes to the weighting system and addition of cell phone interview preclude combining 2011 data with previous years

**United States data:** CDC Behavioral Risk Factor Surveillance System online database.  
<http://apps.nccd.cdc.gov/brfss/index.asp>

## **20. Drug-Overdose Deaths:**

Drug-overdose deaths are deaths caused by an overdose of illicit or prescription drugs. Deaths are then divided by the population and age-adjusted to the 2000 United States standard population.<sup>2</sup>

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	19.5	12.6-26.4
American Indian/Alaska Native	10.5	7.9-13.1
Asian/Pacific Islander	6.6*	2.0-11.3
Hispanic	24.9	23.0-26.9
White	26.0	23.9-28.0

\*Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

### **Source of Data:**

*New Mexico data:* Deaths: Vital Records and Health Statistics, 2002-2011 death files, data compiled by Substance Abuse Epidemiology Unit;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2011.

*United States data -* Centers for Disease Control and Prevention, National Center for Health Statistics, Compressed Mortality File for calendar years 2002-2009, CDC WONDER on-line database, (<http://wonder.cdc.gov/deaths>).

## **21. Alcohol Related Deaths:**

Deaths considered to be either directly or indirectly attributable to alcohol. Directly alcohol-attributable deaths are those for which any of the following are listed as the underlying cause of death on the death certificate: alcoholic psychoses, alcohol dependence syndrome, nondependent abuse of alcohol, alcoholic polyneuropathy, alcoholic cardiomyopathy, alcoholic gastritis, alcoholic fatty liver, acute alcoholic hepatitis, alcoholic cirrhosis of the liver, other alcoholic liver damage, excess blood alcohol level, and accidental poisoning by ethyl alcohol. Indirectly alcohol-attributable deaths (i.e., deaths for which alcohol is sometimes a contributing factor) include respiratory tuberculosis, diabetes mellitus, certain cancers, hypertension, hepatitis, unspecified liver cirrhosis, suicide, homicide, motor vehicle crashes and accidental injury. Alcohol-related deaths identified based on CDC Alcohol-Related Disease Impact (ARDI), version 2.0. Rates were calculated by dividing three-year total deaths by the mid-year population, and age-adjusting to the 2000 United States standard population.<sup>2</sup>

Race	2009-2011 Rate (per 100,000)	Confidence Interval <sup>1</sup>
African-American/Black	33.5	24.4-45.4
American Indian/Alaska Native	97.3	89.2-106.2
Asian/Pacific Islander	16.3*	8.8-31.8
Hispanic	56.9	53.8-60.2
White	38.7	36.5-41.1

\* Rates denoted by an asterisk are based on less than 20 events and may fluctuate greatly from year to year.

#### Source of data:

**New Mexico Data: Deaths:** New Mexico Department of Health, Bureau of Vital Records and Health Statistics, 2002-2010 death files, data compiled by Substance Abuse Epidemiology Unit;

**Population:** Bureau of Business and Economic Research, UNM, 2003-2010;

**United States data** for calendar years 2002-2009: Centers for Disease Control and Prevention, National Center for Health Statistics, Multiple Cause of Death Files;

---

<sup>1</sup> Confidence intervals are estimates of the range of values which will include the true population value.

<sup>2</sup> To account for differences in population age distributions, the age-adjusted death rate is used to compare relative mortality risks between groups and over time. The age-adjusted death rate is calculated by weighting the age-specific death rates and summing the products. Since 1999 the 2000 United States standard population has been used in determining the weights.